- 33 Rountree P, Clayton R. A contextual model of adolescent alcohol use across the rural-urban continuum. *Subst Use Misuse* 1999;**34**:495–519.
- 34 Maddox SJ, Prinz RJ. School bonding in children and adolescents: conceptualization, assessment, and associated variables. *Clin Child Fam Psych* 2003;**6**:31–49.
- 35 Aveyard P, Markham WA, Almond J, et al. The risk of smoking in relation to engagement with a school-based smoking intervention. *Soc Sci Med* 2003;**56**:869–82.
- 36 Fromme K, D'Amico E. Measuring adolescent alcohol outcome expectancies. *Psychol Addict Behav* 2000;**14**:206–12.
- 37 Kloepp M, Hendry L, Ingebrigtsen J, et al. Young people in 'drinking' societies? Norwegian, Scottish and Swedish adolescents' perceptions of alcohol use. *Health Educ Res* 2001;**16**:279–91.
- 38 Catalano R, Hawkins J. The social development model: a theory of antisocial behavior. In: Hawkins J, ed. *Delinquency and crime: current theories*. New York: Cambridge University Press, 1996:149–97.
- 39 Oetting E, Donnermeyer J. Primary socialization theory: the etiology of drug use and deviance. *Subst Use Misuse* 1998;**33**:995–1026.
- 40 Sutherland I, Willner P. Patterns of alcohol, cigarette and illicit drug use in English adolescents. *Addiction* 1998;**93**:1199–208.

THE JECH GALLERY

Children's exposure to tobacco

Although selling tobacco to children <16 years is not allowed, tobacco vendomats are widespread in Germany. By 2007 tobacco vendomats will be equipped with a mechanism that will make it impossible for children to get cigarettes. The technical solution chosen, however, does not guarantee the latter.

Young Germans mostly begin smoking between the ages of 11 and 16 years.¹ The prevalence of smoking among children and youth has risen since the 90s and is now around 20% of their total population.²

Figure 1 Tobacco vendomat beside a candy vendomat.

Figure 2 Tobacco vendomat less than 10 m away from a kindergarten.

Tobacco vendomats represent the main source of cigarettes for children and youth.³ On taking a short walk around the

Figure 3 Tobacco vendomat hung about 50 cm above the ground.

Figure 4 Tobacco vendomat close to a public playground.

quarter where I live in Hamburg, I found at least 10 tobacco vendomats placed near a school (in a radius of 200 m). Some were located right in front of public playgrounds, and others were close to the entrance of kindergartens, or a centre for youth counselling, some beside candy vendomats and some situated very low down. None of the vendomats displayed any warning about the risks of smoking, nor mentioned that the sale of tobacco to young people is prohibited.

One of the principles of health promotion is to facilitate healthy choices.⁴ However, it should be also be to render unhealthy choices more difficult... everywhere.

Marcial Velasco-Garrido

Correspondence to: M Velasco-Garrido, Department of Health Care Management, Technische Universität Berlin, EB 2 Strasse des 17, Juni 145, Berlin 10623, Germany; marcial.velasco@tu-berlin.de

References

- 1 Bundeszentrale für gesundheitliche Aufklärung. *Die Drogenaffinität Jugendlicher in der Bundesrepublik Deutschland 2001* [Drug-Affinity of Youth in the Federal Republic of Germany 2001]. Cologne: Bundeszentrale für gesundheitliche Aufklärung, 2001.
- 2 Schulze A, Lampert T. *Bundes-Gesundheitsurvey: Soziale Unterschiede im Rauchverhalten und in der Passivrauchbelastung in Deutschland* [Federal Health Survey: Social Inequalities in Smoking and Passive Smoking Burden in Germany]. Berlin: Robert-Koch-Institut, 2006.
- 3 Bayerisches Staatsministerium für Arbeit und Sozialordnung Familie, Frauen und Gesundheit. *Gesundheitsverhalten von Jugendlichen in Bayern: Ein Überblick über die repräsentativen Jugendgesundheitsstudien in Bayern von 1973 bis heute* [Health Attitudes of Youth in Bavaria: Overview of representative Youth Health Surveys in Bavaria from 1973 up to date]. München: Bayerisches Staatsministerium für Arbeit und Sozialordnung Familie, Frauen und Gesundheit, 1997.
- 4 World Health Organisation. *Ottawa Charter for health promotion*. Geneva: World Health Organisation, 1986.