

Scottish border), a geographically distinct area with a stable population of three million, and approximately 35 000 deliveries per year.

Participants: 13 738 individuals with at least one congenital anomaly (classified according to the European Surveillance of Congenital Anomalies, EUROCAT, guidelines) born between 01 January 1985 and 31 December 2003, of whom 10 951 were live born and 10 837 (99%) were traced for their survival status on 28 January 2008.

Main Outcome Measures: Congenital anomaly group and subtype specific survival rates at one week, four weeks, one year, five years, 10 years, 15 years and 20 years. Change in survival, and the number of terminations of pregnancy for fetal anomaly, with time.

Results: Survival among children with at least one congenital anomaly was 88.8% (95% CI 88.2 to 89.4) and 85.5 (95% CI 84.8 to 86.3) at one year and 20 years respectively. Twenty year survival among the most common anomaly groups was 89.5% (95% CI 88.4 to 90.6) for cardiovascular, 79.1% (95% CI 76.7 to 81.3) for chromosomal, 93.2% (95% CI 91.6 to 94.5) for urinary, 83.1% (95% CI 79.6 to 86.0) for digestive system, 97.7% (95% CI 96.1 to 98.7) for orofacial clefts, and 66.2% (95% CI 61.5 to 70.4) for nervous system anomalies. Survival varied considerably between subtypes, even within the same anomaly group. The proportion of terminations for fetal anomaly increased throughout the study period, and, together with year of birth, was an independent predictor of survival.

Conclusions: This study presents robust estimates of survival, uniquely to age 20 years, for a range of congenital anomaly groups and subtypes, some of which have been infrequently reported previously. This information will be valuable for families, genetic counsellors and other health care professionals when a congenital anomaly is detected, and will assist in planning for the future care needs of affected individuals.

067 MEASURING LOCOMOTOR DISABILITY: A NEW SCORING MECHANISM FOR SOME FAMILIAR ITEMS

S Muller, E Thomas, G Peat. *Arthritis Research Campaign National Primary Care Centre, Primary Care Sciences, Keele University, Keele, Staffordshire, UK*

doi:10.1136/jech.2009.096727p

Background: Locomotor disability is a critical measure for studying functional decline in later life and has featured in several major epidemiological studies. Yet measurement has mostly been confined to single items or ordinal scales rather than truly unidimensional measures with interval properties, thereby limiting the detection and interpretation of change in locomotor disability over time. Our aim was to combine items from the well-known SF-36 Physical Functioning Scale (PF-10) into a single score using Rasch analysis and test the properties of this new measure.

Methods: A health survey was sent to a random sample of 500 adults aged 50 years and over registered with a single general practice in North Staffordshire. The questionnaire contained the PF-10, from which five items (two relating to stair climbing and three to walking) were selected. Response dependency exists between the items, and so they were formed into two "super-items" (stair-climbing and walking). These items were assessed for their fit to the Rasch measurement model in terms of threshold ordering, unidimensionality, item and person fit, invariance along the latent trait and differential item functioning (DIF) by age-group (50–65, 66+ years) and gender. Rasch analysis was carried out in RUMM2020. The score was confirmed in three external datasets and tests carried out for its repeatability and validity.

Results: 372 people responded to the health survey and of these, 368 responded to at least one of the five PF-10 items. Both super-items displayed disordered thresholds that were corrected by combining adjacent response categories. There was no evidence

against unidimensionality, and item and person fit was good. There was no evidence against invariance along the latent trait and no signs of DIF by either age-group or gender. The items from the external datasets also met Rasch model specifications and the score was shown to be repeatable and valid.

Conclusions: Applications of measurement scales with confirmed interval level properties are advantageous for epidemiological studies using repeated measures over time. We have developed and externally validated a scoring method for deriving a brief self-reported locomotor disability scale with Rasch properties from commonly gathered items. This should allow the interval-level measurement of locomotor disability in general populations and so improve the quality of longitudinal studies in this area. The use of items from the PF-10 means that this improvement in quality is at no extra burden to the respondent or researcher, as in many cases, data can be analysed retrospectively.

068 PHYSICAL CAPABILITY AND SUBSEQUENT MORTALITY: A SYSTEMATIC REVIEW AND META-ANALYSIS OF PUBLISHED DATA

R Cooper, D Kuh, R Hardy, the FALCon study team. *MRC Unit for Lifelong Health and Ageing, Department of Epidemiology and Public Health, Royal Free and University College Medical School, London, UK*

doi:10.1136/jech.2009.096727p

Background: Evidence suggests that measures of physical capability may be predictive of subsequent disability, health and mortality, but the results from existing published studies have not been synthesised.

Objectives: To review published literature examining the associations between physical capability and subsequent all-cause mortality.

Design: Systematic review and meta-analysis.

Sources: Medline (from 1950 to 2008/05), EMBASE (from 1980 to 2008/02) and manual searches of the reference lists of eligible papers.

Included Studies: Any study examining the association between at least one of the specified measures of physical capability (grip strength, chair rises, standing balance, walking speed/time) and mortality in a population who were healthy and community-dwelling at baseline.

Results: Of 2240 unique records screened, 20 papers met the inclusion criteria of the review. An additional 3 papers were identified from reference lists. Within the 23 papers, there were 17 published analyses of grip strength, 9 of walking time/speed, 2 of standing balance and 1 of chair rises. The mean age of participants ranged from 36 to 80 y but the majority of study populations were aged 60 y and above. Of the papers investigating the association between grip strength and mortality, 6 presented Hazard Ratios for a unit change in grip strength with basic adjustments (total $n=10\,171$, deaths = 4267). The pooled Hazard Ratio for mortality per 5 kg increase in grip strength, from a random effects meta-analysis, was 0.90 (95% CI 0.87 to 0.93). Heterogeneity between studies ($I^2=65.2\%$) was largely explained by a single study which had weaker results. The pooled HR when this study was removed was 0.89 (0.86 to 0.92) ($I^2=25.7\%$). Other papers, where comparisons between categories of grip strength had been performed, also generally found weaker grip strength to be associated with higher rates of mortality but these analyses were too heterogeneous to be combined using meta-analysis. Due to differences in the ways in which walking speed had been measured and analysed it was not possible to perform meta-analyses of these data, but the results from all 9 studies showed that slower walking speed/longer walking time was significantly associated with increased mortality risk.

Conclusions: This review demonstrates that grip strength and walking time/speed are useful markers of subsequent mortality risk in community-dwelling populations. The underlying explanations

of these associations needs to be elucidated and further research is necessary to assess whether other measures of physical capability, including chair rises and balance, have similar predictive value.

Ageing

069 THE IMPACT OF CATARACT SURGERY ON HEALTH RELATED QUALITY OF LIFE AND TIME USE IN KENYA, BANGLADESH AND THE PHILIPPINES

¹S Polack, ²C Eusebio, ³W Mathenge, ⁴Z Wadud, ⁵AKM Mamunur, ¹A Foster, ¹H Kuper. ¹London School of Hygiene & Tropical Medicine, Keppel Street, London, UK; ²Cataract Foundation of the Philippines, Bacolod, Philippines; ³Rift Valley Provincial Hospital, Nakuru, Kenya; ⁴Child Sight Foundation, Dhaka, Bangladesh; ⁵CSS Rawm Hospital, Khulna, Bangladesh

doi:10.1136/jech.2009.096727q

Objective: To explore the impact of cataract surgery on health related quality of life (HRQoL) and time-use among adults aged ≥ 50 years Kenya, The Philippines and Bangladesh.

Methods: This was a multi-centre intervention study. Across the three countries at baseline, 651 population-based cases aged ≥ 50 years visually impaired from cataract (visual acuity in the better eye $< 6/24$), and 561 age-gender-cluster-matched controls with normal vision were identified. All participants were interviewed in their homes about vision related quality of life (WHO/PBD VF20), generic HRQoL (Euroqol) and time-use. Cases were offered free/subsidised cataract surgery. Approximately one year later participants were re-interviewed using the same questionnaires. Response rate at follow up was 84% for operated cases, and 80% for controls.

Results: At baseline, cases had substantially poorer vision specific and generic HRQoL compared to controls. Cases were also spent significantly less time on productive activities (paid and non-paid work) and more time in inactivity. Approximately one year after cataract surgery, mean vision specific and generic HRQoL improved ($p < 0.001$) to the level of controls with normal vision. Effect sizes for change in VRQoL were large (> 1.0) regardless of pre-operative VA, but were larger for those who had perception of light at baseline and for people who were operated in both eyes. Poor VA outcome from surgery was a constraint to achieving optimal post-operative VRQoL. At follow-up, operated cases were more likely to undertake and spent 1–2 hours more on productive activities compared to baseline ($p < 0.001$). Time spent in “inactivity” in Kenya and Bangladesh decreased by approximately 2 hours. Frequency of reported assistance with activities was more than halved in each setting among operated cases ($p < 0.001$).

Conclusion: Using three different outcome measures, this study demonstrated positive impacts of cataract surgery on the lives of older adults in three low-income settings, which has advocacy implications for blindness prevention programs. The observed increased time spent on productive activities, reduced time in inactivity and reduced assistance has positive implications for well-being and inclusion and supports arguments of economic benefit at the household level from cataract surgery.

070 POVERTY AND BLINDNESS: AN INTERVENTION STUDY TO ASSESS THE IMPACT OF CATARACT SURGERY ON POVERTY

¹H Kuper, ²S Polack, ³C Eusebio, ²W Mathenge, ⁴Z Wadud, ⁴R Mamun, ²A Foster. ¹Department of Epidemiology and Public Health, London School of Hygiene & Tropical Medicine, London, UK; ²Department of Infectious and Tropical Disease, London School of Hygiene & Tropical Medicine, London, UK; ³Cataract Foundation of the Philippines, Bacolod, the Philippines; ⁴Child Sight Foundation, Dhaka, Bangladesh

doi:10.1136/jech.2009.096727r

Objective: To assess the association between poverty and visually impairing cataract, and the impact of cataract surgery on alleviating poverty.

Design: Multicentre intervention study.

Setting: Centres in Kenya, Bangladesh and the Philippines. Baseline assessment was in 2005–2006, with follow-up one year later.

Participants: We recruited 596 cases and 481 controls, aged ≥ 50 years. Cases had visual acuity (VA) $\leq 3/36$ in the better eye due to cataract. Controls were age-sex-community-matched to cases with normal vision (VA $\geq 6/18$). Controls and most cases (65%) were recruited through population-based surveys, with additional cases (35%) identified through population case finding. 413 cases underwent cataract surgery. At follow-up, we re-interviewed 345 operated cases and 552 controls.

Interventions: Cataract surgery was offered to cases.

Main outcome measures: Poverty was measured through: (1) per capita household expenditure (PCE), (2) household assets and (3) household self-rated wealth.

Results: At baseline, operated cases were significantly more likely than controls to be in the lowest quartile of PCE in Kenya (OR 3.5, 95% CI 1.1 to 11.2), Bangladesh (3.0, 1.5 to 6.1) and the Philippines (4.4, 1.9 to 10.3), with a significant dose-response relationship across quartiles of PCE. The pattern was similar for assets and self-rated wealth. At follow-up, PCE had increased significantly among the operated cases in all countries, but not among the controls. Operated cases were no longer more likely than controls to be in the lowest category of PCE in Kenya and the Philippines, and the association in Bangladesh was weakened (2.0, 1.0 to 4.0). Operated cases remained poorer than controls in terms of assets and self rated wealth, although the association was weaker than at baseline.

Conclusion: The Cataract Impact Study provides evidence that cases with cataract are poorer than controls, and that operating on the cataract may help to alleviate poverty.

071 THE ASSOCIATION BETWEEN PSYCHOSOCIAL STATUS AND MORTALITY IN OLDER ADULTS: EVIDENCE FROM THE ENGLISH LONGITUDINAL STUDY OF AGEING

CC Craigs, AO House, J Hewison, RM West. Leeds Institute of Health Sciences, University of Leeds, Leeds, UK

doi:10.1136/jech.2009.096727s

Objectives: To evaluate the effect of psychosocial status on mortality risk among non-institutionalised older adults in England, controlling for selected demographic, health and lifestyle factors.

Participants: Cohort of 11 392 participants born on or before 29 February 1952 who responded to Wave 1 of the English Longitudinal Study of Ageing (ELSA), collected in 2002, and who participated in the Health Survey for England (HSE) survey in years 1998, 1999, or 2001.

Methods: The following three measures of psychosocial health, which were assessed at Wave 1, were included in the analysis: CASP-19, a measure of quality of life in early old age; the 12 item General Health Questionnaire (GHQ-12); and the Center for Epidemiologic Studies Depression Scale (CES-D). The number of negative statements agreed with or positive statements disagreed with were totalled separately for each of the three measures, resulting in scores in the range 0 to 19 for CASP-19, 0 to 12 for GHQ-12, and 0 to 8 for CES-D. Mortality status up to December 2006, as reported by ELSA in the Index file, was obtained from the Office for National Statistics. Logistic regression modelling was performed separately for each of the three measures, controlling for the following variables assessed at Wave 1: age, sex, marital status, highest educational qualification, smoking status, alcohol consumption, and self reported long-standing illness, disability and infirmity. The analysis was then replicated with a latent construct measured by CASP-19, GHQ-12, and CES-D.

Results: A total of 703 participants had died up to December 2006. The mean scores for participants who were identified as alive