

SP6-38 TOBACCO SMOKE AND THE RISK OF PERTHES' DISEASE IN SOUTH WEST INDIA: A CASE-CONTROL STUDY

doi:10.1136/jech.2011.142976q.9

¹A B Daniel,* ¹H Shah, ¹A Kamath, ²V Guddettu, ¹B Joseph. ¹Kasturba Medical College, Manipal, Karnataka, India; ²Department of Statistics, Manipal University, Manipal, Karnataka, India

Introduction Legg-Calvé-Perthes' disease is an idiopathic avascular necrosis of the femoral head that arises in childhood. There have been many theories on its aetiology, which include the role of genetic factors, environmental factors, poverty, repetitive trauma and abnormalities of thrombosis and fibrinolysis. Recent theories on the aetiology of Perthes' disease have focused on passive smoking as a risk factor. Previous studies in southwest India have determined that the prevalence of Perthes' disease is very high. The prevalence of tobacco use in the area is not higher than in other regions of the country. Based on these observations, the association between passive smoking and Perthes' disease was re-examined.

Methods Cases were children aged <12 years at diagnosis of Perthes' disease. Children presenting to the hospital for other orthopaedic complaints formed the controls. The 128 eligible cases were frequency matched on age and sex with controls, with 3 controls per case. Conditional logistic regression was employed to evaluate the association between the exposures and risk of Perthes' disease.

Results Risk factors for Perthes' disease were having a family member who smokes indoors (adjusted OR 2.07) and indoor use of a wood stove (adjusted OR 2.56).

Conclusion This study adds further evidence for the association between tobacco smoke and Perthes' disease. For the first time the association between wood smoke and Perthes' has been demonstrated.

SP6-39 OVERCOMING BARRIERS OF POOR PERINATAL CARE SERVICES IN URBAN SLUMS: POSSIBLE ROLE OF SOCIAL MOBILISATION NETWORKS

doi:10.1136/jech.2011.142976q.10

S Mehnaz,* Z Khan, A Ansari. Jawaharlal Nehru Medical College, Aligarh, Uttar Pradesh, India

Background Making perinatal care accessible to women in marginalised periurban areas poses a public health problem. Many barriers to utilisation of perinatal care can be overcome by social mobilisation and capacity building at the community level.

Objectives To determine the existing perinatal practices in two urban slums and to identify barriers to utilisation of health services by mothers.

Study design This is a cross-sectional descriptive study.

Setting and Participants The high-risk periurban areas of Nabi Nagar and Mehfooznagar, Aligarh has a population of 87 000 living in 10 480 households. Mothers delivering babies in September 2010 were identified. A total of 202 mothers were interviewed at home. Current perinatal practices and reasons for health services were the topics of enquiry. Statistical Analysis done by SPSS 18.

Results Analysis revealed that 80.4% of mothers had received antenatal care. However, this did not translate into safe delivery practices. Reasons for preferring home deliveries were mostly tradition (41.9%) or related to economics (30.7%). A total of 56% of the deliveries were conducted in the squatting position and in 25% of the cases, umbilical cord was cut using the edge of a broken cup. Inappropriate early neonatal feeding practices were common. Pre-lacteal feeds were given to nearly 50% of the babies and feeding was delayed beyond 24 h in 8% of the cases.

Conclusion Barriers to utilisation of available services leads to hazardous perinatal practices in urban slums.

SP6-40 DEATH AND HOSPITALISATION HAVE DIFFERENT SEASONALITY

doi:10.1136/jech.2011.142976q.11

L T Barnard,* M Baker, S Hales, P Howden-Chapman. University of Otago, Wellington, New Zealand

Introduction Seasonality of disease has been long recognised in epidemiology. More recently, researchers have established standard ways of measuring seasonality. In temperate countries there has been a particular interest in excess winter mortality (EWM), and, to a lesser degree, excess winter hospitalisations (EWH). Understanding the aetiology of seasonality is important for identifying interventions and potential future climate change effects.

Method We measured EWM and EWH in 60–95 year olds between 2000 and 2008, by ICD-10 chapter, and the contribution of each chapter to all-cause winter excess. We then compared indices and percentage contributions for mortality to those for hospitalisation.

Results Indices do not lie in the same direction for all ICD-10 chapters. Excesses lay within the same range only for respiratory deaths and hospitalisations. Circulatory illness showed an excess for both hospitalisation and death, but the excess was higher for mortality than for morbidity. However, neoplasms, digestive diseases, and genitourinary diseases all showed a winter mortality excess but a non-winter hospitalisation index. Other chapters also lacked a relationship between indices for hospitalisation and death. Similarly, there was a difference in contribution to winter excess by chapter.

Conclusion The comparison of EWM and EWH indicates that caution may be needed in extrapolating causal results from one to the other. As their disease and excess distribution is different, factors which contribute to EWH may play a different role in EWM, and vice versa. Research that can identify the reasons for these differences would improve our understanding of the mechanisms causing disease seasonality.

SP6-41 DESCRIPTIVE EPIDEMIOLOGY OF MORTALITY AMONG THE NEW ZEALAND EXPEDITIONARY FORCES IN WORLD WAR I

doi:10.1136/jech.2011.142976q.12

J A Summers,* N Wilson, M G Baker. University of Otago, Wellington, New Zealand

Introduction The military personnel in the New Zealand (NZ) Expeditionary Forces (NZEF) during World War I experienced high mortality (18 000+ deaths). There is little research on these deaths so we aimed to provide a more detailed historical and epidemiological account of this mortality burden.

Methods Various NZEF datasets and sources were used to examine mortality patterns and the robustness of the available data with historical accounts. Extensive coding work was required to allow epidemiological analyses.

Results The majority of NZEF deaths (80.9%) were a direct result of being killed in battle or being wounded. Deaths from disease also represented a substantial NZEF loss of life (at 17.9% of all deaths). The majority of these disease deaths were from pandemic influenza but other outbreaks occurred (eg, dysentery, measles). Māori (indigenous New Zealanders) and Pacific peoples in the NZEF experienced significantly higher mortality rates from disease compared to European/Other NZEF. Healthcare workers experienced significantly lower mortality from all causes, including disease, compared to other military units. This difference is possibly the result of less frontline presence and perhaps prior immunity from occupationally-related exposure to infectious agents.

Conclusions This study found substantial variation in mortality rates, by time, place and person. Both military factors and a

pandemic played important roles in the mortality burden. The datasets based on archival sources provide a rich foundation for both historical and epidemiological research of the mortality burden among these military personnel during World War I.

SP6-42 PHYSICAL ACTIVITY IS ASSOCIATED WITH REDUCED MORTALITY IN A RURAL SOUTH INDIAN COHORT

doi:10.1136/jech.2011.142976q.13

^{1,2}V Raman Kutty,* ²S Safraj, ^{2,3}K Vijayakumar, ³T S Anish, ²K Ajayan. ¹Achutha Menon Centre, SCTIMST, Trivandrum, Kerala state, India; ²Health Action by People, Trivandrum, Kerala state, India; ³Department of Community Medicine, Medical College, Trivandrum, Trivandrum, Kerala state, India

Introduction Physical activity is now recognised as an important determinant of health status. Thus it is important to explore its relation to mortality in transitional populations.

Methods We examined the association between self reported physical activity and mortality in a rural adult cohort numbering over 70 000 in southern India in the first 5 years of observation, from 2001 to 2006. The scores assigned under occupational and non-occupational recreational categories based on previous studies, were added to arrive at a total activity score. Based on this, we categorised subjects into those with mild, moderate, and heavy physical activity.

Results There is an inverse relationship between death rate and physical activity, with a much greater effect in women, true for both all cause and cardiovascular mortality. The effect is more pronounced in smoking males compared to non-smoking males. The protection offered by physical activity is also more evident in subjects aged 40–65 years. Heavy physical activity does not seem to confer proportionately greater protection as compared to moderate activity.

Conclusion Our observations confirm published reports on the protective effect of physical activity on health. Our data support the possibility that subjects with poor physical activity have more chronic morbidity, which underlies higher mortality. Clear cut relationships would emerge only with longer term follow-up of the cohort with physical activity profiles being captured at regular intervals.

SP6-43 ETHNIC GRADIENT IN MORTALITY AMONG THE NEW ZEALAND MILITARY PERSONNEL IN WORLD WAR I

doi:10.1136/jech.2011.142976q.14

J A Summers,* N Wilson, M G Baker. University of Otago, Wellington, New Zealand

Introduction The ethnic distribution in mortality rates for New Zealand (NZ) military personnel in World War I has never been described.

Methods Mortality data for military personnel in the NZ Expeditionary Forces (NZE) were obtained from an electronic dataset (Roll-of-Honour) covering all deaths in these personnel during World War I and the immediate post-war period (1914–1923). All NZE were allocated (using a classification system based on names and background) into the following ethnic groups: European/Other, Māori (indigenous New Zealanders) and Pacific peoples.

Results The highest all-cause mortality (deaths from enemy action, wounds, disease, and other) was experienced by Māori personnel. The majority of deaths among European/Other and Māori personnel were a result of being killed in action. Pacific personnel experienced the lowest all-cause mortality rate. Mortality rates from disease varied greatly by ethnic group, with Māori and Pacific personnel both experiencing higher rates than European/Other personnel.

When deaths from the 1918 to 1919 influenza pandemic are excluded from the analysis, Māori and Pacific personnel still had significantly higher mortality rates from disease compared to European/Other personnel (rate ratio 3.92, 95% CI 3.14 to 4.91 and rate ratio 3.45, 95% CI 2.42 to 4.91).

Conclusions We document for the first time higher all-cause mortality burden among Māori compared with European/Other for NZ's World War I effort. Also identified was the large variation in disease mortality, with Māori and Pacific personnel suffering to a disproportionately greater extent. These historical results serve as a reminder that tackling health inequalities is a long-term commitment that requires ongoing public health attention.

SP6-44 "AN APPROPRIATE APPROACH FOR RCH SERVICES DELIVERY THROUGH INCENTIVE BASED FEMALE ACTIVIST (ASHA) FOR RURAL AREA IN KANPUR DISTRICT, UTTAR PRADESH, INDIA"

doi:10.1136/jech.2011.142976q.15

S C Saxena,* A Singh, V K Srivastava, D S Martolia, P Varma, R P Sharma. GSVM Medical College, Kanpur University, Kanpur, Uttar Pradesh, India

Introduction In India, in the year 2005, the concept of Accredited Social Health Activist (ASHA)—incentive based female volunteer was implemented under NRHM with an objective that these female volunteers in co-operation with already existing other healthcare functionaries will facilitate the proper implementation of MCH programmes having an impact in the reduction of infant mortality rate and maternal mortality rate.

Methodology All the 135 female health volunteers of randomly selected primary health centres of Kanpur were studied by pre-designed and pretested questionnaire.

Results Predominantly the selected female health volunteers belonged to age group of 30–35 years (45.2%), Hindu by religion (99.3%), OBC caste (54.1%), educated upto middle standard (57.8%) and from Social Class IV (68.9%). About 2/3rd these volunteers had received the required two rounds of training but anyhow the remaining were partially trained. All the female health volunteers performed 100% activities regarding antenatal, intranatal and postnatal care and to lesser extent on adolescent's education about RTIs/STIs as well as supplementary nutrition. For improving their performance, all the health volunteers suggested about making their jobs permanent with fixed salary. Nearly half to 3/4th of them were in favour of providing transport as well as adequate training and mobile phone facility.

SP6-45 SCREENING FOR GENITAL CHLAMYDIA FOR YOUNG PEOPLE IN COMMUNITY PHARMACIES: A META-ANALYSIS OF CHLAMYDIA PREVALENCE

doi:10.1136/jech.2011.142976q.16

M Z Kapadia,* P Warner, K Fairhurst. Centre for population Health sciences, University of Edinburgh, Edinburgh, Scotland, UK

Background With traditional specialist sexual-health services overstretched, community pharmacies have been deployed to provide chlamydia testing. We have undertaken a systematic-review to estimate prevalence of chlamydia infection among those screened for chlamydia in community pharmacies. We also reviewed the feasibility/acceptability of this service.

Methods Systematic searches were conducted in electronic databases and grey literature was solicited from experts. Data were extracted on study population, sample size and prevalence of chlamydia to report pooled proportion of chlamydia infection using random effect model.