



OPEN ACCESS

Advice given by NHS Direct in Wales: do deprived patients get more urgent decisions? Study of routine data

Julie Peconi,¹ Steven Macey,² Sarah Rodgers,¹ Ian Russell,¹ Helen Snooks,¹ Alan Watkins¹

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/jech-2017-208978>).

¹Swansea University Medical School, Institute of Life Sciences 2 (ILS2), Swansea University, Swansea, UK

²Action on Smoking and Health (ASH) Wales, Cardiff, UK

Correspondence to

Julie Peconi, Institute of Life Sciences 2 (ILS 2), Medical School, Swansea University, Singleton Park, Swansea, SA2 8PP, Wales, UK; j.peconi@swansea.ac.uk

Received 20 January 2017

Revised 11 April 2017

Accepted 8 June 2017

ABSTRACT

Background In the UK, National Health Service Direct Wales (NHSDW) uses computerised decision support software to advise patients on appropriate care. However, the effect of deprivation on the advice given is not known. We aimed to estimate the effect of deprivation on advice given by nurses in NHSDW adjusting for confounding variables.

Methods We included 400 000 calls to NHSDW between January 2002 and June 2004. We used logistic regression to model the effect of deprivation on advice given by nurses in response to calls seeking advice or information. We analysed two outcomes: receiving advice to phone 999 emergency care rather than to seek other care and receiving advice to seek care face to face rather than self-care.

Results After adjustment for covariates, an increase in deprivation from one-fifth of the distribution to the next fifth increased by 13% the probability that those calling for advice rather than information received advice to phone 999 (OR 1.127; 95% CI from 1.113 to 1.143). Deprivation increased the corresponding probability of being advised to seek care face to face rather than self-care by 5% (OR 1.049; 95% CI from 1.041 to 1.058) within advice calls and by 3% (OR 1.034; 95% CI from 1.022 to 1.047) within information calls.

Conclusions Deprivation increased the chance of receiving more urgent advice, particularly advice to call 999. While our dataset may underestimate the 'need' of deprived patients, it yields no evidence of major inequity in advice given to these patients.

INTRODUCTION

One of the founding principles of the National Health Service (NHS) in the UK was equality of access to, and provision of, healthcare.^{1,2} Yet, in the UK and internationally, inequalities in health persist with people living in economically deprived areas known to have poorer health, including higher levels of depression³ and poorer physical function.⁴ People living in deprived areas are also less likely to have access to good quality medical care than those in more affluent areas, and consultations with General Practitioners (GPs) and other health practitioners may be less clinically effective.⁵ Evidence from providers of healthcare out of hours suggests that those from more deprived backgrounds are recommended more urgent care.^{6–8}

In Wales, the national, nurse-led telephone advice and information line NHS Direct Wales

(NHSDW) is in theory well placed to help those at socioeconomic disadvantage. For the cost of a local phone call, the service aims to provide accessible, standardised advice and information. Similar services existed in England (NHS Direct) and in Scotland (NHS 24) and now operate through '111' as a simpler number to ease entry into the complex emergency care system.⁹ In Wales, '111' is currently being trialled as a method of linking NHSDW and GP out-of-hours services.¹⁰

In the NHS Direct (NHSD) and 111 services, nurses or trained advisors generally use computerised decision support software (CDSS) to advise callers on the most appropriate form of healthcare or how to treat their symptoms themselves, working through a series of questions and answers to a decision. Although they can override this decision, the aim of this software is to give consistent advice in similar circumstances independent of patient or nurse characteristics.

However, there is evidence about variable provision in NHSD: nurses with over 20 years' experience were more likely to advise callers to care for themselves,¹¹ and Registered Sick Children's Nurses were more likely to refer children with fever or rash to routine GP appointments.¹² However, both studies lacked evidence about the influence of patient's characteristics on outcomes. Patient's deprivation has differentially affected the use of GP services that provide telephone advice out of hours, with both the likelihood of being subsequently seen by a GP falling with increasing deprivation^{13,14} to slightly increasing for those in the most deprived areas.¹⁵ However, once advised to see a GP, those in deprived areas were all more likely to receive home visits.^{13–15} We know of no study reporting the effect of patient deprivation on advice given by NHSD nurses or 111 call advisors. This paper therefore aims to describe how deprivation affects advice given by NHSDW controlling for other variables that may affect this advice.

METHODS

Time and place

Following approval by The South East Wales Local Research Ethics Committee in September 2004, we obtained anonymous data on all 615 739 calls to NHSDW originating from Wales between January 2002 and June 2004. Before receiving the data, an NHSDW analyst linked each patient's postcode to the corresponding Welsh Index of Multiple



CrossMark

To cite: Peconi J, Macey S, Rodgers S, et al. *J Epidemiol Community Health* Published Online First: [please include Day Month Year]. doi:10.1136/jech-2017-208978

Research report

Table 1 Codes used in logistic regression analysis

Study variable	Reference	Explanation
Advice to seek emergency care	All other care	999 call or emergency care
Advice to seek care face to face	Self-care	Care face to face (including emergency care)
Gender	Male	Female
Main symptom (from International Classification of Primary Care-2)	Not digestive	Digestive
Relationship of patient to caller	Surrogate caller	Self
Other ethnic background	White or unknown	Other
Ethnicity known	Unknown	Known (white or other)
Call occurred on Sunday	All other days	Sunday
Call occurred on Monday	All other days	Monday
Call occurred on Tuesday	All other days	Tuesday
Call occurred on Wednesday	All other days	Wednesday
Call occurred on Thursday	All other days	Thursday
Call occurred on Friday	All other days	Friday
Call occurred on Saturday	all other days	Saturday
Deprivation fifth	Least deprived	Second least deprived (2=3rd most deprived; 3=2nd most deprived; 4 = most deprived)

Deprivation (WIMD)¹⁶ and ward name before removing the postcode from the dataset, thus making the data anonymous. We excluded duplicate calls, calls transferred from Emergency Departments (EDs) or GP out-of-hour services and calls without deprivation scores. Initial exploration revealed that most calls from Flintshire and Rossett in Wrexham were handled by NHSD in England; therefore we excluded all calls from this area. This left 409 611 calls for analysis.

Data

NHSDW provided data on date (and thus day of call) and its type—whether for advice or information, patient's age, gender, ethnicity, symptom and relationship to caller, and the advice of the NHSDW nurse advisor. We did not receive data on time of call or duration. We coded patient's symptoms according to the International Classification of Primary Care-2 (ICPC-2).¹⁷ We supplemented call data with variables available only at ward level, notably deprivation score, distance to nearest ED and population density.

Table 1 shows how we defined our variables for logistic regression analysis. As preliminary data exploration showed digestive symptoms were most frequent, we grouped the rest to simplify analysis. Similarly, we coded the relationship of caller to the patient as self or surrogate for calls on behalf of someone else. As data on patient's ethnicity were available for only the final year, we created two dummy variables consistent with categories in the 2001 Census: 'white or unknown ethnicity' versus 'any other ethnicity' and 'known' (white or other) versus 'unknown'.

The original dataset contained 244 different types of advice. We were able to reduce these down to 28 using NHSDW's algorithm (see online Supplementary Appendix 1). This algorithm specified the grouping of similar advice into categories, for example, for the advice 'accident and emergency' or 'casualty' the algorithm grouped these into one advice: ED. We then recoded these categories into the six ordered categories used to evaluate NHSD in England¹⁸ and in Wales.¹⁹ These categories are based on resource use with the most expensive services at the top. Using this hierarchy, we ranked advice by urgency as follows: from (1) 999 call, (2) ED or other self-referral to hospital, (3) GP or dentist within 4 hours (labelled as 'emergency'), (4) GP or dentist less

urgently, (5) other healthcare and (6) self-care (the least expensive). We added an additional label as '(7) not assessed' to calls with no specific advice, including calls in which the nurse could not contact the caller again after several attempts. More than 50 000 early calls used a previous version of the NHSDW system that recorded up to four different categories of advice per call. To include these calls in analysis, one of us (JP) assigned each to its highest level of advice; for example, a call yielding advice to 'contact GP' and undertake 'self-care' in the mean time received a final classification of 'contact GP'.

Outcome variables

We treated calls which had been coded as 'not assessed' as self-care as they had not received any other advice from NHSDW. As advice could thus take one of the six forms, we summarised it by two binary variables: first whether the patient received advice to phone 999 versus any other care (contact hospital, GP, dentist, other healthcare or care for oneself) and second whether the advice was to contact any healthcare professional (care face to face) versus self-care. Thus, calls that received advice to phone 999 were always in the more urgent category. Following a previous study,¹¹ we chose these variables to represent the riskiest decisions for the nurse.

Ward-level variables

Our main explanatory variable was the summary score of the WIMD, the deprivation index used in Wales during data collection.¹⁶ To each call with a defined postcode (each of which covered an average of 18 residents), the NHSDW data analyst assigned the WIMD score for the corresponding electoral ward (with an average population of 3300). We then assigned each ward to its 'deprivation fifth' within the full range of deprivation scores. We estimated the distance from each ward geographical centroid to the nearest ED^{20 21} and estimated population density from the 2001 Census information for the 2003 administrative boundaries, which we converted to 1998 wards by Geoconvert.²²

As climatic variables like temperature^{23 24} and pollutants²⁵ affect patient's health, we considered including the average of the maximum and minimum monthly temperatures and air

Table 2 Characteristics of patients and their calls (n=409 611)

NHSDW call variable	n	%
Call type		
For advice	281 223	68.7
For information only	128 388	31.3
Day on which call occurred		
Sunday	66 297	16.2
Monday	61 502	15.0
Tuesday	56 341	13.8
Wednesday	55 863	13.6
Thursday	55 488	13.5
Friday	52 836	12.9
Saturday	61 284	15.0
Relationship of caller to patient		
Self	237 356	58.0
Surrogate	172 064	42.0
Not recorded	191	<0.1
Gender		
Male	155 279	38.0
Female	253 843	62.0
Not recorded	489	0.12
Ethnicity		
White background	3929	1.0
Any other background	180 308	44.0
Not recorded (mainly before July 2003)	225 374	55.0
Symptom (from ICPC-2)		
Digestive	67 190	16.4
General and unspecified	32 262	7.9
Skin	30 304	7.4
Musculoskeletal	27 982	6.8
Respiratory	27 325	6.7
Neurological	21 260	5.2
Female genital	6929	1.7
Eye	6390	1.6
Ear	6410	1.6
Psychological	6106	1.5
Urological	5964	1.5
Pregnancy and childbearing	4266	1.0
Cardiovascular	2620	0.6
Male genital	2387	0.6
Not recorded (mainly information calls)	162 216	39.6
Advice given		
(1) 999 or ambulance	12 791	3.1
(2) ED or other hospital	29 865	7.3
(3) Emergency GP or dentist	89 902	21.9
(4) Other GP or dentist	82 149	20.1
(5) Other	27 131	6.6
(6) Self-care	154 584	37.7
(7) Not assessed	13 189	3.2
Deprivation (from WIMD)		
Least deprived fifth	83 071	20.3
Second least deprived fifth	64 652	15.8
Third least deprived fifth	74 167	18.1
Fourth least deprived fifth	85 024	20.8

Continued

Table 2 Continued

NHSDW call variable	n	%
Most deprived	102 697	25.1

ED, emergency department; GP, general practitioner; ICPC-2, International Classification of Primary Care-2; NHSDW, National Health Service Direct Wales; WIMD, Welsh Index of Multiple Deprivation.

quality measures including the pollutants ozone, particulate matter 10, sulfur dioxide and nitrogen dioxide for each ward. Unfortunately, the paucity of weather stations (n=24) and air quality measuring stations (n=7) in Wales and on the border reduced the value of these data in initial analysis, so we excluded them from final analysis.

Statistical methods and sensitivity analyses

As calls made for advice differ in purpose and practice from calls only for information, we analysed these types of call separately using analysis of variance and χ^2 tests. Both also yielded two separate models: for the likelihood of receiving advice to call 999 over any other advice and for receiving face-to-face care (including emergency care) over self-care. We undertook three logistic regressions for each combination of call type and care model: first, we entered all variables except day of the week and deprivation, then we added weekday and finally we entered 'deprivation fifth' as an ordinal variable since that is simpler but less discriminatory than as a continuous variable. By adding deprivation to the statistical model at the final step, we were able to estimate its true contribution after accounting for known potential confounding variables.

For the majority of NHSDW variables, missing data were fewer than 1% with some exceptions. NHSDW collected data on ethnicity only for the final year. As expected, the majority (96.9%) of those calling for information (eg, how to give up smoking or the location of the nearest open pharmacy) did not have a symptom recorded. Thus, when analysing calls for information, we did not include symptom as a potential confounding variable. We conducted all analyses in SPSS V.16.0.

RESULTS

Table 2 describes the characteristics of the individual data. Most calls (69%) were for advice; more than half were made by the caller themselves. Most patients (62%) were female; 55% had no ethnicity recorded and the mean age of patients was 33.4 years. Sunday was the most popular day for calls (16.2%). More symptomatic calls concerned digestive symptoms (16.4%) than any other group. Over 40% of callers were advised to contact a GP or a dentist. When WIMD scores were analysed in fifths, 25.1% of calls came from the most deprived fifth. Distances to ED ranged from 0.2 km (from Aberystwyth East in Ceredigion to Bronglais General Hospital) to 56.0 km (from Aberdaron in Gwynedd to Gwynedd Hospital in Bangor). Population density ranged from 0.04 people/ha in Llanuwchllyn in Gwynedd to 100 in Plasnewydd in Cardiff.

Initial exploration showed statistically significant differences between mean WIMD scores by advice given. Calls for advice gave patients living in deprived areas more chance of being told to phone 999: the mean WIMD score of those so advised was 26.4 (95% CI 26.1 to 26.7), while that of those advised to care for themselves was 22.7 (95% CI 22.6 to 22.9). For information calls, the corresponding mean WIMD scores were 24.4 (95% CI 21.7 to 27.1) and 22.1 (95% CI 22.0 to 22.1). When we classified deprivation scores in fifths, these differences

Research report

Table 3 Advice given by WIMD deprivation fifths

Advice given	1 (least deprived)	2	3	4	5 (most deprived)
Calls for advice*	n (%)	n (%)	n (%)	n (%)	n (%)
(1) 999 or ambulance	2086 (3.6)	1540 (3.7)	2114 (4.3)	2671 (4.7)	4235 (5.6)
(2) ED or hospital	6309 (10.9)	4339 (10.5)	5077 (10.4)	6048 (10.6)	7776 (10.3)
(3) Emergency GP or dentist	17330 (29.8)	12335 (29.7)	15164 (31.0)	17907 (31.3)	25239 (33.4)
(4) Other GP or dentist	17057 (29.4)	12203 (29.4)	13915 (28.5)	15881 (27.8)	20388 (27.0)
(5) Other professional	3664 (6.3)	2531 (6.1)	2906 (5.9)	3481 (6.1)	4545 (6.0)
(6) Self-care	10118 (17.4)	7356 (17.7)	8278 (16.9)	9569 (16.7)	11385 (15.0)
(7) Not assessed	1494 (2.6)	1207 (2.9)	1393 (2.9)	1601 (2.8)	2081 (2.8)
Total	58058 (100)	41511 (100)	48847 (100)	57158 (100)	75649 (100)
Calls for information only†	n (%)	n (%)	n (%)	n (%)	n (%)
(1) 999 or ambulance	31 (0.1)	24 (0.1)	20 (0.1)	32 (0.1)	38 (0.1)
(2) ED or hospital	71 (0.3)	48 (0.2)	56 (0.2)	67 (0.2)	74 (0.3)
(3) Emergency GP or dentist	355 (1.4)	281 (1.2)	339 (1.3)	424 (1.5)	528 (2.0)
(4) Other GP or dentist	577 (2.3)	442 (1.9)	511 (2.0)	581 (2.1)	594 (2.2)
(5) Other professional	2032 (8.1)	1627 (7.0)	1746 (6.9)	2116 (7.6)	2483 (9.2)
(6) Self-care	20905 (83.6)	19727 (85.2)	21560 (85.2)	23504 (84.3)	22182 (82.0)
(7) Not assessed	1042 (4.20)	992 (4.3)	1088 (4.3)	1142 (4.1)	1149 (4.2)
Total	25013 (100)	23141 (100)	25320 (100)	27866 (100)	27048 (100)

* χ^2 (df=24)=847; p<0.001.† χ^2 (df=24)=214; p<0.001.

ED, emergency department; GP, general practitioner; WIMD, Welsh Index of Multiple Deprivation.

became clearer, both for advice and for information (table 3). Only 3.6% of callers for advice from the least deprived areas received advice to call 999, compared with 5.6% from the most deprived areas. Advice to contact an emergency GP or dentist was also more frequent in the most deprived areas (33.4% vs 29.8%). Similarly, the frequency of advice to care for themselves was 17.4% in the least deprived areas but 15.0% in the most deprived areas. For information calls, differences were similar but less marked.

These initial findings continued when we included other explanatory variables within logistic regression analyses. This was particularly true for advice calls (table 4): an increase in deprivation from one-fifth to the next increased by 13% the probability of receiving advice to call 999 (OR 1.127; 95% CI 1.113 to 1.143). Moving from one deprivation fifth to the next also increased the probability in advice calls of receiving advice to seek care face to face but by less (OR 1.049; 95% CI 1.041 to 1.058). For information calls (table 5), the impact of deprivation on the advice to call 999 was not significant (OR 1.024; 95% CI 0.912 to 1.149) although there was a slight increase in the probability of receiving advice to seek face-to-face care with deprivation fifth (OR 1.034; 95% CI 1.022 to 1.047).

For variables which appeared in most models, the direction of effect was mainly consistent with the exception of day of the week. For advice calls, the probability of receiving advice to seek face-to-face care increased on Sunday while the probability of receiving advice to seek emergency care increased on Mondays and Thursdays. In all models, those who called NHSDW for themselves always had less probability of receiving more urgent advice; ORs ranged from 0.420 to 0.888 (tables 4 and 5). In both advice and information calls, those whose ethnic status was recorded as non-white had less chance of receiving advice to seek face-to-face care (advice calls: OR 0.818; 95% CI 0.729 to 0.918; information calls: OR: 0.815; 95% CI 0.672 to 0.988).

Most other variables had ORs close to 1, showing little change in the probability of receiving more urgent advice (tables 4 and 5).

DISCUSSION

Main findings

Initial analysis showed that those in more deprived areas generally received more urgent decisions. This trend was consistent across both advice and information. However, this trend was generally weaker in the more comprehensive regression models. Indeed, within information calls deprivation did not affect the probability of receiving advice to seek emergency care. Although other findings were all highly significant statistically, most practical differences were quite small, for example, the odds of being advised to seek care face to face by 5% for each transition from one 'deprivation fifth' to the next highest. There was one exception to this: for advice calls, moving from one deprivation fifth to the next increased the probability of receiving advice to call 999 by 13%. Generally, the same explanatory variables appeared consistently across models with no change in direction and little change in effect size. For example, callers who rang NHSDW for themselves consistently had less chance of receiving more urgent advice than callers on behalf of the patient.

Strengths and limitations of the study

Telephone-based healthcare is an integral part of the urgent and emergency care landscape in Britain both through '111' in England and Scotland and NHSDW in Wales. With those from deprived areas shown to have less clinically effective consultations with other health practitioners,⁵ understanding the outcomes of telephone-based healthcare and how this varies according to patient's characteristics is extremely important. This is the first large study of relationships between patient deprivation and the consequences of telephone-based healthcare, with data on 400 000 calls over 30 months. To explore the influence of deprivation on advice given, we included known

Table 4 Advice calls: likelihood of advising 999 call rather than any other care and advising care face to face rather than self-care

Variable	Advising 999 call rather than any other care			Advising care face to face rather than self-care		
	Stage 1 OR (95% CI)***	Stage 2 OR (95% CI)***	Stage 3 OR (95% CI)***	Stage 1 OR (95% CI)***	Stage 2 OR (95% CI)***	Stage 3 OR (95% CI)***
Distance to emergency department (per mile)	1.006 (1.004 to 1.009)	1.006 (1.004 to 1.009)	1.006 (1.003 to 1.008)	0.990 (0.988 to 0.991)	0.990 (0.988 to 0.991)	0.990 (0.988 to 0.991)
Population density (people per hectare)	1.002 (1.001 to 1.004)	1.002 (1.001 to 1.004)	1.002 (1.001 to 1.003)**	1.001 (1.001 to 1.002)**	1.001 (1.001 to 1.002)**	1.001 (1.001 to 1.002)**
Patient age (per year)	1.026 (1.026 to 1.027)	1.027 (1.026 to 1.027)	1.027 (1.026 to 1.027)	1.010 (1.010 to 1.011)	1.010 (1.009 to 1.011)	1.010 (1.009 to 1.011)
Gender						
Male	1.00	1.00	1.00	1.00	1.00	1.00
Female	0.873 (0.839 to 0.908)	0.874 (0.840 to 0.909)	0.871 (0.837 to 0.906)	1.128 (1.100 to 1.156)	1.126 (1.099 to 1.154)	1.125 (1.098 to 1.153)
Relationship						
Surrogate caller	1.00	1.00	1.00	1.00	1.00	1.00
Self-caller	0.418 (0.401 to 0.436)	0.414 (0.397 to 0.432)	0.420 (0.403 to 0.438)	0.845 (0.821 to 0.869)	0.860 (0.836 to 0.885)	0.864 (0.840 to 0.890)
Non-white race						
White or unknown	1.00	1.00	1.00	1.00	1.00	1.00
Non-white	1.123 (0.923 to 1.366)*	1.122 (0.922 to 1.364)*	1.13 (0.929 to 1.375)*	0.806 (0.718 to 0.905)	0.813 (0.725 to 0.913)	0.818 (0.729 to 0.918)
Known race						
Unknown	1.00	1.00	1.00	1.00	1.00	1.00
Known (white or other)	1.212 (1.166 to 1.260)	1.213 (1.166 to 1.260)	1.211 (1.165 to 1.259)	1.169 (1.141 to 1.198)	1.170 (1.142 to 1.199)	1.169 (1.140 to 1.197)
Symptom						
All other symptoms	1.00	1.00	1.00	1.00	1.00	1.00
Digestive symptom	0.307 (0.289 to 0.326)	0.305 (0.287 to 0.325)	0.305 (0.287 to 0.324)	0.627 (0.612 to 0.643)	0.632 (0.616 to 0.648)	0.631 (0.616 to 0.648)
Day of call						
Sunday	Not entered	1.00	1.00	Not entered	1.00	1.00
Monday	Not entered	1.075 (1.004 to 1.150)**	1.076 (1.005 to 1.151)**	Not entered	0.778 (0.745 to 0.812)	0.779 (0.746 to 0.813)
Tuesday	Not entered	1.058 (0.986 to 1.135)*	1.058 (0.986 to 1.135)*	Not entered	0.779 (0.745 to 0.814)	0.778 (0.745 to 0.813)
Wednesday	Not entered	1.062 (0.991 to 1.138)*	1.063 (0.991 to 1.139)*	Not entered	0.774 (0.740 to 0.809)	0.774 (0.740 to 0.809)
Thursday	Not entered	1.136 (1.061 to 1.215)	1.126 (1.053 to 1.205)**	Not entered	0.832 (0.796 to 0.870)	0.830 (0.794 to 0.867)
Friday	Not entered	1.054 (0.983 to 1.131)*	1.056 (0.984 to 1.133)*	Not entered	0.740 (0.708 to 0.773)	0.740 (0.708 to 0.773)
Saturday	Not entered	0.903 (0.848 to 0.962)**	0.902 (0.847 to 0.961)**	Not entered	0.844 (0.810 to 0.880)	0.844 (0.810 to 0.879)
Deprivation fifth (ordinal)	Not entered	Not entered	1.127 (1.113 to 1.143)	Not entered	Not entered	1.049 (1.041 to 1.058)

*p>0.05 (therefore non-significant), **p<0.05, ***p<0.001 unless otherwise indicated.

ORs are the odds of receiving advice to call 999 or seek care face-to-face relative to their respective baselines in the logistic regression models comprising the above variables.

Table 5 Information calls: likelihood of advising 999 call rather than any other care and advising care face to face rather than self-care

Variable	Advising 999 call rather than any other care			Advising care face to face rather than self-care		
	Stage 1	Stage 2	Stage 3	Stage 1	Stage 2	Stage 3
	OR (95% CI)***	OR (95% CI)***	OR (95% CI)***	OR (95% CI)***	OR (95% CI)***	OR (95% CI)***
Distance to ED (per mile)	0.989 (0.970 to 1.009)*	0.991 (0.972 to 1.010)*	0.991 (0.971 to 1.01)*	0.986 (0.984 to 0.988)	0.988 (0.986 to 0.990)	0.988 (0.986 to 0.990)
Population density (people per hectare)	1.003 (0.993 to 1.014)*	1.003 (0.992 to 1.014)*	1.003 (0.992 to 1.014)*	1.003 (1.002 to 1.005)	1.003 (1.002 to 1.004)	1.003 (1.002 to 1.004)
Patient age (per year)	1.013 (1.005 to 1.021)**	1.013 (1.005 to 1.021)**	1.013 (1.005 to 1.021)**	1.001 (1.000 to 1.002)**	1.001 (1.000 to 1.002)**	1.001 (0.973 to 1.049)**
Gender						
Male	1.00	1.00	1.00	1.00	1.00	1.00
Female	1.036 (0.726 to 1.479)*	1.03 (0.721 to 1.470)*	1.029 (0.721 to 1.470)*	1.018 (0.980 to 1.057)*	1.011 (0.973 to 1.050)*	1.010 (0.973 to 1.049)*
Relationship						
Surrogate caller	1.00	1.00	1.00	1.00	1.00	1.00
Self-caller	0.431 (0.296 to 0.627)	0.458 (0.314 to 0.669)	0.460 (0.315 to 0.672)	0.673 (0.644 to 0.704)	0.713 (0.682 to 0.746)	0.716 (0.685 to 0.749)
Non-white race						
White or unknown	1.00	1.00	1.00	1.00	1.00	1.00
Non-white	0.888 (0.122 to 6.449)*	0.904 (0.124 to 6.566)*	0.906 (0.125 to 6.585)*	0.794 (0.655 to 0.963)**	0.811 (0.669 to 0.983)**	0.815 (0.672 to 0.988)**
Known race						
Unknown	1.00	1.00	1.00	1.00	1.00	1.00
Known (white or other)	0.762 (0.542 to 1.070)*	0.766 (0.546 to 1.076)*	0.766 (0.545 to 1.076)*	0.950 (0.917 to 0.983)**	0.955 (0.922 to 0.989)**	0.954 (0.921 to 0.988)**
Symptom						
All other symptoms	Not entered	Not entered	Not entered	Not entered	Not entered	Not entered
Digestive symptom	Not entered	Not entered	Not entered	Not entered	Not entered	Not entered
Day of call						
Sunday	Not entered	1.00	1.00	Not entered	1.00	1.00
Monday	Not entered	0.734 (0.388 to 1.389)*	0.735 (0.388 to 1.390)*	Not entered	0.598 (0.560 to 0.639)	0.599 (0.561 to 0.639)
Tuesday	Not entered	0.622 (0.316 to 1.224)*	0.623 (0.317 to 1.225)*	Not entered	0.590 (0.551 to 0.630)	0.590 (0.552 to 0.631)
Wednesday	Not entered	0.912 (0.487 to 1.705)*	0.912 (0.488 to 1.706)*	Not entered	0.625 (0.585 to 0.668)	0.625 (0.585 to 0.669)
Thursday	Not entered	0.591 (0.294 to 1.187)*	0.591 (0.294 to 1.188)*	Not entered	0.669 (0.626 to 0.715)	0.670 (0.627 to 0.716)
Friday	Not entered	0.889 (0.470 to 1.680)*	0.889 (0.470 to 1.681)*	Not entered	0.626 (0.585 to 0.670)	0.627 (0.585 to 0.670)
Saturday	Not entered	1.349 (0.724 to 2.513)*	1.349 (0.724 to 2.513)*	Not entered	1.007 (0.941 to 1.078)*	1.007 (0.941 to 1.078)*
Deprivation fifth (ordinal)	Not entered	Not entered	1.024 (0.912 to 1.149)*	Not entered	Not entered	1.034 (1.022 to 1.047)

*p<0.05 (therefore non-significant); **p<0.05; ***p<0.001 unless otherwise indicated.

ORs are the odds of receiving advice to call 999 or seek care face-to-face relative to their respective baselines in the logistic regression models comprising the above variables.

confounding variables from previous studies of deprivation and healthcare.^{13–15} We used accepted methods to overcome methodological issues like ranking advice by urgency and inferring distances to hospitals. We modelled the relationships between deprivation and advice by separating calls for advice from those for information only. The main limitation was the lack of any measure of symptom severity. Although we used the ICPC-2 coding system to summarise patient symptoms, this does not address severity. Thus, this dataset cannot tell whether two different callers with ‘digestive’ symptoms had similar levels of dysfunction. Similarly, we do not know whether those calling from deprived areas had worse health and how this affected the advice given. Another limitation was the absence of personal addresses from our dataset, with the result that inferences about individual characteristics stemmed from ward-level data.²⁶ Although we have used proxies like the geometric centre of a ward to calculate distance to ED, this method does not discriminate between types of journey, for example, mountainous or motorway. Finally, we could not include those 59 523 calls (12%) without recorded wards; these may represent a distinctive group of callers, for example, genuine emergencies without time to collect all information or uncooperative callers who refused to give their address. Despite these limitations, our findings have remained consistent: in this national dataset, there is a small to moderate effect of deprivation on the advice given.

Findings in context

One of the difficulties in analysing advice given using routine data from NHSD is the focus on one simplified outcome. Many aspects of calls, for example, time or psychological state of the caller, could have influenced the final advice. Although we have analysed the most urgent advice given, any other advice given within the phone call disappears unless specifically recorded. We do not know who made the final decision—the CDSS or the nurse advisor by overriding that system. Furthermore, as we cannot identify repeat callers, we cannot infer how the advice affected future contacts.

Although data are from a period of time when NHSDW was relatively new, there has been no new research or policy changes related to socioeconomic deprivation that suggest our findings are not still relevant. Indeed, our findings are generally consistent with literature suggesting that those more deprived receive more urgent outcomes, both from other emergency healthcare services^{6–8} and from telephone-based healthcare.²⁷ In particular, our findings resemble those of O’Reilly and colleagues¹⁵ who reported that the probability of seeing a GP out of hours is only slightly increased by deprivation. In short, by specifying a fuller range of independent variables, our models better estimate the true effect of deprivation.

Implications

The tendency for those living in more deprived areas to receive more urgent outcomes from NHSDW has important consequences for policy, practice and research. Although all differences were small, they were highly significant statistically. With those living in deprived areas known to have poorer health, there is a need for further research to understand these findings. For example, do these differences result from inequalities in health or in healthcare-seeking behaviour: is this tendency a true consequence of poorer health or an artefact of the pattern of communication between nurse advisor and caller? Are callers from more affluent areas more likely to define their concerns more clearly, thus avoiding the need for care face to face? There is also a need

to characterise and quantify these types of results in terms of financial effects on the NHS and individual patients. Though we compared advice to call 999 with all other forms and care face to face with self-care, we recommend future researchers model the full range of advice and thus continue to build fully specified regression models of advice given in telephone healthcare out of hours. Qualitative interviews with callers and nurse advisors and analysis of call transcripts should explore reasons for contact with NHSD, explanations for variations in advice given across groups and the relationship between nurse advisors and patients.

Conclusions

Telephone advice for first contact healthcare is now playing a large and growing role in the emergency care landscape. In order to provide better care and to address public health priorities related to inequalities, it is vital to understand and to respond to these new findings about the level of advice given to people living in deprived areas. We do not know from this study whether the advice is warranted and therefore is an indicator of good practice or is a reaction to unknown factors which do not reflect clinical need. While our findings show that NHSDW is not disadvantaging to those living in deprived areas, there is a strong need for further research in this area.

What is already known on this topic

- ▶ National Health Service (NHS) Direct and other providers of healthcare by telephone use computerised decision support software to advise patients on the care most appropriate to their needs.
- ▶ Those living in deprived areas generally receive more urgent healthcare out of hours, but the effect of deprivation on advice given by NHS Direct is not known.

What this study adds

- ▶ Analysis of over 400 000 calls to National Health Service (NHS) Direct Wales showed that after adjustment for confounding variables, increased patient deprivation had a small to moderate positive effect on receiving more urgent advice.
- ▶ Other factors that made patients more likely to receive advice to take urgent action included calls made on their behalf, calls on Sundays, and calls about white patients.
- ▶ While this study suggests that advice given by NHS Direct Wales is more equitable than feared, it advocates more research into the influence of patient and call characteristics.

Contributors JP and HS designed the study. JP cleaned, managed and analysed the data, drafted and revised the paper. She is the guarantor. SM provided expertise in informatics, SR expertise in medical geography, IR and AW expertise in statistics and HS expertise in the field of emergency care research. All authors revised the draft paper.

Competing interests None declared

Patient consent Detail has been removed from this case description/these case descriptions to ensure anonymity. The editors and reviewers have seen the detailed information available and are satisfied that the information backs up the case the authors are making.

Ethics approval South East Wales Local Ethics Committee.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement A more detailed description of the data is contained in Dr Julie Peconi’s PhD Thesis ‘The epidemiology of demand for and outcomes of contacts with telephone based healthcare with particular reference to ward

Research report

deprivation scores: Analysis of calls to NHS Direct Wales 2002–2004'. Please contact j.peconi@swansea.ac.uk for more details.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2017. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- Goddard M, Smith P. Equity of access to health care services: theory and evidence from the UK. *Soc Sci Med* 2001;53:1149–62.
- Morris S, Sutton M, Gravelle H. Inequity and inequality in the use of health care in England: an empirical investigation. *Soc Sci Med* 2005;60:1251–66.
- Balfour JL, Kaplan GA. Neighbourhood environment and loss of physical function in older adults: evidence from the Alameda County study. *Am J Epidemiol* 2002;155:507–15.
- Galea S, Ahern J, Nandi A, et al. Urban neighbourhood poverty and the incidence of depression in a population-based cohort study. *Ann Epidemiol* 2007;17:171–9.
- Hart JT. The inverse care law. *Lancet* 1971;i:405–12.
- Pollock AM, Vickers N. Deprivation and emergency admissions for cancers of colorectum, lung, and breast in south east England: ecological study. *BMJ* 1998;317:245–52.
- O'Donnell CA, McConnachie A, Moffat K, et al. Cross sectional study of social variation in use of an out of hours patient transport service. *BMJ* 1999;318:566–7.
- Beattie TF, Gorman DR, Walker JJ. The association between deprivation levels, attendance rate and triage category of children attending a children's accident and emergency department. *Emerg Med J* 2001;18:110–1.
- Department of Health. *High quality care for all: NHS next stage review final report: strategic health authorities' visions for better healthcare*. London: DH, 2008.
- Welsh Ambulance Service NHS Trust | 111 Wales. <http://www.was-tr.wales.nhs.uk/Default.aspx?pagelid=315&lan=en> (accessed 04 Apr 2017).
- O'Cathain A, Nicholl J, Sampson F, et al. Do different types of nurses give different triage decisions in NHS Direct? A mixed methods study. *J Health Serv Res Policy* 2004;9:226–33.
- Monaghan R, Clifford C, McDonald P. Seeking advice from NHS Direct on common childhood complaints: does it matter who answers the phone? *J Adv Nurs* 2003;42:209–16.
- Munro J, Maheswaran R, Pearson T. Response to requests for general practice out of hours: geographical analysis in north west England. *J Epidemiol Community Health* 2003;57:673–4.
- Turnbull J, Pope C, Martin D, et al. Management of out-of-hours calls by a general practice cooperative: a geographical analysis of telephone access and consultation. *Fam Pract* 2011;28:677–82.
- O'Reilly D, Stevenson M, McCay C, et al. General practice out-of-hours service, variations in use and equality in access to a doctor: a cross-sectional study. *Br J Gen Pract* 2001;51:625–9.
- Welsh Index of Multiple Deprivation. *Local Authority profiles*. Cardiff: Welsh 322 Office, 2000. <http://wales.gov.uk/statistics-and-research/welsh-index-multiple-deprivation/?lang=en> (accessed 6 Dec 2013).
- World Health Organisation. *International classification of primary care*. 2nd ed. Geneva, Switzerland: WHO, 2003. (ICPC-2). <http://www.who.int/classifications/icd/adaptations/icpc2/en> (accessed 6 Dec 2013).
- Munro J, Clancy M, Knowles E, et al. *Evaluation of NHS direct: impact and appropriateness*. Sheffield University: Medical Care Research Unit, 2003. http://sheffield.ac.uk/content/1/c6/02/40/50/nhsd_phase2.pdf (accessed 6 Dec 2013).
- Snooks H, Peconi J, Munro J, et al. An evaluation of the appropriateness of advice and healthcare contacts made following calls to NHS Direct Wales. *BMC Health Serv Res* 2009;9:178.
- Hanigan I, Hall G, Dear KB. A comparison of methods for calculating population exposure estimates of daily weather for health research. *Int J Health Geogr* 2006;5:38.
- Judge A, Welton NJ, Sandhu J, et al. Equity in access to total joint replacement of the hip and knee in England: cross sectional study. *BMJ* 2010;341:c4092.
- Geoconvert: UK data service census support. www.geoconvert.mimas.ac.uk (Accessed 6 December 2013).
- Hajat S, Kovats RS, Lachowycz K. Heat-related and cold-related deaths in England and Wales: who is at risk? *Occup Environ Med* 2007;64:93–100.
- Gasparrini A, Armstrong B, Kovats S, et al. The effect of high temperatures on cause-specific mortality in England and Wales. *Occup Environ Med* 2012;69:56–61.
- Choi M, Curriero FC, Johantgen M, et al. Association between ozone and emergency department visits: an ecological study. *Int J Environ Health Res* 2011;21:201–21.
- Morgenstern H. Uses of ecologic analysis in epidemiologic research. *Am J Public Health* 1982;72:1336–44.
- Sullivan CO, Omar RZ, Forrest CB, et al. Adjusting for case mix and social class in examining variation in home visits between practices. *Fam Pract* 2004;21:355–63.



Advice given by NHS Direct in Wales: do deprived patients get more urgent decisions? Study of routine data

Julie Peconi, Steven Macey, Sarah Rodgers, Ian Russell, Helen Snooks and Alan Watkins

J Epidemiol Community Health published online July 21, 2017

Updated information and services can be found at:

<http://jech.bmj.com/content/early/2017/07/20/jech-2017-208978>

These include:

References

This article cites 21 articles, 8 of which you can access for free at: <http://jech.bmj.com/content/early/2017/07/20/jech-2017-208978#ref-list-1>

Open Access

This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Topic Collections

Articles on similar topics can be found in the following collections

[Open access](#) (327)

Notes

To request permissions go to:

<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:

<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:

<http://group.bmj.com/subscribe/>