Prostitution in Lagos: a sociomedical study

U. G. OLERU

From the Department of Community Health, College of Medicine, University of Lagos

SUMMARY A study of 150 hotel prostitutes in Lagos was undertaken to determine their socioeconomic identity and, through an assessment of their health knowledge, attitude, and practice, their probable impact on public health. The subjects were selected from 15 hotels, representing 20% of the estimated hotel universe in Lagos and different socioeconomic strata. Information was obtained from the subjects with the help of the hotel staff, who arranged pre-interview conferences with them. Prostitution in Lagos is not organised and there are no figures on the prostitute population. Ninety-eight per cent of the women in the study population were aged between 15 and 44 and 70% of them had children. Forty per cent were married and 24% were either divorced or separated. The most important reason given for prostitution was unhappiness in the home (50%) but the undercurrent of poverty, especially in a low-income society, is undoubtedly a major motive. More than half (55.3%) of the subjects selected their clients and the most important criteria for selection were physical fitness and ability to pay. Physical fitness was defined as the absence of signs of gonorrhea. The women's level of education was a significant factor in selection of clients and in mode of health care. More than 60% of those who selected their clients had had some education or had completed secondary schooling. Thirty-three per cent of those who did not select their clients had had no education at all and 45% had received primary education only. The attitude of the subjects to health care suggests an opportunity for the control of disease among prostitutes if prostitution is organised.

Social and legal attitudes to prostitution vary greatly between countries. In a few countries where prostitution has been legalised, simultaneous attempts have been made to eliminate the threat to public health. In these countries, the production of records of periodic medical examination has been made a licensing requirement.¹

In Nigeria prostitution is regarded as a social evil, but rapid urbanisation of Nigerian communities and the strains this has imposed on traditional family structures² have helped to reduce hostility to it. In Nigerian cities, especially in Lagos, prostitution is of the street and hotel type. Prostitutes are not registered so their numbers are not precisely known.

The objective of this study was to determine the socioeconomic identity of prostitutes in Lagos, and their probable impact on public health, through an assessment of their health knowledge, attitude and practice.

Materials and Methods

Fifteen hotels were randomly selected from a list of 50. There is no accurate register of hotels and establishments in Lagos and the sample size probably accounts for 20% of the hotel universe in the city. The breakdown of the hotels according to the socioeconomic status of their patrons is shown in Table 1.

The hotel staff provided the investigating team with an estimate of the number of prostitutes who ply each hotel. The estimated population of prostitutes in 15 hotels where interviews were conducted was 295. One hundred and fifty subjects were included in the study.

All interviews were conducted in rooms provided by the hotel staff. The interview team consisted of three workers who were dressed in the medical outfits of a doctor, a nurse, and a laboratory

Table 1 *Hotels by socioeconomic category, number of subjects, and estimated population of prostitutes in study sample*

HOTELS			SUBJECTS		ESTIMATED POPULATION OF PROSTITUTES
Category	No.	%	No.	%	
High income, international grade	3	20	34	22.67	50
Middle income, limited international patronage	5	33.33	56	37.33	90
Middle to low income	7	46.66	60	40.00	155
TOTAL	15	99.99	150	100.00	295

technician, so as to emphasise to the subjects that the interviewers were not a law enforcement team. Subjects were invited into the room one at a time to eliminate peer influence.

A questionnaire was administered to each of the 150 subjects to elicit information on age, marital status, family history, gravidity, parity, educational background, motives for prostitution, health care, symptoms of sexually transmitted diseases, drug use, and housing.

Results

MARRIAGE, FAMILY, AND PROSTITUTION

Table 2 shows the marital status of the subjects. Sixty-one (40.7%) were currently married and over 70% had been married at some time.

Table 2 Marital status of subjects

	No.	%
Married	61	40.7
Single	40	26.7
Separated	24	16.0
Divorced	12	8.0
Widowed	13	8.6
TOTAL	150	100.00

The age distribution of the subjects indicated that 147 (98%) were between 15 and 44, the active reproductive years. Only 12 of the 147 were aged between 15 and 19.

Forty-one (27.3%) were childless; 70 (46.7%) had one to three children and 34 (22.7%) had four to six children. In all, 109 subjects had to raise families in

addition to practising prostitution and 63 of these (57.8%) had their children living with them.

Table 3 shows the association between marital status and the motive given for prostitution. It seems that the association is significant ($\chi^2 = 34.83$; $P < 0.001$). For women with a living spouse (that is, married, separated or divorced), the most important reason given for adopting prostitution was desertion, unhappiness in the home, or a broken home. Among the widowed subjects, poverty (30.77%) and unhappy homes (33.4%) were stated to be the most important reasons. Among single women, the most important reason given for prostitution was sexual pleasure. The Nigerian civil war, 1966–70, did not have a significant impact on the choice of prostitution by the subjects.

EDUCATION, HEALTH, AND PROSTITUTION

Table 4 shows the association between the level of education and the motive for prostitution. With the exception of the 34 women with no formal education, the proportion of individuals reporting an unhappy home as the motive for prostitution decreased as the level of education increased. This suggests that a higher level of education may have offered subjects from unhappy homes better options for making a living than prostitution.

Table 5 shows the association between educational level and choice of client. Among the 83 subjects who selected their clients this association was statistically significant ($P < 0.01$). The criteria for selection were both physical fitness and ability to pay, ability to pay

Table 3 Association between marital status and motives for prostitution

MARITAL STATUS	POVERTY		UNHAPPY HOME, DESERTION, BROKEN HOME		SEXUAL PLEASURE		CIVIL WAR		TOTAL	
	No.	%	No.	%	No.	%	No.	%	No.	%
Single	4	10	11	27.5	19	47.5	6	15	40	100
Married	12	19.67	33	54.10	13	21.31	3	4.92	61	100
Separated	5	20.83	18	75.0	—	—	1	4.17	24	100
Divorced	3	25.0	8	66.66	—	—	1	8.33	12	100
Widowed	4	30.77	5	38.46	1	7.69	3	23.08	13	100
TOTAL	28	18.77	75	50.00	33	22.00	14	9.33	150	1000

$\chi^2 = 34.83$ $P < 0.001$

Table 4 Association between level of education and motives for prostitution

EDUCATION	BROKEN OR UNHAPPY HOME		PLEASURE		POVERTY		CIVIL WAR		TOTAL	
	No.	%	No.	%	No.	%	No.	%	No.	%
No schooling	16	47.06	4	11.76	10	29.4	4	11.76	34	100
Primary	32	62.74	6	11.76	8	15.69	5	9.8	51	100
Incomplete secondary education	17	48.57	10	28.57	4	11.43	4	11.43	35	100
Complete secondary education	10	33.33	13	43.33	6	20.0	1	3.33	30	100
TOTAL	75	50.0	33	22.0	28	18.67	14	9.33	150	100

$\chi^2 = 17.68$ $P < 0.05$

Table 5 Association between level of education and selection of patrons

EDUCATION	PHYSICAL FITNESS AND ABILITY TO PAY		PHYSICAL FITNESS ALONE		ABILITY TO PAY ALONE		TOTAL	
	No.	%	No.	%	No.	%	No.	%
No schooling	2	22.22	2	22.22	5	55.56	9	100
Primary	6	25.9	5	20.83	13	54.16	24	100
Incomplete secondary education	17	58.62	5	17.24	7	24.14	29	100
Complete secondary education	14	66.66	3	19.05	4	19.05	21	100
TOTAL	39	46.99	15	18.07	29	34.94	83	100

 $\chi^2 = 19.44$ $P < 0.01$

Table 6 Association between level of education and selection of health care

EDUCATION	DOCTOR OR HOSPITAL		DRUG STORE PRESCRIPTION		SELF-MEDICATION		HERBALIST		TOTAL	
	No.	%	No.	%	No.	%	No.	%	No.	%
No schooling	11	32.35	15	44.11	5	14.7	3	8.82	34	100
Primary	18	35.3	21	41.18	12	23.53	—	—	51	100
Incomplete secondary education	18	51.43	10	28.57	4	11.43	3	8.57	35	100
Complete secondary education	24	80.0	3	10.0	3	10.0	—	—	30	100
TOTAL	71	47.33	49	32.67	24	15.99	6	3.99	150	100

 $\chi^2 = 26.90$ $P < 0.005$

alone, and physical fitness alone. Physical fitness was defined as absence of signs of venereal disease, especially gonorrhea, as evidenced by the urethra. Thirty-nine (47%) of the women took into account both the physical fitness of their clients and their ability to pay. Among those who did not choose their clients, 22 (32.83%) had had no formal education at all, 30 (44.78%) had had primary education only, eight (11.94%) did not complete secondary education, and seven (10.45%) did.

Only 35 (52.22%) of those who did not select their clients could positively identify the signs of gonorrhea in a man.

Table 6 shows the relationship between education and approach to health care. This association was statistically significant ($P < 0.005$). Nearly half of the subjects (47%) chose to visit the hospital or doctor's clinic, but women who had completed their secondary education were more likely to do this than less well-educated women. This relationship between selection of hospital or clinic care and level of education probably holds true for the general population, although there are no available data to support this assumption.

The next most important choice of health care was a visit to the chemist or drug store. The latter are sometimes manned by trained pharmacists but more often by untrained personnel whose prescription of particular drugs is as likely to be influenced by the availability of those drugs in the store as by their customers' complaints.

Table 7 Conditions for which subjects had self-medication or medication prescribed by a doctor in the six months before the study

CONDITION	No.	%
Vaginal itching and discharge	62	41.33
Vaginal discharge and joint pain	40	26.67
Skin diseases	30	20.00
Sores	6	4.00
Cough	12	8.00
TOTAL	150	100.00

FREQUENCY OF DISEASE

All the subjects stated that in the six months before the study they had treated themselves for an illness. Seventy-two subjects (48%) said they had treated themselves only once, and 56 subjects (37.3%) said they had treated themselves twice; the rest had treated themselves more frequently.

Table 7 shows their conditions. It is evident that at least about two-thirds of these were venereal. This conclusion is suggested by the fact that 61 (40.7%) had taken antibiotics, 46 Panadol, 29 (19.33%) had used vaginal pessaries, and 16 (10.66%) had used skin cream.

DRUG USE, SMOKING, AND PROSTITUTION

The average daily income of the subjects ranged between N 2400 and N 4000. At the time of the survey N 1.00 was equivalent to \$1.70. This income is substantially higher than average; only 30% of the Nigerian population earn more than N 1000 a year (Shobanjo 1979, personal communication).

None of the women were dependent on drugs before becoming prostitutes. However, the use of non-addictive drugs developed among them in the course of prostitution and their substantial income provided them with the means to support the habit. One hundred and twenty-one of the subjects (80.7%) admitted to taking drugs. Of these, 32 (26.4%) took them to be more relaxed or to sleep, 25 (20.7%) took them to keep awake, and 16 (13.2%) to be 'high'. The preparations used included Valium, amphetamines, and several containing ephedrine. These suit the demands of prostitution—be alert and active when there are clients, sleep when business is slow. No subject admitted to using 'hard' drugs.

Discussion

In a study in Denver and Philadelphia, Weissman and File³ observed a tendency for women who were drug addicts to turn to prostitution and other crimes to support their drug habit. The women in this study had no history of serious crimes and opted for prostitution on the grounds of unhappiness in the home (50%) and poverty (18.67%). Although only 18.7% of the subjects specifically identified poverty as the main reason for prostitution, poverty probably had a lot to do with the decision to become a prostitute. The use of drugs differed from that described by Weissman and File.³ The subjects took drugs acting on the central nervous system, dependence-producing drugs to satisfy the dictates of prostitution to be awake or to sleep when necessary.

Although 54 (36%) of the women refused to accept clients whom they suspected of having venereal disease, more than 100 admitted to having suffered

signs and symptoms of sexually transmitted conditions at least once in the six months before the study.

Only 71 (66.7%) of the women sought health care from doctors or hospitals and this choice was significantly influenced by the level of education. Regular medical examination could be introduced as a means of controlling these diseases.

The family history of the subjects indicated that a large proportion (69.4%) had children to care for. Some of these children lived with the subjects and others with relatives, especially grandmothers. The psychological impact of prostitution on the children of the subject is the topic of another study.

We thank the managements of the hotels where the study was conducted and also Mr. P. Alabunge of the Data Processing Unit of the Department of Community Health for technical assistance.

Reprints from Dr. O. G. Oleru, Department of Community Health, College of Medicine, University of Lagos, Private Mail Bag 12003, Lagos, Nigeria.

References

- ¹Borelli S. Problems of prostitution. *Mater Med Nord* 1969; **10**: 489–501.
- ²Mabogunje AL. *Urbanisation in Nigeria*. London: University of London Press, 1968.
- ³Weissman JC, File KN. Criminal behaviour patterns of female addicts: a comparison of findings in two cities. *Int J Addict* 1976; **16**: 1063–75.