Ischaemic heart disease in England and Wales around the year 2000

C Osmond, D J P Barker

Abstract

Study objective—The aim was to predict death rates from ischaemic heart disease in England and Wales during the next 20 years.

Study design—Age and sex specific death rates for 1968–1987 were used to predict national and regional trends until 2007.

Setting—The nine standard regions of England and Wales.

Results—There will be a large fall in deaths from ischaemic heart disease. The fall will be greater in men and women below 65 years of age. There will, however, be a sharp worsening of the north/south divide. The largest fall in any region will be in Wales.

Conclusions—The campaign to change the national diet should give highest priority to the northern regions. Because infant growth is inversely related to adult risk of ischaemic heart disease, the poor growth of young children in some northern areas is a cause for concern.

Death rates from ischaemic heart disease are now falling in Britain,¹ but they remain high in comparison with other countries.² Future rates can be predicted from recent trends. This prediction must distinguish changes in successive generations from other sources of year to year variation. We present predicted rates for England and Wales, in men and women, in each age group and within each region, into the next century.

Methods

We abstracted numbers of deaths from ischaemic heart disease (International Classification of Diseases codes 410–414 in 8th and 9th revisions) and population sizes from 1968 to 1987 for England and Wales and for the nine standard regions by age and sex. These data were used to predict future death rates, allowing for cohort (generation) effects. The statistical model used for prediction partitioned the age and sex specific rates for 1968 to 1987 according to age, year of death, and cohort.³ The year of death and cohort components were extended into the future, and then combined to predict age specific death rates for 1988 to 2007.

We calculated regional rates for deaths in middle age, which is by convention from 45 to 64 years. These were directly standardised to the European standard population,⁴ and are presented as mortality ratios based on the figure of 100 for England and Wales.

Results

The figure shows age specific death rates during 1968–87 by year of birth. Rates among men fell progressively at all ages in generations born from 1928 onwards. Similarly among women rates fell in generations born from 1933 onwards. This demonstrates a cohort effect.

Table I shows death rates during 1968–87 and predicted rates for 1988–2007. Rates will fall in both men and women and in each age group. They will fall more in younger than older people. Up to 2003–7 there will be a 43% fall in men aged 45–49 years in relation to 1983–7 rates. The fall at ages 65–69 will be 25%. The corresponding figures for women will be 44% and 18%.

Table II shows observed and predicted mortality ratios by region among middle aged people. There will be a widening in regional differences. The highest death rates will continue to be in the three northern regions, but their ratios to the national average will increase. A large rise in the north west will take the ratio to 137. The ratios in both midland regions will rise to exceed 100. The lowest death rates will continue to be in East Anglia and the two southern regions, but their ratios to the national average will fall. The fall in East Anglia will take the ratio to 70. The position of Wales will improve sharply, the ratio falling from 121 to 90.

Discussion

If the trends over the past 20 years continue there will be a large fall in premature deaths from ischaemic heart disease in England and Wales. The fall at ages over 65 years will be less. There will be a sharp worsening of the north-south divide. Mortality in the three northern and two midland regions will worsen in comparison with East Anglia and the two southern regions. In Wales death rates at younger ages have fallen sharply in successive generations and predict a continuing large fall in premature deaths. There may be additional benefits arising from the prevention project 'Heart-Beat Wales'.

Studies of both fatal and non-fatal myocardial infarction in the United States of America, Finland, Australia, and New Zealand have shown that death rates are falling and are accompanied by falls in incidence.⁵ Except in New Zealand, rates are falling more in younger than older age groups.

Two recent surveys of adult diet have shown that dietary fat intake does not differ greatly between one part of England and Wales and another.^{6,7} Nevertheless because of their greater susceptibility, however caused, implementation of the recommendations made by the Committee on Medical Aspects of Food Policy is more urgent

MRC Environmental
Epidemiology Unit,
University of
Southampton,
Southampton General
Hospital,
Southampton SO9
4XY, United Kingdom
C Osmond
D J P Barker

Correspondence to:
Dr Osmond

Accepted for publication May
1990

Ischaemic heart disease death rates in England and Wales, ages 35–74 years, 1968–1987. Rates are per million person-years. (A) men, (B) women.

Table 1 Observed and predicted death rates per million years at risk from ischaemic heart disease among men and women aged 35–74 years in England and Wales, 1968 to 2007.

Period	Age group (years)						
	35–39	40–44	45–49	50–54	55–59	60–64	65–69
Men							
Observed:							
1968–1972	344	945	1972	3432	5524	8697	13674
1973–1977	326	908	1976	3624	5681	8793	13651
1978–1982	295	793	1765	3435	5657	8771	13094
1983–1987	232	623	1466	2834	5027	8154	12482
Predicted:							
1988–1992	205	563	1281	2592	4583	7923	12607
1993–1997	178	488	1105	2206	4050	7007	11898
1998–2002	153	422	959	1903	3448	6193	10524
2003–2007	133	365	829	1651	2975	5272	9301
Women							
Observed:							
1968–1972	55	147	290	621	1226	2608	5065
1973–1977	54	157	339	666	1346	2697	5083
1978–1982	47	130	286	670	1371	2683	4915
1983–1987	39	91	235	568	1299	2646	4791
Predicted:							
1988–1992	33	94	192	508	1197	2734	5032
1993–1997	29	78	185	401	1038	2447	5080
1998–2002	25	67	153	385	819	2122	4546
2003–2007	21	58	131	319	787	1674	3942

Table 11 Observed and predicted mortality ratios for ischaemic heart disease among men and women aged 45–64 years in the regions of England and Wales, 1968–2007

Region	Observed in			
	1968–77	1978–87	1988–97	1998–2007
North	121	124	128	128
North West	119	123	127	137
Yorkshire and Humberside	115	116	118	120
East Midlands	97	100	106	115
West Midlands	99	102	107	112
East Anglia	80	76	72	70
South East	87	85	82	80
South West	92	88	83	77
Wales	121	114	107	90

among men and women in the northern and midland regions. Currently in England only around 25% of men and women aged 35–54 years have fat intakes below the recommended 35% of total energy intake, or ratios of polyunsaturated to saturated fatty acids above the recommended 0.45.⁶

The distribution of mortality from ischaemic heart disease throughout England and Wales is closely related to the distribution of neonatal mortality 50 and more years ago.⁹ This indication that growth and development during prenatal life and infancy are important risk factors for the disease is supported by the results of a prospective study that showed a strong inverse relation between infant weights and death rates.¹⁰ Given the inverse relation between infant growth and ischaemic heart disease the current poor growth of young children in the inner city area of Liverpool,¹¹ and in other northern towns such as Middlesbrough (Harland E, personal communication), must be a cause for concern. It may foreshadow regional differences in mortality that are even larger than those predicted from recent trends.

- Registrar General. *Statistical review of England and Wales*. Part I: tables, medical. London: HMSO, 1968 et seq.
- Tunstall-Pedoe H, Smith WCS, Crombie IK. Level and trends of coronary heart disease mortality in Scotland compared with other countries. *Health Bull* 1986; 44: 153–61.
- Osmond C. Using age, period and cohort models to estimate future mortality rates. *Int J Epidemiol* 1985; 14: 124–9.
- Breslow NE, Day NE. *The design and analysis of cohort studies*. Lyon: IARC, 1987.
- Martin CA, Hobbs MST, Armstrong BK, de Klerk NH. Trends in the incidence of myocardial infarction in Western Australia between 1971 and 1982. *Am J Epidemiol* 1989; 129: 655–68.
- Office of Population Censuses and Surveys. *Dietary and nutritional survey of British adults*. London: HMSO, 1990.
- Cade JE, Barker DJP, Margetts BM, Morris JA. Diet and inequalities in health in three English towns. *BMJ* 1988; 296: 1359–62.
- Department of Health and Social Security. *Report on Health and Social Subjects 28. Diet and cardiovascular disease*. London: HMSO, 1984.
- Barker DJP, Osmond C. Infant mortality, childhood nutrition, and ischaemic heart disease in England and Wales. *Lancet* 1986; i: 1077–81.
- Barker DJP, Winter PD, Osmond C, Margetts BM, Simmonds SJ. Weight in infancy and death from ischaemic heart disease. *Lancet* 1989; ii: 577–80.
- Hall AJ, Barker DJP, Dangerfield PH, Osmond C, Taylor JF. Small feet and Perthes' disease: a survey in Liverpool. *J Bone Joint Surg* 1988; 70-B: 611–3.