College of Physicians is developing a common format for the everyday use of the recommended scales. A scientific network has been formed in Europe that may have a future opportunity to develop an EC-wide consensus on standard assessment scales (further details from the author).

E J DICKINSON

Royal College of Physicians Research Group
London Hospital Medical College
London E1

- 1 Research Unit of the Royal College of Physicians and British Geriatrics Society. *Standardised assessment scales for elderly people*. London: Royal College of Physicians, 1992.

- 2 Mahoney FI, Barthel DW. Functional evaluation: the Barthel Index. *Maryland State Med J* 1965; 14: 61-5.
- 3 Collin C, Wade DT, Davies S, Hovue V. The Barthel ADL Index: a reliability study. *Int Disabil Stud* 1988; 10: 61-3.
- 4 Hodkinson HM. Evaluation of a mental test score for assessment of mental impairment in the elderly. *Age Ageing* 1972; 1: 233-8.
- 5 Peach H, Green S, Locker D, *et al*. Evaluation of a postal screening questionnaire to identify the physically disabled. *Int Rehabil Med* 1980; 2: 189-93.
- 6 Yesavage JA, Brink TL, Rose TL, *et al*. Development and validation of a geriatric depression screening scale—a preliminary report. *J Psychiatr Res* 1983; 17: 37-49.
- 7 Davies B, Challis D. *Matching resources to needs in community care*. Canterbury: Personal Social Sciences Research Unit, University of Kent, 1986.
- 8 Katz S, Branch LG, Branson MH, Papsidero JA, Beck JC, Greer DS. Active life expectancy. *N Engl J Med* 1983; 309: 1218-24.

Predicting mortality from cervical cancer

Sir,—A few months ago 1990 mortality data for England and Wales were interpreted as providing some reassurance that the cervical cancer screening programme was working, though given the dearth of information about sexual behaviour, “safer sex” might also be playing a part in the reversal of trends under age 50 years.^{1 2}

A preliminary assessment of 1991 mortality data is encouraging. Population estimates revised in the light of the 1991 census are not yet available, only projections and estimates based on the 1981 census, so we have concentrated our attention on the number of deaths observed.

We recently predicted deaths over the decade 1991–2001, by modelling the death rates from 1959–88. The lowest estimate for deaths expected in 1991 was 674 for women aged 20–49 and 1329 for those aged 20–69 years.³ In fact, 461 and 1030 deaths at these ages respectively were observed (compared to 488 and 1119 in 1990), considerably lower than predicted. Although we acknowledged

that our predictions might be too high (as other forecasts have been) we are gratified they have turned out to be so much greater than the observed numbers.

It seems likely that the screening programme is finally having really noticeable effects, though without monitoring trends in sexual behaviour, we are still guessing.

M MURPHY

J CHARLTON

Unit of Health Care Epidemiology
Department of Public Health and Primary Care,
University of Oxford

- 1 Murphy M, Milne R. Trends in cervical cancer mortality. *Lancet* 1991; 338: 1081-2.

- 2 Sasieni P. *Lancet* 1991; 338: 818-9 (correspondence).

- 3 Murphy M, Osmond C. Predicting mortality from cancer of the uterine cervix from 1991–2001. *J Epidemiol Community Health* 1992; 46: 271-3.