Case-control studies in cancer patients as a surveillance system of occupational exposure in the European Community

R J Rona, N A Taub, Siri Rasmussen

Abstract

Study Objective—The main aim was to detect known relationships between lung and blood cancers and various occupational exposures (using job titles as proxies) using a case-control design. The suitability of this system for routine surveillance could then be assessed.

Design—A case-control study was carried out in 1989.

Setting—Hospitals in eight European Community countries.

Subjects—Men aged 25 to 75 years with incident and prevalent cancer of the lung (190 cases), haematopoietic system (210 cases), or gastrointestinal tract (245 controls) were studied.

Measurements and main results—The crude estimate of the overall odds ratio exposure (OR) for relevant occupational exposure of lung cancer relative to gastrointestinal cancer was 1.20 (95% confidence interval (CI) 0.82, 1.77). In a logistic regression analysis adjusting for country, age at diagnosis, smoking, and alcohol consumption, the overall OR was not greatly changed. A significant interaction of occupational exposure and age at diagnosis showed that lung cancer patients diagnosed at a younger age had a higher OR than patients diagnosed at an older age. Thus, the overall, insignificant result may have been due to a low reliability of occupational history in older age or to a selective mechanism related to age. The overall OR for occupational exposure of cancer of the blood relative to gastrointestinal cancer was 0.88 (95% CI 0.60, 1.31). The logistic regression analysis did not alter these results.

Conclusion—A surveillance based on a case-control design using job titles would not be sensitive enough to detect possible occupational risks.

information was evaluated.¹ It was concluded that the repeatability of industrial occupation information was satisfactory for research purposes (Kappa=77%). The second stage aimed to determine the utility of case-control studies using a standardised questionnaire in detecting occupational exposure related to cancer.

Doll and Peto,² in their review of avoidable risks of cancer in the US, emphasised the poor quality and limitations of current data in estimating the impact of occupational hazards on cancer deaths. They recommended a case-control design. Siemiatycki *et al*³ provided theoretical arguments for case-control studies in this context. There are two commonly used controls for this type of study, community controls⁴ and patients with another severe disease, for example another type of cancer or chronic disease.⁵ In a surveillance system the collection of information would be greatly helped by selecting hospital controls. In an overview of this subject, Smith *et al*⁶ concluded that this design has the disadvantage of possible selection bias but the advantages of minimising recall and interview bias and of reducing costs. In this phase of the study the "cases" were patients with cancers known to have some relation to occupation; the "controls" were patients with types of cancers seldom associated with occupation. We wanted to assess whether in common cancers for which approximately 10 to 15% of cases can be ascribed to industrial exposure² the appropriate association could be detected by a case-control study.

Methods and subjects

The study questionnaire was originally developed for earlier stages of the study¹ and was revised to reduce deficiencies that had become apparent. The items included were the patient's age; marital status; smoking behaviour (duration, number of cigarettes a day, and details of any recent change in behaviour); alcohol consumption; occupational and industrial history⁷; and the ICD-O coding of their disease⁸ in terms of site, morphology, behaviour, differentiation, and year of diagnosis.

The data on occupation relate to the total number of jobs that the patient has performed and to details of his most recent job, his previous job, and any earlier job lasting at least 10 years. The details requested are duration of job (in years), the EC status group, and the industrial classification for the 1981 census in Britain.⁷ The industrial classification was based on a four digit classification; the first two digits give the class and the composite four digit number gives the activity.

The questionnaire was administered to male patients aged 25-75 years. The rationale for the case-control design of the study was to ascertain

Division of
Community Health,
Department of Public
Health Medicine
United Medical and
Dental Schools, St
Thomas's Hospital,
Lambeth Palace Road,
London SE1 7EH
R J Rona
N A Taub
Siri Rasmussen
On behalf of the
European Community
Working Party whose
members are listed at the
end of this report

Correspondence to:
Dr R J Rona

Accepted for publication
November 1992

J Epidemiol Community Health 1993; 47: 320-325

A collaborative European Community (EC) study to develop a surveillance system to assess occupational and industrial risks based on case-control studies in cancer patients was carried out between 1986 and 1990. An economical surveillance system across EC countries would be helpful in detecting occupational hazards, especially in less common work activities or where the level of risk may be very low.

In the first stage of the study a new questionnaire was developed and its acceptability, repeatability, and validity of occupational

whether such a design would be robust enough to identify well established occupational hazards. The suggested number of cases in each country was 25 patients with lung cancer and 30 patients with cancer of the haematopoietic system (non-Hodgkin's lymphoma, Hodgkin's disease, or leukaemia). The controls were 35 patients with stomach, colon, or rectal cancer, diseases that have rarely been reported to be related to an occupational hazard.

The sample size for lung cancer cases was estimated by assuming that 24% of the control group would be exposed to occupations associated with lung cancer (taken from the corresponding rate among bladder cancer patients in previous stages of the project). For a relative risk of 2.0 to be detected with an 80% chance with a two-sided test at the 5% significance level, the minimum sample required is 149 subjects in each group. If eight EC countries take part in the study approximately 20 patients with lung cancer and 20 control patients in each country will be required (25 patients in each group will be required to allow for adjustments such as smoking). The sample size for patients with cancer of the haematopoietic system was based on a 16% proportion of the control group being exposed to occupations associated with blood cancers, likewise estimated from the rate in bladder cancer patients. A relative risk of 2.0 can be detected with a sample of 187 subjects in each group as the minimum sample with an 80% chance in a two sided test at a 5% significance level. If eight EC countries take part in the study, 25 patients with cancer of the blood and 25 control patients (a total of 30 patients in each group to allow for adjustments) will be required. The decision to include lung cancer patients as cases was taken despite the very strong relation between this cancer and smoking behaviour, because of the sheer number of cases and the confirmation that in industries known to be a hazard for lung cancer this risk still persists.⁹

A review of the published reports was carried out to identify the occupations that were currently reported to be related to the diseases of the cases. The occupations associated with lung cancer are listed in table I, those associated with cancers of the blood are listed in table II. The tables give the occupation, coding, and relevant references. In

Table I Significant associations between lung cancer and occupation

Occupation	Coding	References
Farmer*	0100	10
Mine worker*	2100	11, 12, 13
Steel worker*	2200	12, 14
Structural metal production	2210	15
Copper smelter	2246	16
Uranium mining, Non-ferrous ore mining, Underground miners*	2300	12, 13, 16-18
Mining and milling of crocidolite	2396	19
Stone clay*	2400	20
Asbestos goods	2440	10, 17, 21, 22
Stone cutting	2450	13
Ceramic worker	2489	13
Chemical worker*	2500	23
Foundry worker	3111	13
Shipyard/dockyard	3610	24
Bakers and pastrycooks	4136	25
Printing worker	4754	26
Painters construction, plumbers and electricians, concrete and asphalt workers†	5000	16, 17
Railway worker*	7100	27, 28
Motor vehicle driver, tracking service*	7200	8, 29
Transport equipment operator*	7700	8
Cleaning services	9230	16

*These are occupation classes, ie include all occupations starting with the two digits

†This is a division class

Table II Associations between cancer of the haematopoietic system and occupation

Occupation	Coding	References
<i>Leukaemia</i>		
Agriculture*	100	30
Manufacture of product of coal	1115	30
Petroleum products	2512	30, 31
Photographic material and chemicals	2591	31
Chemical co workers	2599	31
Manufacturing*	3100-3700	32
Tobacco manufacture	4290	30
Shoe-leather: (i) Leather goods	4420	31
(ii) Footwear	4510	31
Rubber worker	4812	31
Construction*	5000	32
Sales*	6400-6600	32
Utilities and communication*	7100-7900	32
Finance, insurance, and real estate*	8100-8300	32
Research*	9400	30
<i>Hodgkin's disease</i>		
Farmers*	0100	33, 34
Fisherman*	0300	34
Cementer	2420	34
Glass	2478-2479	30, 35
Pottery	2489	35
Chemical processing	2500	35
Dyer	2516	34
Chemist sprayer	2551	35
Photography	2591	34
Spinning, weaving	4321	30
Leather worker	4410	34
Leather goods	4420	34
Woodmakers	4600	34, 36
Carpenter	4630	37
Plastic worker	4836	34
<i>Non-Hodgkin's lymphoma</i>		
Agriculture*	0100	38
Forestry*	0200	33
Waterworks and supply*	1700	30
Manufacturing of non-metallic products*	2400	30
Benzene	2512	34
Canning, preserving and processing of fish etc.	4115-4150	33
Leather worker*	4400	34
Fencing work	4630	38

*These are occupation classes, ie include occupations starting with the two digits

relation to lung cancer, we included only those references that were based on data from industrially developed countries and which quoted a confidence interval for the estimated risk. With regard to cancer of the blood, we relaxed the requirement that the relevant report should quote a confidence interval for the risk, since there are fewer relevant papers on cancer of the blood than on lung cancer. Proportional mortality ratio studies were excluded. These can be misleading as differences in mortality from other cancers may influence the association between exposure and the relevant disease.

In the statistical analyses, subjects were considered exposed to a high risk occupation if they had worked in any of the relevant occupations for at least three years (see tables I and II). Odds ratios (ORs) were used to estimate the association of occupational exposure with lung cancer and with blood cancer, using gastrointestinal cancer patients as controls. Where there are (a) exposed cases, (b) unexposed cases, (c) exposed controls, and (d) unexposed controls, the crude, unadjusted estimate of the odds ratio is given by $OR = \frac{ad}{bc}$.

Confidence intervals were calculated using the formula:

$$\text{Var} [\log_e(OR)] = \frac{1}{a} + \frac{1}{b} + \frac{1}{c} + \frac{1}{d}$$

and the assumption that $\log_e(OR)$ is normally distributed. Logistic regression models³⁹ were

developed to adjust for the suspected confounding variables: age at diagnosis, country, amount of smoking, alcohol consumption, and the dates of diagnosis and of interview (to distinguish between prevalent and incidence cases). The variables for the analysis of lung cancer were divided into groups of approximately equal size as follows:

Age at diagnosis: 25–56 years, 57–61 years, 62–66 years, and 67 years or over;

Length between time of diagnosis and interview: up to 36 months or more than 36 months;

Length of time smoking: 0 years, 1–34 years, 35–44 years and more than 44 years;

Cigarette consumption per day: 0, 1–19, 20–24, and more than 24;

Alcohol consumption (g/week): 0, 1–120, 121–299, and more than 299;

Age at time of survey: 25–56 years, 57–61 years, 62–66 years, and more than 66 years;

EC job status grouped into eight categories as a proxy for socioeconomic group;

Which of the eight EC countries.

The exposure variable was divided into less than 36 months or at least 36 months in the relevant high risk occupations.

The variables for the analysis of blood cancer were divided in a similar manner except that for age at diagnosis the classes were 21–49 years, 50–58 years, 59–64 years, and more than 64 years, since patients with this cancer were younger than for lung cancer.

Results

DESCRIPTION OF THE SAMPLE

Table III gives the distribution of patients in relation to type of cancer and country after excluding 41 patients who did not satisfy the entry criteria. The age distributions for patients with lung cancer and gastrointestinal cancer were very similar both between countries and between these two types of cancer. Patients with cancer of the blood were on average younger than patients in

the other two groups. In general, patients with Hodgkin's disease were the youngest (mean age 46.8 years), while patients with lymphoma and leukaemia had mean ages of 54.0 and 53.4 years respectively. In terms of cancer site, the distributions of patients with a gastrointestinal tract cancer were as follows: stomach (n=49), small intestine (n=1), colon (n=102), and rectum or rectosigmoid junction (n=77). Belgian patients could not be included in the distributions by site as they were given a diagnosis without further classification according to site or morphology.

The types of cancer of the blood were: lymphomas not otherwise specified or diffuse (n=57), lymphomas (nodular or follicular) (n=23), Burkitt's lymphoma (n=1), leukaemias (n=55), reticulosarcoma (n=5), and Hodgkin's disease (n=41).

Most patients were diagnosed in the last year of the study. For gastrointestinal cancer 114 patients were diagnosed less than a year before interview, 68 between one and three years before interview, and 40 more than three years before interview. Similar figures for lung cancer were 128, 11, and 19 respectively and for cancer of the blood 84, 40, and 49 respectively. This information was not available for 92 patients, most of whom were from Belgium.

For each country in the study, both the length of time smoking and the number of cigarettes smoked were higher in patients with lung cancer than in patients with gastrointestinal cancer or cancer of the blood. When patients who smoked pipes or cigars were included, only one patient in the lung cancer group was a non-smoker. Smoking was also highly prevalent in the group of patients with cancer of the gastrointestinal tract. Even in patients with cancer of the blood, the observed percentage of non-smokers was low in comparison with that often reported in the population.

Patients with lung cancer had on average a higher level of alcohol intake than patients with other types of cancer. The variation in alcohol consumption between countries was very large, ranging from 88 g per week in Britain to 328 g per week in Germany.

Table IV gives the number of years accounted for by the three jobs that patients were asked to describe—that is the last job, previous job, and earlier jobs lasting more than 10 years. These occupations together represented approximately 93% of the total time that patients had spent in employment. On average, the jobs represented approximately 35 years' work for patients with gastrointestinal cancer or cancer of the lung and 30 years for patients with blood cancer. The SDs were, however, very large, indicating that for some patients the number of years in any employment was low.

CASE-CONTROL ANALYSIS

Lung cancer (cases) and gastrointestinal tract cancer (controls)

Table V gives the frequencies of cases and of controls in relation to country and exposure category. The overall unadjusted estimate of OR for occupational exposure was not significantly different from 1, OR=1.20 (95% CI 0.82, 1.77). Although the unadjusted ORs varied widely in

Table III Distribution of patients in relation to mean (SD) age (years), type of cancer, and country

Country	Gastrointestinal cancer		Lung cancer		Blood cancer	
	No	Mean (SD)	No	Mean (SD)	No	Mean (SD)
Belgium	16	59.8 (10.1)	25	61.4 (7.4)	28	57.3 (10.9)
Denmark	33	59.8 (6.8)	21	60.9 (6.4)	25	50.1 (15.1)
France	35	61.1 (8.1)	24	59.5 (7.3)	25	44.0 (13.9)
Germany	35	58.9 (9.9)	25	61.1 (6.1)	27	53.5 (13.2)
Ireland	21	56.0 (13.3)	19	60.4 (8.9)	19	51.5 (14.5)
Italy	33	58.7 (7.4)	28	58.1 (8.1)	31	52.0 (11.9)
United Kingdom	36	62.2 (8.9)	25	60.9 (7.7)	31	57.2 (11.2)
Spain	36	61.4 (8.3)	23	64.8 (6.8)	24	55.6 (13.9)
Total	245	60.0 (9.0)	190	60.8 (7.5)	210	52.8 (13.4)

Table IV Mean (SD) numbers of years accounted for in the occupational history explored in the study

Country	Gastrointestinal cancer		Lung cancer		Blood cancer	
	No	Mean (SD)	No	Mean (SD)	No	Mean (SD)
Belgium	16	34.4 (28.5)	25	36.5 (11.5)	28	36.6 (21.4)
Denmark	33	34.9 (8.4)	21	36.5 (9.7)	25	27.4 (15.9)
France	35	32.0 (11.3)	24	28.9 (11.6)	25	20.4 (13.3)
Germany	35	33.0 (9.9)	25	37.9 (8.0)	27	26.1 (13.9)
Ireland	21	31.4 (12.5)	19	28.8 (16.4)	19	28.1 (17.0)
Italy	33	38.8 (8.2)	28	36.6 (10.3)	31	30.7 (11.3)
United Kingdom	36	36.3 (10.8)	25	34.0 (11.4)	31	33.2 (9.8)
Spain	36	36.1 (10.4)	23	43.2 (8.4)	24	34.3 (13.3)
Total	245	34.8 (12.2)	190	35.5 (11.6)	210	29.8 (15.3)

Table V Numbers occupationally exposed and unexposed for cases and controls in relation to country for lung cancer (cases) and gastrointestinal tract cancer (controls)

Country	Lung cancer		Gastrointestinal tract cancer		Total	Odds ratio
	Exposed cases a	Unexposed cases b	Exposed controls c	Unexposed controls d		
Belgium	6	19	6	10	41	0.53
Denmark	8	13	14	19	54	0.84
France	9	15	12	23	59	1.15
Germany	12	13	9	26	60	2.67
Ireland	9	10	10	11	40	0.99
Italy	15	13	14	19	61	1.57
United Kingdom	13	12	17	19	61	1.21
Spain	11	12	14	22	59	1.44
Overall	83	107	96	149	435	1.20 (95% CI 0.82, 1.77)

Table VI Logistic regression model for lung cancer and gastrointestinal cancer: estimated coefficients, *b* of smoking behaviour

	No	<i>b</i>	<i>se(b)</i>	Odds ratio	95% CI
Time smoking (y):					
Non-smokers	39	0.00	—	1.00	—
1–34	133	2.45	1.16	11.54	1.20, 111.67
35–44	129	2.93	1.16	18.70	1.94, 179.43
45+	134	3.80	1.17	44.70	4.54, 440.68
Daily cigarette consumption:					
Non-smokers	62	0.00	—	1.00	—
1–19	99	0.03	0.57	1.03	0.33, 3.16
20–24	123	0.61	0.56	1.84	0.61, 5.49
25+	151	1.02	0.56	2.78	0.93, 8.26

individual countries, from 0.53 (Belgium) to 2.67 (Germany), none were significantly different from 1 ($p > 0.05$).

Logistic regression analysis was carried out to adjust the OR for the potentially confounding variables, giving the estimate $OR = 0.99$ (95% CI 0.63, 1.58). The only independent factors that were significantly associated with case-control status were: duration of smoking ($p < 0.001$), cigarette consumption ($p < 0.01$), and job status ($p < 0.05$) (table VI). Two-way interaction terms between smoking and drinking were then included in the model and retained, although their overall significance was marginal ($p = 0.05$).

The presence of any effect modifications was now assessed by backwards stepwise deletion of the interaction terms between each independent factor and occupational exposure. The only factor found to modify the association of high risk jobs with lung cancer was the patient's age at diagnosis. Although this effect is not highly significant ($p = 0.03$), a gradient where occupational exposure seems to increase the risk for younger men but to decrease the risk for older men was shown (table VII).

Blood cancer (cases) and gastrointestinal tract cancer (controls)

The overall unadjusted estimate of OR was not significantly different from 1 ($OR = 0.88$ (95% CI 0.60, 1.31)). As with lung cancer, the variation in unadjusted ORs in different countries was wide but was not statistically significant (table VIII).

In the logistic regression analysis, adjustment was made for the same set of potentially confounding variables as for the previous analysis of lung cancer except for time smoking because this variable would be strongly associated with age.

Table VII Logistic regression model for lung cancer and gastrointestinal cancer: age specific odds ratios of occupational exposure

Age (y)	No	Odds ratio	95% CI
22–56	132	1.76	0.73, 4.22
57–61	95	1.58	0.58, 4.30
62–66	126	0.73	0.29, 1.82
67+	82	0.25	0.08, 0.78

The only significant confounder in the model was age at diagnosis ($p < 0.001$) (table IX). Interaction terms between smoking and drinking were then included in the model, although they were not significant ($p > 0.05$).

The search for effect modification did not identify any factor that interacted with occupational exposure.

Discussion

The time available for recruiting patients into the study was only four months. Most countries complied with the number of patients suggested in the protocol. Although 41 patients had to be excluded from the analysis because they did not fulfil the protocol requirements, some departures from the rules for age and diagnosis were permitted. Thus, reticulosarcoma was included within the definition of lymphoma. After excluding the patients that did not fulfil the requirements, the number of patients for the analysis was 90% of the target number.

In general, the percentage of non-smokers was very low for all types of cancers, although, as expected, patients with a lung cancer had higher tobacco consumption levels. In terms of alcohol intake, there were very large differences between countries. There are several sources of artefact: the wide variation in alcohol content of beverages, the variation between countries in terms of the size of glass used commercially and in the home, and the conversion of beverage consumption (as given in the questionnaire) to that of total alcohol content. We minimised the second and third types of artefact by allowing for different sizes of glasses and by standardising for alcohol content by beverage.

The occupational history explored in this study accounted for around 90% of the participants' total time in employment. The reasons for choosing lung and blood cancers as the case diseases in the study were: (i) they were sufficiently common, (ii) several reports of their relation with occupation were available, and (iii) they had contrasting characteristics. Lung cancer is a fairly homogeneous entity characterised by short survival and it is strongly related to tobacco consumption. Cancer of the blood consist of a very large number of disorders, each of them low in frequency and with a survival time that can be large. A cure can be achieved for some of these blood disorders, such as Hodgkin's disease.

The controls used in this study include both an aggressive type of cancer, such as stomach cancer, and a cancer with a fairly good survival such as cancer of the colon. In general, the published reports available have failed to find many occupations related to this type of cancer.

The results from the case-control analysis of lung cancer and gastrointestinal cancer patients indicated no overall risk as a result of occupational exposure, which indicates that this type of design is not sufficiently sensitive for a surveillance system. There was a significant interaction between occupational exposure and age at diagnosis; the OR for the younger patients was higher than that for the older patients. There are several possible explanations—chance finding, artefact, a selective factor, or a real difference due to some change in

Table VIII Crude data for blood cancer (cases) and gastrointestinal tract cancers (controls)

Country	Exposed cases a	Unexposed cases b	Exposed controls c	Unexposed controls d	No	Odds ratio
Belgium	10	18	7	9	44	0.71
Denmark	17	8	27	6	58	0.47
France	20	5	22	13	60	2.36
Germany	21	6	26	9	62	1.21
Ireland	18	1	15	6	40	7.20
Italy	13	18	10	23	64	1.66
United Kingdom	19	12	30	6	67	0.32
Spain	17	7	27	9	60	0.81
Overall	135	75	164	81	455	0.88 (95% CI 0.60, 1.31)

Table IX Logistic regression model for blood cancer and gastrointestinal cancer estimated coefficients, b, of age effect

Age at diagnosis (y)	No	b	se(b)	Odds ratio	95% CI
<50	120	0.00	—	1.00	—
50–58	117	–1.23	0.29	0.29	0.16, 0.52
59–64	108	–1.81	0.31	0.16	0.09, 0.30
64+	110	–1.47	0.31	0.23	0.13, 0.42

the nature of occupational exposure over time. It seems that differential reliability and selection over time are the most plausible explanations for this finding. The differential reliability of occupational history (if it were shown in other studies) would suggest that patients in this type of study be restricted to a narrow age range, although this would increase the difficulty of recruitment. This may be a valid finding for lung cancer and should not therefore affect the general design of surveillance.

The case-control analysis of blood cancers and gastrointestinal cancers did not show a significant overall risk due to occupational exposure either.

The study has been helpful in identifying a number of problems with this sort of surveillance system. It was a great advantage that in this study we could group together the many occupations already known to be associated with a type of cancer. In a full surveillance system of occupational exposure, however, each separate occupation will have to be tested for an association with a disease. The diversity of occupations in the modern world means that a very large number of patients will be required.

In this study the control group was mixed and would be expected to be a reasonable group to choose for this purpose. However, there may be two caveats. Some of these cancers are associated with lower socioeconomic status groups and would include a higher proportion of patients in semiskilled and unskilled occupations than would, say, cancers of the blood. We checked this possibility by including EC job status (as a proxy for socioeconomic background) in the logistic modelling. In a surveillance programme there may be an advantage in using an additional control group which is not known to be associated with social status.

The case-control study related to blood cancers illustrated another possible limitation of this sort of system. Patients with cancer of the blood are younger than patients with other types of cancer. We have adjusted for age in our analysis but other characteristics that are more prevalent in younger than in older patients, or vice versa, might have escaped adjustment.

In conclusion, the study has shown that the power of a surveillance system for occupational exposure based on a case-control design would be

rather low and, if implemented, it should include a very large number of collaborating institutions in each country. We fear that many of the true associations will be missed because of the very large number of occupations in modern societies. One of the possible problems, as pointed out by Siemiatycki *et al.*³ is that job title covers a group of workers whose activities and exposures are heterogeneous. A risk may go undetected because only a subgroup in a job title may have been exposed to a carcinogen.

Members of the working party were as follows: Belgium—Dr M Kornitzer (Laboratoire d'Epidemiologie et de Medicine Sociale, Universite Libre de Bruxelles), Dr P De Vuyst (Hopital Erasme, Universite Libre de Bruxelles). Denmark—Dr J Mosbech (Chairman) (National Board of Health), Dr M Anderson (Institute of Cancer Epidemiology). France—Professor R Pariente (Clinique Pneumologique de l'Hôpital Beaujon), Dr F Neukirch (Laboratoire de Physiopathologie Respiratoire, Hôpital Beaujon). Germany—Dr C Duting (Institut für Soziale Medizin, Freie Universität Berlin), Professor Gustav Schake (Institute of Occupational medicine, Freie Universität Berlin). Ireland—Dr P Kirke (The Health Research Board), Dr B Herity (Department of Community Medicine and Epidemiology, University College). Italy—Professor S Marinoni (Istituto de Scienze Sanitarie Applicate, Università degli Studi di Pavia), Dr L Casali (Presidio Pneumotisiologico). Netherlands—Dr D J van Puter (Nederlands Instituut voor Praeventieve Gezondheidszorg TNO), Dr H P A van de Walter (Nederlands Instituut voor Praeventieve Gezondheidszorg TNO). Spain—Dr S Moncada (Centre de Salut Laboral, Institut Municipal de Salut), Dr A Arias (Servei d'Informacioni Sanitaria Institut Municipal de la Salut). United Kingdom—Professor W W Holland (Department of Public Health Medicine, UMDS), Dr M Farebrother (Medway Hospital); Dr R J Rona (Department of Public Health Medicine, UMDS).

- 1 Rona RJ, Mosbech J. Validity and repeatability of self-reported occupational and industrial history from patients in EEC countries. *Int J Epidemiol* 1989; 18: 674–9.
- 2 Doll R, Peto R. The causes of cancer. Quantitative estimates of avoidable risks of cancer in the United States today. *J Natl Cancer Inst* 1981; 66: 1191–308.
- 3 Siemiatycki J, Nicholas ED, Fabry J, Cooper JA. Discovering carcinogens in the occupational environment: a novel epidemiologic approach. *J Natl Cancer Inst* 1981; 66: 217–25.
- 4 Hoover RN, Hartge Strasser P. Artificial sweeteners and human bladder cancer. *Lancet* 1980; i: 837–840.
- 5 Acheson ED, Cowdell RH, Rang E. Adenocarcinoma of the nasal cavity and sinuses in England and Wales. *Br J Ind Med* 1972; 29: 21–30.
- 6 Smith AH, Pearce NE, Callas PW. Cancer case-control studies with other cancers as controls. *Int J Epidemiol* 1988; 17: 298–306.
- 7 Office of Population Censuses and Surveys. *Classification of occupations* 1980. London: HMSO, 1980.
- 8 World Health Organisation. *International classification of disease for oncology*. ICD-O. Geneva: WHO, 1976.
- 9 Benhamou S, Benhamou E, Flamant R. Occupational risk factors of lung cancer in a French case-control study. *Br J Ind Med* 1988; 45: 231–3.
- 10 McDuffie HH, Klaassen DJ, Cockcroft DW, Dosman JA. Farming and exposure to chemicals in male lung cancer patients and their siblings. *J Occup Med* 1988; 30: 55–9.
- 11 Dave SK, Edling C, Jacobson P, Axelsson O. Occupation, smoking and lung cancer. *Br J Ind Med* 1988; 45: 790–2.
- 12 Lerchen ML, Wiggins CL, Samet JM. Lung cancer and occupation in New Mexico. *J Natl Cancer Inst* 1987; 79: 639–45.
- 13 Lynge E, Kurppa K, Kristofersen L, Malker H, Sauli H. Silica dust and lung cancer: results from the Nordic occupational mortality and cancer incidence registers. *J Natl Cancer Inst* 1986; 77: 883–9.
- 14 Beaumont JJ, Leveton K, Knox K, *et al.* Lung cancer mortality in workers exposed to sulfuric acid mist and other acid mists. *J Natl Cancer Inst* 1987; 79: 911–21.
- 15 Ronco G, Ciccone G, Mirabelli D, Troia B, Vineis P. Occupation and lung cancer in two industrialized areas of northern Italy. *Int J Cancer* 1988; 41: 345–58.
- 16 Damber LA, Larsson LG. Occupation and male lung cancer: a case-control study in northern Sweden. *Br J Ind Med* 1987; 44: 446–53.

- 17 Damber LA, Larsson LG. Underground mining, smoking and lung cancer: a case-control study in the iron ore municipalities of northern Sweden. *J Natl Cancer Inst* 1985; 74: 1207–13.
- 18 Samet JM, Pathak DER, Morgan MV, Marbury MC, Key CR *et al*. Radon progeny exposure and lung cancer risk in New Mexico uranium miners: a case-control study. *Health Physics* 1989; 56: 415–21.
- 19 Armstrong BK, De Klerk NH, Musk AW, Hobbs MST. Mortality in miners and millers of crocidolite in Western Australia. *Br J Ind Med* 1988; 45: 5–13.
- 20 Buiatti E, Kriebel D, Geddes M, Santucci M, Pucci N. A case control study of lung cancer in Florence, Italy. I Occupational risk factors. *J Epidemiol Community Health* 1985; 39: 244–50.
- 21 Gardner MJ, Powell CA, Gardner A, Winter PD, Fletcher AC. Continuing high lung cancer mortality among ex-amosite asbestos factory workers and a pilot study of individual anti-smoking advice. *J Soc Occup Med* 1988; 38: 69–72.
- 22 Vineis P, Thomas T, Hayes RB, *et al*. Proportion of lung cancers in males, due to occupation, in different areas of the USA. *Int J Cancer* 1988; 42: 851–6.
- 23 Bond GG, Flores GH, Shellenberger RJ, Cartmill JB, Fishbeck WA, Cook RR. Nested case-control study of lung cancer among chemical workers. *Am J Epidemiol* 1986; 124: 53–66.
- 24 Sanden A, Naslund P, Jarvholm B. Mortality in lung and gastro-intestinal cancer among shipyard workers. *Int Arch Occup Environ Health* 1985; 55: 277–83.
- 25 Wicksell L, Carstensen JM, Eklund G, Gustafsson JA. Lung cancer incidence among Swedish bakers and pastry cooks: geographical variation. *Scand J Soc Med* 1988; 16: 183–6.
- 26 Schoenberg JB, Stenmhagen A, Mason TJ, Patterson J, Bill J, Altman R. Occupation and lung cancer risk among New Jersey white males. *J Natl Cancer Inst* 1987; 79: 13–21.
- 27 Garshick E, Schenker MB, Munoz A, *et al*. A retrospective cohort study of lung cancer and diesel exhaust exposure in railroad workers. *Am Rev Respir Dis* 1988; 137: 820–5.
- 28 Garshick E, Schenker MB, Munoz A, *et al*. A case-control study of lung cancer and diesel exhaust exposure in railroad workers. *Am Rev Respir Dis* 1987; 135: 1242–8.
- 29 Damber L, Larsson LG. Professional driving, smoking, and lung cancer: a case referent study. *Br J Ind Med* 1985; 42: 246–52.
- 30 Olsen J, Jensen OM. Occupation and risk of cancer in Denmark. *Scand J Work Environ Health* 1987; 13(suppl 1): 1–19.
- 31 Linet MS. *The leukaemias: epidemiological aspects*. New York. Oxford University Press, 1985.
- 32 Williams RR, Stegens NL, Goldsmith JR. Associations of cancer site and type with occupation and industry from the third national cancer interviews. *J Natl Cancer Inst* 1977; 59: 1143–85.
- 33 Dubrow R, Poulson JO, Indian RW. Farming and malignant lymphoma in Hancock Country, Ohio. *Br J Ind Med* 1989; 45: 25–8.
- 34 Vianna NL, Polan A. Lymphomas and occupational benzene exposure. *Lancet* 1979; i: 1394–5.
- 35 Olsson H, Brandt L. Occupational handling of chemicals preceding Hodgkin's disease in men. *Br J Ind Med* 1979; 279: 580–1.
- 36 Fonte R, Grigis L, Grigis P, Franco G. Chemicals and Hodgkin's disease. *Lancet* 1982; ii: 50.
- 37 Brownson RC, Reif JS. A cancer registry-based study of occupational risk for lymphoma, multiple myeloma and leukaemia. *Int J Epidemiol* 1988; 17: 27–32.
- 38 Pearce NE, Smith AH, Fisher DO. Malignant lymphoma and multiple myeloma linked with agricultural occupations in a New Zealand cancer registry-based study. *Am J Epidemiol* 1985; 121: 225–37.
- 39 Baker RJ, Nelder JA. *Generalised linear interactive modelling (GLIM) system*. Release 3. Oxford: Oxford Numerical Algorithms Group, 1978.