late 1970s and is now rising. After allowing for the proportion of births to mothers over 35 years, the trend with year remains significant at the 5% level, $OR=0.9996$, (95%CI: 0.9995, 0.9997). Over the period there has also been a steady increase in the proportion of single mothers. As single mothers are more likely to have boys,⁴ it is likely that the decline in the sex ratio of children born to other women is even greater.

This decline in the sex ratio remains unexplained. Possibilities are a general fall in the frequency of intercourse, perhaps due to the increasing stress of society or changes in male or female hormone levels. The latter could be consequence of air pollution which was shown by Williams *et al*⁵ to be associated with a lower sex ratio.

H O DICKINSON
L PARKER

University of Newcastle
Sir James Spence Institute of Child Health
Royal Victoria Infirmary
Newcastle upon Tyne

This research was supported by grants from the United Kingdom Coordinating Committee on Cancer Research, the North of England Children's Cancer Research Fund and Westlakes Research Institute.

- 1 Harrison PTC, Humphrey CDN, Litchfield M, Peakall D, Schuker LK. *Environmental oestrogens: consequences to human health and wildlife*. Leicester: MRC Institute for Environment and Health, University of Leicester, 1995.
- 2 James WH. The human sex ratio. Part 2: A hypothesis and a program of research. *Hum Biol* 1987;59:873-900.
- 3 James WH, Rostrom J. Parental age, parity and sex ratio in births in England and Wales 1968-1977. *Journal of Biosocial Science* 1985;17:47-56.
- 4 James WH. The human sex ratio. Part 1: A review of the literature. *Hum Biol* 1987;59:721-52.
- 5 Williams FL, Lawson AB, Lloyd OL. Low sex ratios of births in areas at risk from air pollution from incinerators, as shown by geographical analysis and 3-dimensional mapping. *Int J Epidemiol* 1992;21:311-19.

Comparing measures of variation

Sir - We recently reported levels of geographical variation in hospital admission rates in the Oxford region.¹ In discussing our results, we briefly compared systematic component of variation (SCV) values from our data with those published by Wennberg *et al*² for hospital service areas in Maine, USA. Wennberg *et al* described their method by citing an earlier paper in which SCVs had been calculated using a multiplication constant of 100.³ We now believe that in the 1984 study² they in fact used a multiplication factor of 1000, although it is impossible to discover this from their paper.

The reported differences between our results and those from Maine persist, however, after taking account of the 10-fold multiplication factor. Only 10% of admissions in Maine were for conditions with an SCV (100)

of less than 5.0, compared with 44% of surgical workload in Oxford.

Our conclusion remains unchanged. There was substantially less variation in admission rates in the Oxford region than in Maine, USA.

JOHN NEWTON
VALERIE SEAGROATT
MICHAEL GOLDACRE
Unit of Health-Care Epidemiology,
Department of Public Health,
and Primary Care,
University of Oxford.

- 1 Newton JN, Seagroatt V, Goldacre M. Geographical variation in hospital admission rates: an analysis of workload in the Oxford region, England. *J Epidemiol Community Health* 1994;48:590-95.
- 2 Wennberg JE, McPherson K, Caper P. Will payment based on diagnosis-related groups control hospital costs? *N Engl J Med* 1984;311:295-300.
- 3 McPherson K, Wennberg JE, Horind OB, Clifford P. Small-area variations in the use of common surgical procedures: an international comparison of New England, England, and Norway. *N Engl J Med* 1982;307:1310-14.

Perinatal mortality in a first generation immigrant population and its relation to unemployment in The Netherlands

Sir - In the introduction of the above article¹ we stated in error that a previous study by Doornbos and Nordbeck of the same population² showed an odds ratio of 1.50 for perinatal mortality for infants of Surinam origin and of 1.42 for infants of other non-Dutch origin. In fact, these authors reported crude odds ratios of 1.23 and 1.22 for the two groups. The error resulted from a misreading of data provided by these authors.

Our re-analysis of the role of various factors associated with perinatal mortality and ethnic origin therefore confirms the simpler Doornbos/Nordbeck analysis with respect to the marginal role of infant origin itself.

The main finding of our report regarding the important role of parental employment status as a predictor of perinatal mortality is not affected.

L H LUMEY
S A REIJNEVELD
American Health Foundation
New York
USA

- 1 Lumeij LH, Reijneveld SA. Perinatal mortality in a first generation immigrant population and its relation to unemployment in The Netherlands. *J Epidemiol Community Health* 1995;49:434-9.
- 2 Doornbos JPR, Nordbeck HJ. *Perinatal mortality. Obstetric risk factors in a community of mixed ethnic origin in Amsterdam* (proefschrift). Amsterdam: Universiteit of Amsterdam, 1985.

NOTICES

European Journal of Public Health (1996;6)

The contents of the *European Journal of Public Health* for June 1996 are as follows.

Looking back and ahead: a bright view for the journal (ed) Palm J, Svensson P-G.

ORIGINAL ARTICLES:

Sharing syringes in Madrid: a social phenomenon Rodrigues-Arenas MA, Zunzunegui Pastor MV, Friedman SR, *et al*.

Prevalence and trends of cigarette smoking in different occupational groups: results of the Minnesota heart survey 1980-1982 and 1985-1987 Knutsson A, Luepker RV, Sprafka JM, Virnig B.

The effectiveness of treatment for the prolapsed lumbar intervertebral disc: a review of the literature Deane M, Moore AJ, Long AF, Harrison S.

Working in the field: the psychiatric nurse as a libero in the community health services Hummelvoll JK.

Proposals for collaboration in European public health training Köhler L, Bury JA, de Leeuw E, Vaughan P.

Health expectancy in the Netherlands 1983-1990 van de Water HPA, Boshuizen HC, Perenboom RJM.

Functional ability scales for the elderly. A validation study Avlund K, Kreiner S, Schultz-Larsen K.

Detection of depressive complaints in children Kroesbergen HT, de Wit C, Stijnen T.

The French breast cancer screening programme: epidemiological and economic results of the first round of screening Wait SH, Allemand HM.

A study of general practitioners' prescribing behaviour to the elderly in Wallonia, Belgium Taziaux P, Franck J, Ludovicy R, Albert A.

The incidence of hypertension and the effects of a regular high-dose alcohol intake on blood pressure in a medium-sized Hungarian town (Csonhrád) Mohácsi G, Abrahám G, Remes P, Lovas S, Sonkodi S.

Uncertainty and lack of trust with Parkinson's disease Nijhof G.

Book reviews, Calendar of Events, EIPHA pages