Figure 1 Age specific mortality rates in relation to selected cohorts in Taiwan. The 1941 curve is close to the 1951 curve. It is excluded to avoid visual confusion. The legends indicate the mid-points of the five year cohorts.

post war cohorts have an almost identical mortality level in the age range 15–29, but the 1961 cohort has much better child mortality level. Due to the lack of an interaction term, the model would overestimate child mortality in the 1961 cohort and underestimate its adult mortality level in relation to the 1951 cohort's. Note that child mortality is more heavily weighted in the calculation of life expectancy since it affects the number of person-years lived in all subsequent age intervals. In contrast adult and old age mortality affect fewer intervals. Therefore, the estimates of life expectancy at birth based on the age-cohort model may be biased. We should not read too much into figure 1 because the data are scanty. Nor should we overlook it because it is consistent with our epidemiological knowledge.

Obviously the age-cohort model cannot accommodate an interaction term because of the missing data for some cells. Therefore, its use should be limited to the age ranges and/or cohort ranges for which we expect the assumption of constant age pattern to hold. For instance, we can apply it to the estimation of life expectancy at age 60 with reasonable confidence. Of course, the assumption must be empirically checked whenever possible.

YIN BUN CHEUNG
Institute for Human Services Research,
PO Box 73815, Kowloon Central Post Office,
Hong Kong

- 1 Lee WC, Hsieh RL. Estimating life expectancy using an age-cohort model in Taiwan. *J Epidemiol Community Health* 1996;50:214–7.
- 2 Clayton D, Schifflers E. Models for temporal variation in cancer rates. I: age-period and age-cohort models. *Stat Med* 1987;6:449–67.
- 3 Philips MA, Feachem RG, Murray CJ, Over M, Kjellstrom T. Adult health: a legitimate concern for developing countries. *Am J Public Health* 1980;70:1527–30.

Reply

Dr Cheung points out correctly that the age-cohort model we used to estimate cohort life expectancy in Taiwan¹ has an implicit assumption of constant age pattern across cohorts. Dr Cheung also explains why such

an assumption is unlikely to be met, especially in a population under epidemiologic transition. I fully agree with his views.

In the context of age-period-cohort analysis, the problem of non-parallelism in age patterns has been explicitly addressed.^{2,3} Unfortunately, the proposed methodologies are a bit too complex for the present purpose. Also, simple addition of an interaction term to the age-cohort model is not feasible because of unavoidable missing cells from age-by-cohort cross tabulation. Instead of resorting to models with interaction terms, I feel that a practical solution to this problem may be to perform a "separate analysis". For example, we can break down the data by age into, say, "childhood", "adulthood", and "elderly" group, each modelled separately, and then the results combined. By this method, different segments of the population assume different cohort trends. Alternatively, we can break down the all-cause data into their component causes, modelled separately and then combined – an approach suggested in our paper.¹ Clearly, such an approach pays due respect to the different roles of communicable and the non-communicable diseases during the epidemiologic transition. However, the benefits we gain from these more "sophisticated" approaches must be weighed against what we can infer from "crude" but "simple" age-cohort modelling as shown in our paper.¹

Finally, Dr Cheung has stated that the assumption of parallelism is not met in our data after the 1961 cohort. I wish to point out that we also refute the results after 1961, though for a different reason.

WEN-CHUNG LEE
Graduate Institute of Epidemiology,
National Taiwan University,
Taiwan

- 1 Lee WC, Hsieh RL. Estimating life expectancy using an age-cohort model in Taiwan. *J Epidemiol Community Health* 1996;50:214–7.
- 2 James IR, Segal MR. On a method of mortality analysis incorporating age-year interaction, with application to prostate cancer mortality. *Biometrics* 1982;38:433–43.
- 3 Lee WC, Lin RS. Analysis of cancer rates using excess risk age-period-cohort models. *Int J Epidemiol* 1995;24:671–77.

Tuberculosis among homeless people at a temporary shelter in London

SIR – Each year the charity Crisis sets up temporary Christmas shelters, not for "the homeless" but for homeless people. Kumar *et al* use the former term on 23 occasions in their report of a chest x ray screening programme.¹

Issues of concern were the high prevalence of tuberculosis and suboptimal management of known and presumed cases due to patient "default". These problems have been previously documented in a very similar group of homeless people in London during the mid 1980s.² In the earlier survey, there was a notable association between tuberculosis and alcohol (ab)use,² which may have an impact on treatment compliance. Data from the Crisis Open Christmas¹ could perhaps be used to ascertain whether there was an association between "loss to follow up" and accommodation status of "no fixed abode" or self reported "regular alcohol consumption".

The documented health care needs of single homeless, or homeless and roofless people are multiple.^{3,4} The prevalence of mental illness noted by Kumar *et al*,¹ low by comparison with other survey data,^{4,7} may have been significantly underestimated by selection or reporting bias.

A comparably wide range of accessible and acceptable health care provision therefore seems appropriate and is, indeed, a feature of most dedicated services for homeless people.^{8,9} Recommendations to improve current provision, based on specialist outreach respiratory/tuberculosis services, will have limited capacity to address important and potentially confounding comorbidity.

The most effective means of delivering comprehensive health care to homeless people, including case finding and management of those with tuberculosis and their contacts, needs formal evaluation. However, the views of professionals in primary and community health services and in the voluntary sector, who best understand the motivating factors and influential social networks of homeless people, should not be overlooked. Indeed, the suggestion of patient held records was recently raised in *The Big Issue*.¹⁰

MAGGIE HARDING
NIKKI BROWN
West London Initiative on Single
Homelessness (WISH),
17 Whitton Dene, Hounslow,
Middlesex TW3 2JN

- 1 Kumar D, Citron KM, Leese J, Watson JM. Tuberculosis among the homeless at a temporary shelter in London: report of a chest x ray screening programme. *J Epidemiol Community Health* 1995;49:629–33.
- 2 Ramsden SS, Baur S, El Kabir DJ. Tuberculosis among the central London single homeless. *J R Coll Physicians* 1988;22:16–17.
- 3 Shanks NJ. Medical morbidity of the homeless. *J Epidemiol Community Health* 1988;42:183–86.
- 4 George SL, Shanks NJ, Westlake L. Census of single homeless people in Sheffield. *BMJ* 1991;302:1387–89.
- 5 Timms PW, Fry AH. Homelessness and mental illness. *Health Trends* 1989;21:70–71.
- 6 Marshall M. Collected and neglected: are Oxford hostels for the homeless filling up with disabled psychiatric patients? *BMJ* 1989;299:706–9.
- 7 Reed A, Ramsden S, Marshall J, *et al*. Psychiatric morbidity and substance abuse among residents of a cold weather shelter. *BMJ* 1992;304:1028–29.
- 8 El Kabir DJ. Great Chapel Street Medical Centre. *BMJ* 1982;284:480–81.
- 9 Toon PD, Thomas K, Doherty M. Audit of work at a medical centre for the homeless over one year. *J R Coll Gen Pract* 1987;37:120–22.
- 10 *The Big Issue* 1996;170:25.