SHORT REPORT

Changing sex ratio in Iran, 1976–2000

M Ansari-Lari, M Saadat

J Epidemiol Community Health 2002;**56**:622–623

Sex ratio—that is, the ratio of male to total live births—is a subject of scientific interest but little is known about the factors that affect the sex ratio of humans. Many demographic and environmental factors have been suggested to affect the sex ratio. Some of these include birth order, race, coital rate, certain hormonal treatment,^{1,2} parental age,^{1,3} and exposure to environmental toxins.² Studies on secular trends in the human sex ratio at birth have shown that there are rapid increase during and just after war.⁴ The sex ratio can vary between populations and changes within populations over time, for example it has been reported that it decreased in recent decades in some countries such as Denmark⁵ and the United States⁶ and increased in Italy³ and Japan.⁷ There is no investigation about the sex ratio in Iran. To determine if the sex ratio has changed in Iran we examined the live births data from 1976 to 2000.

METHODS

Data about live births are routinely compiled by the statistic centre of Iran (SCI). The annual number of live births by sex for the period 1355 to 1378 Hejrae Shamsi (HS) (Iranian calendar) was obtained from the annual publications of SCI. It should be mentioned that this period was from 20 March 1976 to 19 March 2000. The war between Iran and Iraq began at mid-1359 HS (September 1980) and ended in early 1367 HS (July 1988). To determine the possible effect of the war on the sex ratio, the total studied period was divided to three distinct parts as before, during, and after war. The odds ratio (OR) (the odds of being male) was calculated, considering the exposure to the war as a risk factor. An OR > 1.0 indicates an increase in the sex ratio and an OR < 1.0 indicates a decrease in the sex ratio.

RESULTS AND DISCUSSION

The sex ratio for all live births in Iran was 0.517 in 1355 HS (1976–77) and declined gradually to 0.514 in 1358 HS (1979–80). It had decreased to 0.500 in 1359 HS (1980–81), which was the beginning of the war, and with some degree of fluctuation reached 0.503 in 1367 HS (1988–89), which was the end of the war. It increased thereafter and reached to 0.510 in 1378 HS (1999–2000) (fig 1). The sex ratio decreased significantly in Iran during war when compared with prewar (OR = 0.9287; 95% CI 0.9271 to 0.9303) and when the comparison was done between wartime and postwar years it showed significant increase (OR = 1.0440; 95% CI: 1.0431 to 1.0460). Another point is that the sex ratio decreased in postwar period when compared with prewar period (OR = 0.9701; 95% CI: 0.9683 to 0.9718).

This is not in complete agreement with previous studies on the first world war and second world war. It has been shown that the sex ratio increased in practically all the belligerent European countries during the first world war and in the succeeding two or three years; also an increase in England and Wales during second world war was detected.⁴

There are other factors such as parental age and birth order, which is suggested to affect the sex ratio. Although we were limited by availability of precise data about birth order and parental age of live births, but as suggested by Erickson the association between birth order and sex ratio accounts for only a very minor (<10%) proportion of the total variation in the

Figure 1 Sex ratio for live births in Iran, 1976–2000. HS means the Iranian calendar.

ratio.¹ Certainly this is true for parental age, the effect of which on sex ratio was not established by some studies.¹ According to population census data, the average Iranian family size was 4.9 in 1355 HS (1976–77) without change in 1365 HS (1986–87) and decreased significantly ($t = -35.8$, $p < 0.0001$) to 4.6 in 1375 HS (1996–97). Taken together, it is clear, that demographic changes did not provide an obvious explanation for the remarkable decline during wartime.

Of interest are the animal studies that have suggested that under favourable environmental and maternal conditions male infants would be favoured and the probability of delivering a female infant would increase under unfavourable conditions,⁸ which surprisingly seems reasonable to explain the findings of our study. There is little doubt that the environmental conditions were disturbed during wartime and became unfavourable because of psychological tensions and stressful life. However, it is possible that unidentified factors would be responsible for this variation and it therefore needs further studies to explore those factors.

Authors' affiliations

M Ansari-Lari, Department of Community Medicine, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran
M Saadat, Department of Biology, College of Sciences, Shiraz University, Shiraz, Iran

Funding: the study was supported by Shiraz University.

Conflicts of interests: none.

Correspondence to: Dr M Saadat, Department of Biology, College of Sciences, Shiraz University, Shiraz 71454, Iran

Accepted for publication 11 March 2002

REFERENCES

- 1 Erickson JD. The secondary sex ratio in the United States 1969–1971: association with race, parental age, birth order, parental education and legitimacy. *Ann Hum Genet* 1979;**40**:205–12.
- 2 James WH. The human sex ratio. Part I: a review of the literature. *Hum Biol* 1987;**59**:721–5.

- 3 **Ulizzi L**, Zonta LA. Factors affecting the sex ratio in humans: multivariate analysis of the Italian population. *Hum Biol* 1995; **67**:59–67.
- 4 **McMahon B**, Pugh TF. Sex ratio of white birth in the United States during second world war. *Am J Hum Genet* 1954; **7**:284–92.
- 5 **Moller H**. Change in male:female ratio among newborn infants in Denmark. *Lancet* 1996; **348**:828–9.
- 6 **Marcus M**, Kiely J, Xu F, *et al*. Changing sex ratio in the United States, 1969–1995. *Fertil Steril* 1998; **70**:270–3.
- 7 **Imaizumi Y**, Murata M. The Changing sex ratio in Japan. *Jpn J Hum Genet* 1981; **26**:71–81.
- 8 **Trivers RL**, Willard DE. Natural selection of parental ability to vary the sex ratio of offspring. *Science* 1973; **179**:90–2.

Call for peer reviewers

Clinical Evidence is a regularly updated evidence based journal available world wide both as a paper version and on the internet. *Clinical Evidence* urgently needs to recruit a number of new contributors. Contributors are health care professionals or epidemiologists with experience in evidence based medicine and the ability to write in a concise and structured way.

Clinical Evidence needs to recruit a number of new peer reviewers. Peer reviewers are health care professionals or epidemiologists with experience in evidence based medicine. As a peer reviewer you would be asked for your views on the clinical relevance, validity and accessibility of specific topics within the journal, and their usefulness to the intended audience (international generalists and health care professionals, possibly with limited statistical knowledge). Topics are usually 2000–3000 words in length and we would ask you to review between 2–5 topics per year. The peer review process takes place throughout the year, and our turnaround time for each review is ideally 10–14 days.

If you are interested in becoming a peer reviewer for *Clinical Evidence*, please complete the peer review questionnaire at www.clinicalevidence.com or contact Polly Brown (pbrown@bmjgroup.com).