REVIEW

Excess in cardiovascular events on Mondays: a meta-analysis and prospective study

Adrian G Barnett, Annette J Dobson

J Epidemiol Community Health 2005;**59**:109–114. doi: 10.1136/jech.2003.019489

The aim of this paper was to summarise the reported excess in coronary events on Mondays, and examine the evidence for three competing explanations: stress, alcohol consumption, or registration errors. A review of the literature found 28 studies covering 16 countries and over 1.6 million coronary events. The overall Monday excess was small; in a population experiencing 100 coronary events per week there was one more event on Monday than other days. The excess was larger in men and in studies including sudden cardiac death or cardiac arrests. In a prospective study an increase in events on Mondays was associated with greater alcohol consumption, lower rainfall, and the month of January. The excess in coronary events on Mondays is a persistent phenomenon. The size of the effect varies widely between populations. There is some evidence of an association with alcohol consumption, but a definitive explanation remains elusive and is likely to remain so because of the smallness of the effect and the paucity of high quality data.

Differences among studies may be useful for identifying the real cause (or causes) of the Monday excess.⁹ If the hypotheses of alcohol consumption or stress were true, then we would expect the differences between studies to be attributable to study populations. If the excess was caused by data collection and recording procedures then the differences between studies should be related to study design and methods.

To date no individual study has simultaneously investigated the three hypotheses. We attempt to do this using a prospective community based study of coronary events, conducted as part of the WHO MONICA Project (to monitor trends and determinants in cardiovascular disease).¹⁰

METHODS

Meta-analysis

We followed the meta-analysis guidelines for observational studies given by the MOOSE group.¹¹

We searched the databases of Medline, Pubmed, the Cochrane Library, and Web of Science using combinations of the words "Monday", "Hebdomadal", "Cardiovascular", and "Infarction". The search had no time period restriction. For every relevant publication we searched all related references. We contacted three authors of relevant publications about other work in the area, but this did not yield any extra studies. Two studies (one German, one Italian) were translated into English. One paper was an abstract.¹²

Prospective study

In this study we used data from the MONICA centre in Newcastle, Australia.¹³ The study population was adults age 25–69 years living in the Hunter Region of New South Wales; this is a temperate area on the east coast, with a mixed economy of manufacturing, mining, and agriculture. Ethical approval for the study was obtained from the University of Newcastle and Hunter Region Health Service Ethics Committees.

All suspected CHD events in the study population were identified prospectively from surveillance of hospital admissions, death records, and other sources. Those events confirmed as eligible were added to the coronary register.

The date of coronary event onset was recorded as the day symptoms began. Instructions were

Rates of coronary heart disease (CHD) vary over the day, the week, and the year.¹ A decrease in risk at night and increase in the morning is thought to be attributable to tachycardia and an increase in blood pressure after waking. The size of the effect has been estimated as a 40% increase in events in the morning (6 am to noon) compared with the rest of the day.² An increase in coronary events in colder months and decrease in warmer months has been linked to changes in the weather.³ For example, in Ireland a 45% increase in cardiovascular deaths was observed in winter compared with summer.⁴ The weekly variation of an increase on Monday has yet to be convincingly explained, nor has the size of the effect been adequately described. Some hypothesised reasons for this variation include:

- (1) an increase in stress on Mondays in employed people⁵
- (2) weekend alcohol consumption (because high use may cause arrhythmias and thrombosis^{6 7})
- (3) anomalies in data registration.⁸

All, some, or none of these hypotheses may be correct. To investigate them further—and to clarify the size of the excess—we first summarise the existing findings of the Monday excess.

See end of article for authors' affiliations

Correspondence to: Dr A G Barnett, School of Population Health, University of Queensland, Herston, QLD 4006, Australia; a.barnett@sph.uq.edu.au

Accepted for publication 23 April 2004

Abbreviations: CHD, coronary heart disease; SCD, sudden cardiac death; CVD, cardiovascular disease

Table 1 Summary of studies presenting results on the weekly distribution of fatal coronary events, ordered by total number of events

Reference	Population(s)	Period	Diagnostic category	Number of events (N)	Days with a significant excess	Monday excess $\hat{M}\%$	95% CI
Rogot	USA	1962–66	CHD	>2 million	Monday*	NA	
AlHW	Australia	1979–99	Acute MI	429255	Monday	2.0	1.3 to 2.8
Macfarlane	England and Wales	1969–72	Acute MI	373598	Monday	3.2	2.4 to 4.0
Evans	Scotland	1986–95	CHD	170244	Monday, Saturday†, Sunday†	3.1	1.9 to 4.3
Chenet, 1998	Moscow	1993–95	CVD	117286	Monday‡, Saturday‡, Sunday‡	0.8	–0.6 to 2.2
Näyhä	Finland	1961–72	IHD	113246	Monday, Saturday	3.4	2.0 to 4.8
Massing	Saxony, Germany	1968–77	MI	84538	Monday§, Saturday§	0.8	–0.9 to 2.4
Barnett	39 centres worldwide	1980–95	MI	74068	Monday	6.8	5.0 to 8.6
Chenet, 2001	Lithuania	1988–97	CVD and IHD	41452	Monday, Saturday, Sunday	9.7	7.3 to 12.2
Arntz	Berlin	1987–91	SCD	24061	Monday	8.3	5.1 to 11.5
Macfarlane	Greater London	1965–66	IHD	22813	Monday	4.0	0.8 to 7.2
Beard	Rochester, USA	1950–75	SCD	1054	None	–3.0	–17.6 to 11.6
Pasqualetti	Aquila, Italy	1970–87	SCD	269	None	24.9¶	–7.1 to 56.9
Myers	Newcastle, UK	1971–72	SCD	100	None	26.0	–26.7 to 78.7
Nicolau	Bucharest	1981–84	CVD	86	Monday	70.9	7.4 to 134.5

CHD, coronary heart disease; CVD, cardiovascular disease; IHD, ischaemic heart disease; MI, myocardial infarction; SCD, sudden cardiac death; NA, not available. *No statistical testing done but numbers involved make it highly likely that observed increase is statistically significant. †Subgroup of CHD deaths outside hospital in men under 65 years old. ‡Subgroup of men ($N=69934$). §Subgroup of men age 25–54 years ($N=6259$). ¶Number of events estimated from graphical results.

also given to code the day as missing if the date was uncertain, or there was a vague onset and a slow increase in severity of symptoms over several days, that is a crescendo onset. If the subject was found dead circumstantial evidence was used to estimate the day of death. Subjects were interviewed soon after the event; if the subject had died then information was obtained from relatives or other informants.

We used employment as a surrogate measure of stress. We assumed that people in part time or full time employment were more likely to be stressed on Mondays than those who were unemployed, retired, studying, engaged in home duties, or unable to work. The employment question was not used

during part of the registration period, so only a subset of the total sample was available for this analysis.

The number of questionnaire responses that were missing or not known was used as an indicator of data quality. We assumed that more guessing of onset dates would take place where fewer details were available.

Because the weather may affect patterns of alcohol consumption, daily temperature and rainfall data were obtained from the Hunter Valley Research Foundation for 1985–90. We summarised both average daily temperature and total daily rainfall over the previous zero to three days to model the possible lag between exposure and onset of the coronary event.

Table 2 Summary of studies presenting results on the weekly distribution of non-fatal coronary events or both fatal and non-fatal events, ordered by total number of events

Reference	Population(s)	Period	Diagnostic category	Number of events (N)	Days with a significant excess	Monday excess $\hat{M}\%$	95% CI
Barnett	39 centres worldwide	1980–95	Non-fatal MI	153305	Monday	9.2	7.9 to 10.5
Peters	32 N American centres and one Swedish	1987–91	Acute MI	22516	Monday	8.5	5.2 to 11.8
Gnecchi	Italy	1988–89	Acute MI	11472	Monday	8.6	4.0 to 13.2
WHO	18 European centres and one Australian	1971	Acute MI	8502	Saturday	4.1	–1.2 to 9.4
Allegra	New Jersey, USA	1988–98	Cardiac arrests	6827	Saturday	5.1	–0.8 to 11.0
Willich	Augsburg, Germany	1985–90	Acute MI and SCD	5596	Monday	11.8	5.1 to 18.5
Peckova	Seattle, USA	1985–93	Cardiac arrests	5248	Monday	8.0	1.2 to 14.9
Genès	France	1995	MI	2348	None	–0.1	–10.0 to 9.8
Thompson	Leicester, UK	1979–88	Acute MI	2254	Monday, Tuesday	12.7	2.1 to 23.4
Spielberg	Dessau, Germany	1980–88	Acute MI	1901	Monday	NA	
Bilora	Northern Italy	1992	Acute MI	817	Monday*	4.5	–12.6 to 21.6
Ku	Taipei	1992–96	Acute MI	540	Tuesday	5.0	–16.1 to 26.1
Zhou	China	1991	Acute MI	428	Saturday	–15.0	–36.6 to 6.7
Rabkin	Canada	1948–77	SCD	358	Monday	34.9	6.3 to 63.5
Paul	Boston, USA	NA	MI	334	Monday and Tuesday combined	29.9†	0.8 to 59.1

MI, myocardial infarction; SCD, sudden cardiac death; NA, not available. *Subgroup of active workers ($N=264$). †No results given for Monday alone, so an estimate of $\hat{M}\%$ was made by halving the combined results from Monday and Tuesday.

Statistical methods

The excess in events on a Monday was defined as

$$\hat{M}\% = 100 \left(\frac{m}{N} - \frac{1}{7} \right) / \frac{1}{7} \quad (1)$$

where m is the observed number of events on a Monday and N is the total number of events over all days. The value $\hat{M}\%$ is the percentage of excess events on a Monday compared with the expected of proportion of 1/7. Confidence intervals for the excess percentage of Monday events were calculated using the normal approximation to the binomial distribution.

Meta-analysis

The Monday excess was estimated using a random effect meta-analysis,¹⁴ and was implemented using the Bayesian WinBUGS software.¹⁵ Results using Bayesian methods are reported with a 95% posterior interval (PI), and results using standard methods are reported with a 95% confidence interval (CI). Both intervals estimate a probable range of the parameter of interest, but the PI has the simpler interpretation as an interval that has a high (95%) probability of containing the true parameter value.

A test of heterogeneity between studies was performed using the I^2 statistic,¹⁶ which is interpretable as the proportion of total variation in the Monday excess that is attributable to differences between studies; the remaining variation is attributable to random error.

To examine possible explanations of any differences between studies we used a regression tree¹⁷ analysis of the Monday excess. The study level explanatory variables were: data collection method (retrospective compared with prospective), study location (Eastern Europe compared with other—because excessive alcohol consumption is higher in Eastern Europe⁷), study time (midpoint in years), study duration (years and months), event outcome (fatal only compared with only non-fatal or both fatal and non-fatal), total number of events, and diagnostic category (see tables 1 and 2). Study time was used to examine any temporal trend in the Monday excess. Study duration was used as a proxy for quality, with shorter studies possibly having greater accuracy. Diagnostic category was not a precise grouping, but it did broadly differentiate the studies. There was probably some misclassification, for example, studies with a diagnostic category of CHD probably included some sudden deaths, and studies with a diagnostic category of sudden cardiac death probably included some unattended sudden deaths from myocardial infarction and other CHD. We classified study outcomes into five diagnostic groups: myocardial infarction (MI) or acute MI, sudden cardiac death (SCD), cardiac arrest, CHD or ischaemic heart disease (IHD), and cardiovascular disease (CVD). Because diagnostic category

was a nominal explanatory variable with five categories we used a regression tree rather than standard regression analysis. The tree method evaluated every possible split of the five categories into two groups, while simultaneously evaluating the effect of the other covariates. We used 10-fold cross-validation to select an optimal tree. The tree analysis was run using the “rpart” library in R.¹⁸

Prospective study

The excess in events on a Monday was examined using equation (1). The distribution of events over the month was also examined to check for excesses on the 1st or 15th of the month that would indicate some systematic imputation of unknown onset dates.⁸

Univariate comparisons between either the missing and non-missing event dates, or Monday and non-Monday events were made using the χ^2 test for categorical variables and the Wilcoxon two sample test for continuous variables.

We used multiple logistic regression to predict an event on a Monday using the explanatory variables of: age, sex, usual frequency and quantity of alcohol consumption, smoking status (non-smoker, ex-smoker, current smoker, unknown), history of previous MI, apparent survival time, average temperature and total rainfall (previous 0–3 days), month, year, and number of missing data items. Employment status was not used in the logistic regression because it was not collected throughout the study period. A model was chosen using a backward stepwise procedure (with a significance level of 10%), and the regression diagnostics of Pregibon.¹⁹

RESULTS

Meta-analysis: description of studies

We found 28 community based studies^{5 6 8 12 20–43} with results on the weekly distribution of CHD events. Two studies^{8 23} presented results in two populations, so there is a total of 30 results summarised in table 1 (fatal coronary events) and table 2 (non-fatal coronary events or both fatal and non-fatal events). The number of days after the onset of symptoms that a death would still be attributable was not always clearly defined, which may cause some inconsistency between studies.

Two studies^{44 45} were found that examined the Monday excess in a population at high risk of a coronary event, and hence were not included.

With three exceptions,^{25 35 38} the studies were retrospective and relied on routinely collected data such as death certificates or hospital records, although five studies^{5 8 12 22 39} used a coronary event register that was updated close to the date of the event. Three studies^{12 31 35} recruited subjects from admissions to an emergency department. The Moscow study³⁶ reported results for cardiovascular deaths based on the Soviet classification system that may include stroke.

Table 3 Meta-analysis of excess Monday events and subgroup meta-analyses by gender (where available), diagnostic category, event category, and study size

Category	Subgroup	Number of studies	Number of events (N)	Amount of heterogeneity* (I^2)	Estimated excess $\hat{M}\%$	95% PI
Gender	Male	14	656730	83.0	9.6	2.3 to 17.7
	Female	12	331064	76.2	4.0	–5.4 to 13.7
Diagnostic category	SCD and Cardiac arrest	8	43513	24.6	11.6	–3.3 to 27.4
	CHD/IHD, CVD, MI, acute MI	19	1628768	90.0	4.8	–0.1 to 9.6
Event category	Fatal events only	14	1452070	85.2	5.3	–1.6 to 12.9
	Fatal and non-fatal events	13	220211	31.6	7.1	–0.6 to 15.1
Study size	Larger studies only (N>400)	23	1671468	88.6	4.9	1.1 to 8.7
All studies		27	1672281	87.3	5.9	2.0 to 9.9

*The percentage of total variation in the Monday excess that is attributable to heterogeneity between studies. CHD, coronary heart disease; CVD, cardiovascular disease; IHD, ischaemic heart disease; MI, myocardial infarction; SCD, sudden cardiac death; PI, posterior interval; N, total number of events.

Figure 1 Meta-analysis of community based studies: overall and study specific estimates of the Monday effect. Studies are listed in order of the total number of events, for the exact *N* see tables 1 and 2. The line at *M*% = 0 represents no excess. *Upper limit of 134.5% for smallest study²⁹ not shown to improve the scale. Lines represent the mean estimate for studies reporting fatal events only, crosses represent the mean estimate for studies that included some non-fatal events.

A statistically significant excess in coronary events on Mondays was reported in 22 of the 30 results (and 20 of the 28 studies). Significant excesses in events were also found in some studies on Tuesday, Saturday, and Sunday. Only four studies reported no significant differences on any days. The excess in events on Mondays has been observed: in both fatal and non-fatal events; in both sexes; in the United States, Canada, Australia, and Europe; and in the 1960s, 70s, 80s, and 90s.

Table 4 Crude estimates of the Monday excess of events in key categorical variables in the prospective study

Variable	Category	Number of events (N)	Monday excess* <i>M</i> %	p Value
Gender	Male	7669	3.2	0.98
	Female	3720	3.4	
Apparent survival time	Death	1081	3.6	0.63
	<1 hour			
	Death	1014	10.5	
	≥1 hour			
Employed‡	Alive at 28 days	8532	2.0	0.84
	Unknown	312	14.4	
	Yes	1602	5.3	
	No	4428	5.8	
Usual frequency of alcohol consumed (days/week)	Unknown	289	-3.1	0.040
	High (>4)	2134	15.1	
	Low (0-4)	7875	-0.2	
	Unknown	930	5.4	
Usual quantity of alcohol consumed (number standard drinks)	High (>8)	753	7.8	0.51
	Low (0-8)	8651	1.9	
	Unknown	1535	9.0	
	Unknown			
Binge drinking	Yes	176	-24.4	0.22
	No	9266	2.9	
	Unknown	1497	9.0	
Month	January	880	22.5	0.0017
	Other	10059	1.6	
Overall		10939	3.3	

**M*% defined in equation (1), larger *M*% indicates a greater excess in events on a Monday. † χ^2 Test; ‡*N* = 6319 as employment status was only available for a subset of questionnaires.

The Monday excess was shown to be greater in younger compared with older subjects in five studies (age <65 years,⁴² age ≤65,⁴⁰ age <50,⁴¹ men <65,²⁴ men <60).²³ An excess in both younger and older age groups was shown in three studies,^{5, 32, 36} while one study³³ reported that the excess was greater in the elderly population (age >65 years). It has been suggested that the excess at younger ages is attributable to employment status and therefore stress.^{23, 40} A significantly greater Monday excess in working compared with non-working subjects was shown in two studies,^{5, 31} but no statistically significant difference was found in another.³³

A greater excess in Monday events in men than women was shown in two studies,^{23, 24} whereas one⁵ had a greater excess in women, and another³³ found no statistically significant difference.

The Monday effect was shown to be larger in cardiovascular deaths outside hospital in Moscow³⁶ and in a subgroup of coronary deaths occurring outside hospital in people with no previous admissions for CHD in Scotland.⁴¹ The authors of both papers suggested that the excess in deaths on Monday was attributable to binge drinking.

Using data from many centres in the WHO MONICA Project we showed that the Monday excess was greater in

Table 5 Distribution of continuous variables among those experiencing, and those not experiencing an event on Monday in the prospective study

Variable	Monday	Median	p Value*
Age	Yes	60	0.30
	No	61	
Total rainfall† (mm) (previous 0-3 days)	Yes	0.75	0.30
	No	0.90	
Average temperature† (°C) (previous 0-3 days)	Yes	23.4	0.23
	No	23.2	
Number of missing data items	Yes	1	0.54
	No	2	

*Wilcoxon two sample test; †*N* = 5796 as weather data were only available from 1985-90.

Key points

- There is a small excess in cardiovascular events on Mondays in many populations worldwide. The excess seems highest in sudden cardiac deaths.
- Some evidence was found of an association with alcohol consumption, but a definitive explanation of the Monday effect remains elusive.

centres with less thorough data collection procedures, and fewer missing dates of coronary onset.⁸ Excesses of events of a similar size to the Monday excess were found on the first of the month, suggesting that when the date of onset was unknown, some centres imputed Monday or the first of the month rather than recording the date as missing.

Meta-analyses: random effects and meta-regression models

Three studies^{12 21 34} were excluded from the meta-analysis as they did not report the total number of events or the number observed on a Monday, which are required for equation (1). Meta-analyses of the remaining studies based on over 1.6 million coronary events from 27 results are shown in table 3.

The estimated mean Monday excess was 5.9% (95% PI: 2.0% to 9.9%) or an excess number of events on a Monday of 0.8 in a population experiencing 100 events per week (95% PI: 0.3 to 1.4). Figure 1 shows a summary plot of the results. The studies are listed in order of total number of events, with a different marker if non-fatal events were included. The very large Monday excesses in the four smallest studies suggested a publication bias. To reduce the possible impact of publication bias we ran a meta-analysis using the 22 results each with more than 400 events and obtained an estimated Monday excess of 4.8% (95% PI: 0.7% to 8.9%).

The Monday excess was smaller for women than men. In women the excess was 4.0% (95% PI: -5.4% to 13.7%), indicating a non-significant excess.

In the regression tree analysis only diagnostic category was selected by the optimal tree; the Monday effect was more apparent in studies of sudden cardiac death or cardiac arrest. In a subgroup meta-analysis of the eight studies with the outcome of SCD or cardiac arrest the Monday excess was 11.6% (95% PI: -3.3 to 27.4).

The heterogeneity between the results was large, in the overall meta-analysis 87.3% of the total variation in the Monday excess was due to differences between studies. There was a much less heterogeneity between the results of studies that included non-fatal events and in studies of SCD or cardiac arrest.

Prospective study

A total of 11 637 events was recorded between 1984 and 1994. We excluded 698 events (6.0%) because the exact date of onset was incomplete. Compared with the events with a confirmed date these events were significantly more likely to be fatal, in older people, and have incomplete information on employment and alcohol consumption.

Of the 10 939 coronary events with an exact onset date, 1614 occurred on Monday, giving an excess of 3.3% (95% CI: -1.4% to 7.9%), which is in the 95% PI of the meta-analysis. There was no significant excess in events on the 1st, 15th, or 30–31st of the month.

Crude associations between events on a Monday and various putative predictors are shown in tables 4 and 5; only alcohol consumption and the month of January showed a statistically significant association. January was specifically

Policy implications

The small excess in events on Mondays does not warrant any policy attention beyond the well established measures to reduce heart disease in general.

chosen as it showed a strong association in exploratory analyses. In binge drinkers fewer events were observed on Monday ($\hat{M}\% = -24.4\%$), although this was not statistically significant. In contrast more events occurred on Friday and Sunday in binge drinkers (Excess > 19%), but again these differences were not significant because of the small total number of binge drinkers (results not shown).

In a multiple logistic regression model, events on a Monday were more likely to occur in January (OR = 1.36, 95% CI: 1.06 to 1.75) and in periods with less rainfall (OR = 1.027, 95% CI: 1.012 to 1.043, for a one millimetre decrease in total rainfall).

DISCUSSION

The meta-analysis showed that the Monday excess in coronary events is small but greater than zero. The excess on Mondays is less than 1 event in 100, or a relative increase of 1%, whereas events in the morning have been shown to increase by 40%,² and in events in winter by 45%.⁴

To investigate the cause of the excess we looked for differences in the published results that might be explained by study features. Differences in the size of the excess were found according to gender (greater in men) and diagnostic category (greater in studies of sudden cardiac death or cardiac arrests). The plot of study results (fig 1) suggested a publication bias, and when only large studies were considered the Monday excess reduced but remained statistically significant.

The link between binge drinking and heart disease is strongest in sudden cardiac deaths,⁴⁶ and therefore the greater Monday excess in studies of sudden cardiac death provides support for the binge drinking hypothesis, and differences in the Monday excess among diagnostic categories explain some of the large variability in results. However, sudden cardiac deaths are also more likely to be unwitnessed, and hence the greater Monday excess found in studies of these events also supports the registration hypothesis.

We did find other evidence to support the hypothesis of binge drinking. Men generally consume more alcohol than women, which could explain their greater Monday excess in the meta-analysis. The prospective study found a direct association between high alcohol consumption and the Monday excess, but binge drinking was negatively associated. We also found associations between the Monday excess and January and low rainfall. Both these factors can be indirectly linked to alcohol consumption as in Australia January is a hot month traditionally devoted to summer holidays with a high incidence of social drinking. Additionally alcohol consumption and high temperatures may interact to increase the risk of a coronary event.⁴⁷ These associations with the Monday excess, and the large variation in the Monday excess in the meta-analysis, suggest that there may be populations in which the Monday effect may be important.

Apart from the link with sudden cardiac deaths we found little evidence to support the registration hypothesis. If this hypothesis were true we would have expected a greater excess in fatal events on Mondays, because the date of onset might be unknown for unwitnessed deaths (and these events could have had the date of onset imputed as Monday). In fact

there was a slightly greater Monday excess in non-fatal events. Similarly, we would have expected retrospective studies to have less accurate onset dates and hence have a higher proportion of Monday events; this was not the case. Furthermore, the prospective analysis showed no difference in the Monday excess according to the number of missing items on the questionnaire (used as a proxy for data quality), although there was a non-significant excess in deaths with an unknown survival time.

Evidence for the employment/stress hypothesis is provided by the greater Monday excess found in men in the meta-analysis (assuming that men are more likely to be employed and exposed to greater stress on a Monday). Similarly the Monday excess in non-fatal events might be attributable to non-fatal events being more common among younger people,¹⁰ who are also more likely to be employed. If data were available for Moslem countries (where the working week begins on Saturday), or in Israel (where the working week begins on Sunday), they could be used to investigate the employment/stress hypothesis further.

The true cause of the Monday excess remains elusive, because of the absence of studies designed to investigate this phenomenon. We estimate that to find the cause would need a study of over 40 000 events. This is based on a proportion of (1/7+1/100) events on Monday and 1/7 on other days of the week, and an 80% power and 5% significance level. The study would need to record accurately each subject's alcohol consumption, employment (or stress) and other factors. Such a large study to examine an apparently small effect seems unjustified.

ACKNOWLEDGEMENTS

Thanks to Elsa Quagliotto for translating a journal article from Italian to English, and Lois Englberger for translating a journal article from German to English. Thanks also to Martin McKee and Susana Sans for helpful comments on the manuscript. We thank the Hunter Valley Research Foundation for providing their weather data free of charge.

Authors' affiliations

A G Barnett, A J Dobson, School of Population Health, University of Queensland, Herston, Australia

Funding: this work was funded by the National Health and Medical Research Council of Australia (grant numbers 100954 and 252834).

Conflicts of interest: none declared.

REFERENCES

- Arntz HR, Müller-Nordhorn J, Willich SN. Cold Monday mornings prove dangerous: epidemiology of sudden cardiac death. *Curr Opin Crit Care* 2001;**7**:139–44.
- Cohen MC, Rohlf KM, Lavery CE, et al. Meta-analysis of the morning excess of acute myocardial infarction and sudden cardiac death. *Am Heart J* 1997;**79**:1512–15.
- The Eurowinter Group. Cold exposure and winter mortality from ischaemic heart disease, cerebrovascular disease, respiratory disease, and all causes in warm and cold regions of Europe. *Lancet* 1997;**349**:1341–6.
- Eng H, Mercer JB. Seasonal variations in mortality caused by cardiovascular diseases in Norway and Ireland. *Am J Emerg Med* 1998;**5**:89–95.
- Willich SN, Lowel H, Lewis M, et al. Weekly variation of acute myocardial infarction: Increased Monday risk in the working population. *Circulation* 1994;**90**:87–93.
- Chenet L, Britton A, Kalediene R, et al. Daily variations in deaths in Lithuania: the possible contribution of binge drinking. *Int J Epidemiol* 2001;**30**:743–8.
- McKee M, Britton A. The positive relationship between alcohol and heart disease in eastern Europe: potential physiological mechanisms. *J R Soc Med* 1998;**91**:402–7.
- Barnett AG, Dobson A, for the WHO MONICA Project. Is the increase in coronary heart disease on Mondays an artifact of registration? *Epidemiology* 2004;**15**:583–8.
- Glasziou PP, Sanders SL. Investigating causes of heterogeneity in systematic reviews. *Stat Med* 2002;**21**:1503–11.
- Tunstall-Pedoe H, ed. *MONICA monograph and multimedia sourcebook*. Geneva: World Health Organisation, 2003.
- Stroup DF, Berlin JA, Morton SC, et al. Meta-analysis of observational studies in epidemiology. *J Am Stat Assoc* 2000;**283**:2008–12.
- Paul SD, Eagle KA, O'Gara PT. Weekly rhythm of acute myocardial infarction: are physicians more aggressive with thrombolysis after the weekend? [Abstract]. *Circulation* 1993;**88**(suppl 1):509.
- McElduff P, Dobson AJ. Case fatality after an acute cardiac event: the effect of smoking and alcohol consumption. *J Clinical Epidemiol* 2001;**54**:58–67.
- Normand ST. Meta-analysis: formulating, evaluating, combining, and reporting. *Stat Med* 1999;**18**:321–59.
- Spiegelhalter DJ, Thomas A, Best NG. *Win BUGS version 1.2 user manual*. London: Technical report, MRC Biostatistics Unit, 1999.
- Higgins JPT, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med* 2002;**21**:1539–58.
- Breiman L, Friedman JH, Olshen RA, et al. *Classification and regression trees*. Belmont, CA: Wadsworth International Group, 1984.
- Ihaka R, Gentleman RR. A Language for data analysis and graphics. *Journal of Computational and Graphical Statistics* 1996;**5**:299–314.
- Pregibon D. Logistic regression diagnostics. *Ann Stat* 1981;**9**:705–24.
- Myers A, Dewar HA. Circumstances attending 100 sudden deaths from coronary artery disease with coroner's necropsies. *Br Heart J* 1975;**37**:1133–43.
- Rogot E, Fabsitz R, Feinleib M. Daily variation in USA mortality. *Am J Epidemiol* 1976;**103**:198–211.
- World Health Organisation. *Myocardial infarction community registers. Public health in Europe no 5*. Technical report. Geneva: World Health Organisation, 1976.
- Macfarlane A, White G. Deaths: the weekly cycle. *Population Trends* 1997;**7**:7–8.
- Näyhä S. Short and medium-term variations in mortality in Finland. *Scand J Soc Med* 1980;**21**(suppl.):21–101.
- Rabkin SW, Mathewson FAL, Tate RB. Chronobiology of cardiac sudden death in men. *JAMA* 1980;**244**:1357–8.
- Beard MC, Fuster V, Elveback LR. Daily and seasonal variation in sudden cardiac death, Rochester, Minnesota, 1950–1975. *Mayo Clin Proc* 1982;**57**:704–6.
- Massing W, Angermeyer MC. Myocardial infarction on various days of the week. *Psychol Med* 1985;**15**:851–7.
- Pasqualetti P, Colantonio D, Casale R, et al. Cronobiologia della morte cardiaca improvvisa. Evidenza di una periodicità circadiana, circasettana e circannuale nella sua incidenza. (In Italian). *Minerva Med* 1990;**81**:391–8.
- Nicolau GY, Haus E, Popescu M, et al. Circadian, weekly, and seasonal variations in cardiac mortality, blood pressure and catecholamine excretion. *Chronobiol Int* 1991;**8**:149–59.
- Thompson DR, Pohl JEF, Sutton TW. Acute myocardial infarction and day of the week. *Am J Cardiol* 1992;**69**:266–7.
- Bilora F, Vigna GB, Manfredini R, et al. Acute myocardial infarction incidence during the week: a chronobiological evaluation. *Minerva Cardioangiol* 1993;**43**:559–62.
- Gnecchi-Ruscone T, Piccaluga E, Guzzetti S, et al. Morning and Monday: critical periods for the onset of acute myocardial infarction. the GISSI 2 study experience. *Eur Heart J* 1994;**15**:882–7.
- Peters RW, Brooks MM, Zoble RG, et al. Chronobiology of acute myocardial infarction: cardiac arrhythmia suppression trial (CAST) experience. *Am J Cardiol* 1996;**78**:1198–201.
- Spielberg C, Falkenhahn D, Willich SN, et al. Circadian, day-of-week, and seasonal variability in myocardial infarction: comparison between working and retired patients. *Am Heart J* 1996;**132**:579–85.
- Genès N, Vaur L, Renault M, et al. Rythme circadien des infarctus du myocarde en France: résultats de l'étude USIK. (In French). *Presse Med* 1997;**26**:603–8.
- Chenet L, McKee M, Leon D, et al. Alcohol and cardiovascular mortality in Moscow: new evidence of a causal association. *J Epidemiol Community Health* 1998;**52**:772–4.
- Ku CS, Yang CY, Lee WJ, et al. Absence of a seasonal variation in myocardial infarction onset in a region without temperature extremes. *Cardiology* 1998;**89**:277–82.
- Zhou RH, Xi B, Gao HQ, et al. Circadian and septadian variation in the occurrence of acute myocardial infarction in a Chinese population. *Jpn Circ J* 1998;**62**:190–2.
- Peckova M, Fahrenbruch C, Cobb L, et al. Weekly and seasonal variation in the incidence of cardiac arrests. *Am Heart J* 1999;**137**:512–15.
- Arntz HR, Willich SN, Schreiber C, et al. Diurnal, weekly and seasonal variation of sudden death. *Eur Heart J* 2000;**21**:315–20.
- Evans C, Chalmers J, Capewell S, et al. "I don't like Mondays" – day of the week of coronary heart disease deaths in Scotland: study of routinely collected data. *BMJ* 2000;**320**:218–19.
- Allegra JR, Cochrane DG, Allegra EM, et al. Calendar patterns in the occurrence of cardiac arrest. *Am J Emerg Med* 2002;**20**:513–17.
- Australian Institute of Health and Welfare (AIHW). *Seasonality of death. Technical report bulletin no 3*. Canberra: AIHW, 2002.
- Bleyer AJ, Russell GB, Satko SG. Sudden and cardiac death rates in hemodialysis patients. *Kidney Int* 1999;**55**:1553–9.
- Peters RW, McQuillan S, Resnick SK, et al. Increased Monday incidence of life-threatening ventricular arrhythmias: experience with a third-generation implantable defibrillator. *Circulation* 1996;**94**:1346–9.
- McKee M, Shkolnikov V, Leon DA. Alcohol is implicated in the fluctuations in cardiovascular disease in Russia since the 1980s. *Ann Epidemiol* 2001;**11**:1–6.
- Keatinge WR. Death in heat waves. *BMJ* 2003;**327**:512–13.