080

SLEEP DURATION AND BREAST CANCER RISK: A META-ANALYSIS

doi:10.1136/jech.2010.120956.80

X Wang, B J Cairns, R C Travis. *Cancer Epidemiology Unit, University of Oxford, Oxford, UK*

Background It is hypothesised that sleep duration affects the production of melatonin and subsequently influences cancer risk. The relationship between sleep duration and risk of breast cancer has been investigated recently but findings from epidemiological studies have been inconsistent and have not been summarised quantitatively. We aimed to conduct a meta-analysis of observational studies of the association between sleep duration and breast cancer.

Methods Relevant publications were identified from reviews and computer-aided searches using PubMed, with keywords "sleep duration", "breast cancer", "survival rate", "mortality", "morbidity", "incidence" and "risk", up to July 23rd, 2009. RR estimates and 95% CIs were extracted for the comparison between the highest exposure group, women who slept ≥ 9 hours, and the reference group that comprised women who had a moderate sleep duration (7 or 8 hours). Summary RRs were estimated by calculating the average of the log RRs, weighted by the inverse variances of the log RRs.

Results Five studies, four with prospective data and one case-control study, were identified on the risk of breast cancer in relation to sleep duration. The published data include 9166 women with incident invasive breast cancer and 147 344 women without breast cancer. When results from these studies were combined, the aggregate RR was 0.96 (95% CI 0.86 to 1.07) for women with the longest sleep duration compared to those in the reference group with a shorter sleep duration. When analysis was restricted to prospective data, the aggregate RR was 0.89 (95% CI 0.78 to 1.01). There was no evidence for significant heterogeneity in this association by menopausal status.

Conclusion Meta-analysis of the published epidemiological data provides no strong evidence for a relationship between sleep duration and risk of breast cancer.

Plenary

081

PREDICTORS OF SURVIVAL IN CHILDREN BORN WITH DOWN SYNDROME

doi:10.1136/jech.2010.120956.81

¹P W G Tennant, ¹M S Pearce, ²M Bythell, ^{1,2}J Rankin. ¹*Institute of Health and Society, Newcastle University, Newcastle-upon-Tyne, UK;* ²*Regional Maternity Survey Office, Newcastle-upon-Tyne, UK*

Objective To investigate the influences of survival in children born with Down Syndrome.

Design Population-based case-series derived from the Northern Congenital Abnormality Survey.

Setting A geographically distinct area of Northeast England, incorporating North Cumbria, Northumberland, Tyne and Wear, Country Durham, and the Tees Valley.

Participants 1101 individuals with Down Syndrome delivered between 01 January 1985 and 31 December 2003, of whom 697 were live born and 664 (95%) were traced for their survival status on 28 January 2008.

Main outcome measures Overall prevalence of Down Syndrome. Frequency of additional structural anomalies among cases of Down Syndrome. 10-year survival for all cases of Down Syndrome, by time-period, and by presence, and type, of additional structural anomalies. Independent influence of year of birth, sex, plurality, gestational age, birthweight (standardised for sex, plurality, and

gestational age), maternal age, index of multiple deprivation, prenatal diagnosis, and presence, and type, of additional structural anomalies on survival.

Results Overall prevalence of Down Syndrome was 16.7 per 10 000 registered births (95% CI 15.6 to 17.6). 697 (63%) cases had no additional structural anomalies, 320 (29%) had isolated cardiovascular anomalies, 27 (2%) had isolated digestive system anomalies, 26 (2%) had both cardiovascular and digestive system anomalies, and 31 (3%) had at least one other structural congenital anomaly. 10-year survival among children born with Down syndrome was 83.9% (95% CI 80.9 to 86.5). Survival increased significantly with time ($p < 0.001$), from a 10-year survival of 78.2% (95% CI 72.4 to 83.0) in 1985–1990 to 91.1% (86.7 to 94.1) in 1997–2003. Presence of an additional structural anomaly significantly reduced survival ($p < 0.001$) from a 10-year survival of 94.1% (95% CI 90.9 to 96.1) among those with no additional anomaly to 75.2% (69.5 to 80.0), 68.4% (42.8 to 84.4), 54.2% (32.7 to 71.4), and 76.9% (44.2 to 91.9) among those with cardiovascular anomalies, digestive system anomalies, both cardiovascular and digestive system anomalies, and at least one other structural anomaly respectively. Of the other factors examined, only gestational age ($p < 0.001$) and standardised birthweight ($p = 0.008$) also independently predicted survival.

Conclusion Independent of significant improvement over time, survival of children born with Down syndrome is influenced by gestational age, birthweight, and the presence of additional structural anomalies. This information will be valuable for families and health care professionals when Down Syndrome is detected, and will assist in planning, assessing, and aiding, the future care needs of those affected.

082

MATERNAL EMPLOYMENT AND CHILD SOCIO-EMOTIONAL BEHAVIOUR: LONGITUDINAL EVIDENCE FROM THE MILLENNIUM COHORT STUDY

doi:10.1136/jech.2010.120956.82

A McMunn, Y Kelly, N Cable, M Bartley. *Department of Epidemiology and Public Health, University College London, UK*

Objective To examine the effects of parental employment in the early years on child socio-emotional behaviour at age 5 in a recent birth cohort study in the UK.

Design Prospective cohort study.

Setting The Millennium Cohort Study (MCS): a large, representative sample of children born in the UK between September 2000 and January 2002 ($n = 18\,819$ at sweep 1).

Participants Singleton births in households in which a mother was present in the first three sweeps of the MCS, when participants were 9 months, 3 years and 5 years. Analysis was restricted to white children as there was large ethnic variation in maternal employment, but inadequate power to stratify by ethnicity.

Main outcome measure The Strengths and Difficulties Questionnaire (SDQ) with clinically relevant cut-points for problem behaviours. The SDQ is a widely used instrument for assessing socio-emotional difficulties in children.

Methods Data on parental employment across the three sweeps were used to investigate: (i) whether children whose mothers were in paid work during their first five years were more likely than children whose mothers were at home full-time to display adverse behavioural symptoms at age 5, independent of maternal education, mental health or economic position; (ii) whether effects of maternal employment on child socio-emotional development were cumulative in nature, or whether children were more sensitive to the effects of maternal employment during their first year; and (iii) the effects of different types of parental work arrangements on child socio-emotional behaviour at age 5.

Results No evidence of detrimental effects of maternal employment in the early years on subsequent child socio-emotional behaviour was seen. There were significant gender differences in the effects of parental work arrangements on behavioural outcomes. Girls whose mothers were not in paid work during their first 5 years were 77% (95% CI 1.21 to 2.57) more likely to have behavioural difficulties at age 5 than girls whose mothers were in paid work throughout their early years, independent of maternal characteristics and household income. For boys this was not the case, but boys in two-parent households in which their father was not in paid work for at least one period during their first five years were at an increased risk for behavioural problems at age 5. The most beneficial working arrangement for both girls and boys was that in which both mothers and fathers were present in the household and in paid work, independent of parental educational attainment and household income.

083

IT IS NOT "JUST DEPRIVATION": WHY DO EQUALLY DEPRIVED UK CITIES EXPERIENCE DIFFERENT HEALTH OUTCOMES?

doi:10.1136/jech.2010.120956.83

¹D Walsh, ²N Bendel, ³R Jones, ⁴P Hanlon. ¹Glasgow Centre for Population Health, Glasgow, UK; ²NHS Manchester/Manchester Joint Health Unit, Manchester, UK; ³Liverpool Primary Care Trust, Liverpool, UK; ⁴University of Glasgow, Glasgow, UK

Background The link between socio-economic circumstances and health is well established, with material deprivation cited by UK politicians and policy-makers as the principal underlying cause of poor health and low life expectancy. However, research has shown that not all poor health can be explained purely in terms of deprivation: Glasgow in particular has been cited in this regard, with a "Glasgow effect" suggested to explain the city's high mortality rates. However, Glasgow is not alone in experiencing relatively high levels of poor health and deprivation within the UK: Liverpool and

Manchester are very similar in this regard. Previous analyses of this "effect" were constrained by limitations of data and geography.

Objectives To establish whether there is evidence of a so-called Glasgow effect: (a) even when compared to its two most similar and comparable UK cities; and (b) when based on a more robust and spatially sensitive measure of deprivation than was previously available to researchers.

Participants and setting Total populations of Glasgow, Liverpool and Manchester.

Design Rates of "income deprivation" (a measure very highly correlated with the main UK indices of multiple deprivation) were calculated for small areas in the three cities. All-cause and cause-specific SMRs were calculated for Glasgow relative to Liverpool and Manchester, standardising for age, sex and deprivation decile. In addition, a range of historical census and mortality data were analysed.

Results The deprivation profiles of Glasgow, Liverpool and Manchester are almost identical. Despite this, premature deaths in Glasgow are more than 30% higher, with all deaths around 15% higher. This "excess" mortality is seen across virtually the whole population: most age groups, both males and females, in deprived and non-deprived neighbourhoods. For <65s, SMRs tended to be higher for the more deprived areas, and around a half of "excess" premature deaths were directly related to alcohol and drugs. Analyses of historical data suggest that the deprivation profile of Glasgow has not changed significantly relative to Liverpool and Manchester in recent decades; however, the mortality gap appears to have widened since the early 1970s, suggesting that the "effect" is a relatively recent phenomenon.

Conclusion While deprivation is a fundamental determinant of health and, therefore, an important driver of mortality, it is, however, only one part of a complex picture. As currently measured, deprivation does not explain the higher levels of mortality experienced by Glasgow in relation to two very similar UK cities. Additional explanations are required.