levels of trust and reciprocity were related to non-smoking and adequate duration of sleep, and high levels of social support to adequate duration of sleep and daily use of vegetables. These associations persisted after controlling for age, gender, education and living arrangements. According to our findings, structural social capital seems to be associated with health behaviour more strongly than cognitive social capital.

Conclusion Irrespective of their social status, people with higher levels of social capital—especially in social participation and networks—engage in healthier behaviour. When trying to reduce health inequalities, one strategy could be to promote social participation especially among people in danger of social exclusion.

P32 USING EVIDENCE TO PRIORITISE AREAS FOR PUBLIC HEALTH ACTIONS FOR TACKLING CHILDHOOD OVERWEIGHT

doi:10.1136/jech.2010.120477.32

¹G McNeill, ¹G Osei-Assibey, ¹S Dick, ¹J Macdiarmid, ^{1,2}S Semple, ³J Reilly, ⁴A Ellaway, ²H Cowie. ¹Population Health Section, University of Aberdeen, Aberdeen, UK; ²Division of Developmental Medicine, University of Glasgow, Glasgow, UK; ³MRC Social and Public Health Sciences Unit, Glasgow, UK; ⁴Institute of Occupational Medicine, Edinburgh, UK

Objective To use evidence from systematic reviews of environmental influences on childhood overweight to prioritise areas for development of public health interventions.

Design Systematic literature reviews of intervention and longitudinal studies of environmental factors which could influence overweight via diet and/or physical activity, followed by a questionnaire-based ranking exercise for (a) the strength of the evidence for a causal association between the environmental factor and childhood overweight and (b) the likely effect size of public health actions on each factor on the prevalence of overweight in children. Environmental factors for which there was both strong evidence of causality and a larger effect size were selected for detailed modelling of the likely impact of public health interventions in children in Scotland.

Setting Children up to 8 years in Scotland.

Participants Academics and policy makers with experience of reviewing evidence of public health interventions on obesity.

Main Outcome Measure Ranking of environmental factors to identify those which should be prioritised for future public health interventions.

Results Of the twenty-four environmental factors initially identified at stakeholder workshops, no studies were found for three factors influencing physical activity and one factor affecting diet. For the remaining twenty factors a median of six (range 1–28) studies were identified and summarised in tabular form. Fourteen academics and policy makers were given the tables to complete the rating exercise, which took approximately 1 h. The strength of the evidence and the likely effect size of actions were rated on a scale of 0 (low) to 5 (high). Eight of the environmental factors obtained mean ratings >3 for both strength of evidence and likely effect size: six were factors related to diet (high energy-dense snacks, sugar-sweetened soft drinks, infant feeding, availability of high fat, sugar and salt foods and portion size of manufactured foods and of restaurant and cafeteria items) and two were related to physical activity (physical education in schools and sedentary leisure activity). The ratings were lowest for access to local amenities, safe routes to schools and provision of healthy foods in schools.

Conclusion The ranking exercise provided a structured approach for obtaining a consensus view on priorities for public health action using tables of evidence from a systematic review. Use of this approach with a larger number academics and policy makers from different sectors would be useful to assess whether the area of expertise and sector (academia vs policy) influences the rating of the evidence.

P33 TOWARDS A SMOKE FREE MINISTRY OF HEALTH IN ITALY

doi:10.1136/jech.2010.120477.33

L Spizzichino, D Galeone, L Pagliari, S Mele. *Italian Ministry of Health, Prevention Department, Rome, Italy*

Background In 2009, a health promotion initiative targeted the employees of the Italian Ministry Health aimed to promote compliance with the smoking ban and provide support for any smokers who wanted to quit, by providing free of charge tobacco cessation programme, available during working hours and in the workplace, for all smoking employees who agreed to participate in the programme. Activities included: an information seminar for employees on the risks of smoking; a questionnaire-based survey on the smoking habits of employees in the Ministry of Health head-quarter; the launch of “Groups to quit smoking” run by personnel from the Italian anti-cancer league (LILT).

Results The programme is still ongoing but some results can be highlighted: The seminar held on February was attended by over 200 employees; all employees received an information booklet on the damage caused by smoking; the survey on the smoking habits of employees was completed. Out of the 1244 questionnaires distributed, 1030 were returned (participation rate 82.8%), analyses were carried out on 994 of them. Data showed that 25.1% of employees are smokers, smoking an average of 13.3 cigarettes a day; 43% of them are heavy smokers. Sixty-three percent of smoking employees claim they want to quit, but only 40% of them were advised to do so by their physician. Non-smokers account for 50.5% of employees, and former smokers 24.4%; second-hand smoke is a serious nuisance for 61% of them. Thirty-six percent of non-smokers is exposed to second-hand smoke in different places within the Ministry, including their own room and other rooms (in 2% and 6% of cases, respectively). Four tobacco cessation courses were launched in May 2009. A total of 74 employees (29% of smoking employees) signed up. Of these, 57 participated, and after having completed the course, 32 (56%) quit smoking and 25 (44%) smoked fewer cigarettes a day. Follow-up activities 6 months after the completion of the course are currently ongoing, and the rate of abstinence from smoking is above 37%, in line with data reported in the literature. A new course has been launched in January 2010, 22 employees participated and at the end of the course 32% quit smoking.

Conclusions The program seems to work, law that bans smoking together with cessation program, free of charge and during working hours, are useful to protect non smokers from passive smoking and to help smokers to quit.

P34 DEFINING A TARGET PROFILE FOR PROMOTING SMOKING CESSATION IN ITALY

doi:10.1136/jech.2010.120477.34

¹C Bietta, ²G Ferrante, ²V Minardi, ³L Spizzichino, ³S Vasselli. ¹Local Health Unit, Region Umbria, Italy; ²Istituto Superiore di Sanità, National Centre of Epidemiology, Surveillance and Health Promotion, Rome, Italy; ³Prevention Department, Italian Ministry of Health, Rome, Italy

Background In Italy, smoking cessation programmes rarely consider the characteristics of people who make quit attempts during life (independently from quitting or still smoking) as an useful starting point for effective intervention. Behavioural risk factor surveillance systems can provide information about sociodemographic and health profile of smokers who express intention to quit smoking and therefore represent a potential target of cessation treatment and health promotion intervention.

Objective To evaluate the percentage of people who reported smoking cessation or at least an attempt once in the last 12 months;