Reference Centre. Women aged 50–52 who had a first call invitation (a first invitation to the national screening programme) in this period, and women aged 50–69 who had a routine recall invitation (after previously having a mammography as part of the screening programme) were analysed. Where ethnicity was not known, multiple imputation was used. First call and routine recall data were analysed separately. Screening uptake in different ethnic groups was assessed using logistic regression, and adjusted for age at invitation, socioeconomic deprivation and screening area. Data for the six individual screening areas were also analysed separately.

Results Data on 159,078 women were included in the first call analysis, and on 496,438 women in the routine recall analysis. Ethnicity information was available for 475,478 (72.5%) of these women. Compared with White British women, all other ethnic groups were less likely to attend their first call screening invitation. White British women were also most likely to attend for routine recall screening sessions. Some screening areas showed less variation, with women from several ethnic groups having similar screening uptake to White British women.

Conclusion Breast cancer screening attendance varies by ethnic group for both the first invitation and for subsequent invitations after previously being screened, with White British women more likely to attend. Collaboration between areas to find successful practices for engaging with different communities should improve breast cancer screening uptake.

PS33 HEALTH-SEEKING BEHAVIOUR IN THE ERA OF FREE HEALTHCARE IN URBAN SLUMS IN SIERRA LEONE

doi:10.1136/jech-2012-201753.132

¹L Nathaniel-Wurie, ¹G Martin, ¹G Cooper, ¹G-L De Bernier, ²T Ajayi, ²F Martineau, ²B Cridford, ²S Lako. ¹Global Health Department School of Medicine and Dentistry, King's College London, London, UK; ²Welbodi Partnership, Freetown, Sierra Leone

Background Sierra Leone has child health statistics which are amongst the worst in the world. In April 2010, the Government of Sierra Leone launched a Free Health Care Initiative (FHCI), eliminating user fees in government facilities for children under the age of five, as well as pregnant and lactating women. Data from Ola During Children's Hospital (ODCH), the country's only government-run tertiary paediatric hospital, suggests that the FHCI has increased service utilisation. However, mortality rates remain high and delayed presentation is a major factor contributing to poor outcomes. This study sought to understand health-seeking behaviour in the era of Free Health Care, as a means of addressing barriers to the delivery of timely paediatric care for the urban poor.

Methods Data was collected via focus group discussions and key informant interviews in Freetown. Men and women with at least one child below the age of five under their care were included. Participants were selected from neighbourhoods in Freetown where Ola During Children's Hospital (ODCH) represented the nearest secondary care facility to their usual place of residence. Focus groups were facilitated in Krio, the national language, guided by a pre-designed survey tool. Discussions were audio-recorded, transcribed and translated into English. Data analysis was conducted by the research team using a thematic framework.

Results Preliminary analysis reveals several factors contributing to delays in accessing appropriate care for sick children:

Indirect costs: Transportation, child-minding, lost income, cost of prescribed medicines and unsanctioned charges for care that continue to be levied.

The negative perceptions of ODCH staff and facilities: Perceived disrespectful, discriminatory or unfair treatment within the health facility.

Numerous healthcare options more easily accessible: Private pharmacies, self-treatment at home using traditional (herbal) or allopathic remedies.

Poor insight into disease: Deficiencies in caregivers' ability to recognise at what time during illness progression seeking healthcare would result in the highest cost benefit ratio.

Non caregiver factors: Long queues, perceived complicated referral process between FHC facilities, and reduced opening hours.

Conclusion User fees represent a major barrier to accessing timely and appropriate care for poor children. However, the elimination of official charges for paediatric care in Sierra Leone has brought to the surface several critical factors, beyond direct costs, which also impact upon healthcare utilisation. Understanding the processes and determinants which modulate health seeking behaviour is fundamental to efforts to improve health outcomes for the most vulnerable children.

PS34 RECORDING AND MANAGEMENT OF OBESITY IN PRIMARY CARE, 1997-2006

doi:10.1136/jech-2012-201753.133

HP Booth, M Gulliford. *Department of Primary Care and Public Health Sciences, King's College London, London, UK*

Background Primary care services play a potentially important role in the diagnosis and management of obesity but little evidence exists on current routine practice. The aim of this study was to describe the recording of body mass index and interventions used in the management of body weight in UK primary care in different categories of body weight.

Methods A retrospective cohort study was implemented using the UK General Practice Research Database (GPRD). Participants were aged between 18 and 100 and had a diagnosis of obesity and/or a record indicating a weight management intervention between 1997 and 2006. Diagnoses of obesity were identified by a medical code for obesity or a body mass index (BMI) record of $\geq 30\text{kg/m}^2$. Interventions were identified by medical and therapy codes and included advice on diet, body weight control or exercise, referral to specialist care for obesity or prescription of anti-obesity drugs.

Results Patients ($n=69,102$) with a diagnosis of obesity or who received a weight management intervention were identified at 128 UK general practices. Average BMI rose slightly over the study period from 35.0kg/m^2 in 1997 to 35.8kg/m^2 in 2006, with a higher proportion of patients gaining rather than losing or maintaining their weight each year. Monitoring of BMI was infrequent with just 48% of patients previously identified as obese having a BMI recorded in 2006. The proportion of patients with one or more interventions for weight control recorded increased from 29% in 1997 to 37% in 2006. The most frequent intervention throughout the study period was the provision of dietary advice, recorded in 13,097 (40.7%) patients in 2006 compared to 3,401 (32.3%) in 1997. Prescription of anti-obesity drugs increased by almost 60-fold over the study period. The proportion of morbidly obese patients ($\text{BMI} \geq 40\text{kg/m}^2$) receiving a weight control intervention in 1997 was 38%, rising to 63% in 2006. In obesity category I ($\text{BMI} 30\text{--}34.9\text{kg/m}^2$) the proportion receiving treatment was 43% in 1997 and 53% in 2006 and in category II ($\text{BMI} 35\text{--}39.9\text{kg/m}^2$) 41% and 56% respectively.

Conclusion Monitoring of obesity in diagnosed patients was inconsistent in this population with less than half having a BMI recorded in each year of the study. The frequency of interventions for weight management increased substantially in primary care between 1997 and 2006.

PS35 THE ROLE OF TIME PREFERENCE IN SMOKING CESSATION: A LONGITUDINAL ANALYSIS OF DATA FROM THE HOUSEHOLD INCOME AND LABOUR DYNAMICS OF AUSTRALIA SURVEY, 2001-08

doi:10.1136/jech-2012-201753.134

J Adams, H Brown. *Institute of Health & Society, Newcastle University, Newcastle, UK*

Background 'Time preference' is an economic concept that describes the trade-offs that individuals make between costs and benefits occurring at different points of time. Related psychological concepts include time perspective, impulsivity and delay of gratification. Present orientated people prefer immediate over future gains and place more value on these. Differences in how much value different individuals place on future outcomes are likely to play some role in present day behavioural decisions.

There is substantial cross-sectional evidence that smokers tend to have shorter term time preferences than non-smokers. However, few studies have explored the longitudinal relationship between time preference and smoking cessation.

We explored the relationship between time preference and smoking cessation in a general adult cohort using data from a large Australian longitudinal panel study: the Household Income and Labour Dynamics of Australia (HILDA) survey.

Methods HILDA is a longitudinal, nationally representative, household survey which began in 2001. The focus of the survey is on economic and subjective well-being, and labour market and family dynamics. Data is collected by annual interview and self-complete questionnaire.

Members of the HILDA panel, aged 15–64, who responded to at least four waves of data collection between 2001 and 2008, and reported any level of tobacco consumption at any wave were included in the analyses.

Smoking cessation was measured using self-report questions. Time preference was measured using self-reported time period for financial planning. A range of sociodemographic (age, number of children, various markers of socio-economic position) and smoking-related (previous quit attempts, years smoked) co-variables were controlled for.

Results 1,817 individuals (representing 7,913 unique observations) were included in the analyses. In univariable analyses, both men and women who reported quitting smoking were more likely to have a longer term time preference in the data collection period immediately prior to quitting than those who did not quit (men: $t(4,126)=4.59$, $p<0.001$; women: $t(3,783)=7.18$, $p<0.001$). These relationships persisted after control for socio-demographic and smoking-related covariates (hazard ratio of quitting in those with longer vs shorter term time preference (95% confidence intervals) = 1.27 (1.03 – 1.57) in men; 1.31 (1.05 – 1.63) in women).

Conclusion Adult smokers with a longer term time preference are more likely to quit smoking. This does not exclude the possibility that quitting smoking simultaneously leads to development of longer term time preference. Interventions that encourage development of longer term time preference may enable smoking cessation.

PS36

PROBATION AND NHS WORKING TOGETHER TO REDUCE HEALTH INEQUALITIES: GREAT EXPECTATIONS OR A TALE OF TWO SERVICES

doi:10.1136/jech-2012-201753.135

J Parkes, P Roderick, S Panozzo. *Public Health Sciences & Medical Statistics, University of Southampton, Southampton, UK*

Background Offenders within the community are considered a socially excluded population experiencing poorer health than the general population and difficulties in accessing health care. There is little research focused on Community based offenders supervised by Probation. This study explored probation staff's knowledge, perceptions and attitudes to offender health needs and access to health services.

Methods Using information derived from qualitative research with probation staff and following a pilot, a national on-line questionnaire survey was conducted of probation staff involved in the management of offenders from a geographically representative

sample of Probation Trusts in England and Wales. The questionnaire covered: offender health issues, enablers and barriers to health service access, and the NHS/CJS partnership. Models of good/innovative practice were sought.

Results Sixteen Probation Trusts participated and 611/2810 (22%) responses were obtained. 72% were female, 58% had 6–15 years of experience in probation service. The main offender health issues were substance misuse, mental health (MH), learning disabilities and chronic disease, with alcohol misuse the most important. Provision of easily accessible mental health (for mild–moderate severity) and alcohol treatment services were considered to be inadequate (often/occasional lack of adequate provision 65% alcohol treatment vs 42% drug treatment vs 90% MH).

The main barrier reported was poor communication between probation and the NHS in particular with Primary Care services. It was felt that many offenders had not registered with a GP and that there was little joint working, understanding and education between primary care and probation. In addition the chaotic life-style of offenders lead to them having difficulties in negotiating health services and having appropriate health seeking behaviours, often accessing health services at a point of crisis (A&E). Responsiveness of health services was perceived to be lacking. Provision of in-house services (particularly mental health) was considered to be very useful where available.

Suggested improvements included undertaking health needs assessments to inform joint strategic planning, focus on improvement of alcohol treatment provision; joint education sessions for GPs and Offender Managers, exploration of in-house mental health services; increased access to health trainers, and greater health focus in probation staff training.

Conclusion Probation staff participants were very supportive of improving health of offenders. This study identified some key barriers to addressing the health needs of offenders and their access to services, with practical solutions identified by participants providing implications for policy within the 'new' NHS, health service commissioning, and future research.

PS37

DOES THE ROUTE TO DIAGNOSIS AFFECT OUTCOMES FOR OESOPHAGO-GASTRIC CANCER PATIENTS

doi:10.1136/jech-2012-201753.136

¹T Palser, ²D Cromwell, ³R Hardwick, ⁴S Riley, ⁵K Greenaway, ²J van der Meulen. ¹Clinical Effectiveness Unit, Royal College of Surgeons of UK, London, UK; ²Department of Health Services Research and Policy, LSHTM, London, UK; ³Addenbrookes Hospital, Cambridge, UK; ⁴Northern General Hospital, Sheffield, UK; ⁵The Information Centre for Health and Social Care, Leeds, UK

Background The UK Cancer Reform strategy aims to improve early diagnosis but there is little evidence on how the route to diagnosis and affects outcomes for patients with oesophago-gastric (O-G) cancer. We investigated the relationship between the route to diagnosis, patient characteristics, treatment intent and one-year survival among O-G cancer patients

Methods The study was undertaken in 142 English NHS trusts and 30 cancer networks, with data being prospectively collected on patients diagnosed with O-G cancer between October 2007 and June 2009. Route to diagnosis was defined as general practitioner (GP) referral - urgent or non-urgent, hospital consultant referral, or after an emergency admission. The association between patients' route of diagnosis, treatment intent and survival was estimated using logistic regression.

Results Among 14,102 cancer patients, 66.3% were diagnosed after a general practitioner (GP) referral, 16.4% after an emergency admission, and 17.4% after hospital consultant referral. Of the 9,351 GP referrals, 68.8% were urgent. Compared to urgent GP referrals, a markedly lower proportion of patients diagnosed after emergency admission had a curative treatment plan (36% v 16%;