

Short reports

General practice consultation in childhood in The Netherlands: sociodemographic variation

Marc A Bruijnzeels, Johannes C van der Wouden, Marleen Foets

Consultation with a GP is the point of entry into the Dutch health care system. We studied the sociodemographic determinants of GP consultation rates for children. The determinants that are examined are age, gender, socioeconomic status, health insurance, degree of urbanisation, and ethnicity.¹

Methods: We used data for children 0 to 14 years from the Dutch national survey of morbidity and interventions in general practice, carried out by The Netherlands Institute of Primary Health Care (NIVEL) between April 1987 and March 1988.² A random, non-proportionally stratified sample of 161 GPs was drawn. All persons listed in the participating practices provided basic sociodemographic data. The 161 GPs (103 practices) were divided into four groups of approximately 40 GPs. Each group registered every contact between the practice and patient during three consecutive months and in this way one whole year was covered. Registration was episode-oriented. Consultations concerning the same health problem were grouped into episodes of care. We considered only episodes in which the first contact took place within the registration period for a new or relapse problem. The diagnosis considered in this study is the diagnosis made in the last contact registered for the episode. To assess the nature of the problem the episode diagnoses are grouped into clusters. The clusters of interest are acute somatic

complaints, infections, chronic problems, and traumas.

We compared odds ratios (OR) resulting from bivariate logistic regressions and a multivariate logistic regression to assess the contributions of all sociodemographic determinants. In addition, the possible interaction effects between socioeconomic status, insurance coverage, ethnicity, and degree of urbanisation were evaluated.

Results: The table shows that 32% of all children consulted the GP at least once during three months. Younger children, boys, children from lower socioeconomic strata, children insured through the sick funds, children living in the larger cities, and non-Dutch children consulted the GP more often. The association between ethnicity and consultation rate, however, disappears in a multivariate analysis. For three of four different diagnosis clusters the relations show the same direction with different magnitudes. For trauma, older children and children in smaller cities yield higher consultation rates.

The analysis including a possible interaction effect shows that the largest difference in consultation rates existed between children who were insured through the sick funds and lived in the larger cities (39%) on one hand and children who were privately insured and lived in the smaller cities (26%) on the other. Within these two groups children whose socio-

Department of
General Practice,
Erasmus University
Rotterdam,
P.O. Box 1738,
3000 DR,
Rotterdam,
The Netherlands
M A Bruijnzeels
J C Van der Wouden

Netherlands Institute
of Primary Health
Care, Utrecht,
The Netherlands
M Foets

Correspondence to:
Mr M A Bruijnzeels.

Accepted for publication
February 1995

J Epidemiol Community Health
1995;49:532-533

Percentages, bivariate, and multivariate odds ratios (OR) of children who consulted the GP for all episodes and in relation to morbidity category by sociodemographic factors

Sociodemographic factor	All episodes (32%)			Morbidity category (multivariate OR)			
	%	Bivariate OR	Multivariate OR	Acute somatic (8.6%)	Infections complaints (20.9%)	Chronic problems (10.1%)	Traumas (4.3%)
Age (reference category: 0-4 (y))	43	1.00					
5-9	31	0.60**	0.60**	0.92*	0.55**	0.64**	0.98 NS
10-14	24	0.42**	0.42**	0.74**	0.26**	0.73**	1.32**
Gender (reference category: Boys)	32	1.00					
Girls	33	1.04*	1.04*	1.19**	1.03 NS	0.99 NS	0.81**
Socioeconomic status (reference category: high)	29	1.00					
Low/middle	33	1.24**	1.15**	1.11*	1.22**	1.07 NS	1.12 NS
Health insurance (reference category: private)	28	1.00					
Sick funds	35	1.40**	1.31**	1.34**	1.27**	1.25**	1.20**
Urbanisation (reference category: <50 000)	31	1.00					
>50 000 inhabitants	36	1.21**	1.23**	1.12**	1.25**	1.14**	0.85*
Ethnicity (reference category: Dutch)	32	1.00					
Non-Dutch†	36	1.14**	0.99 NS	0.99 NS	1.04 NS	1.00 NS	0.86 NS

NS, not significant; * p<.05; ** p<.01

† A child is considered non-Dutch if either the child or one of its parents is born in Turkey, Morocco, Surinam or the Netherlands Antilles or its has one of these nationalities.

economic status was low consulted more often than those whose status was high socioeconomic.

Conclusions: Three conclusions may be drawn from this study. Firstly, there are relatively large differences in consultation rates between different subgroups. The consultation rate for children who live in a large city and are insured through the sick funds is 1.5 times that of children who live outside the large cities and who are privately insured. This difference has implications not only for the children but also for the burden on the health care system in different areas. GPs who work in socioeconomically deprived areas in larger cities are consulted more often for children than their colleagues in better off areas.

Secondly, if the relation between ethnicity and consultation rate is studied multivariately,

the effect of ethnicity disappears. In The Netherlands allochthonous children consult the GP more often than all autochthonous children, but not more often than autochthonous children who live under the same socioeconomic and urban circumstances. The cultural differences between Dutch and autochthonous children do not affect their consultation rates.³

Thirdly, the subdivision into morbidity categories shows some small deviant relations. This subdivision seems relatively unimportant, when studying GP consultation.⁴

1 Starfield B, Berg BJ van den, Steinwachs DM, Katz HP, Horn SD. Variations in utilization of health services by children. *Pediatrics* 1979;63:633-41.

2 Foets M, Velden J van der, Bakker D de. *Dutch national survey of general practice. A summary of the survey design*. Utrecht: Netherlands Institute of Primary Health Care, 1992.

3 Senior PA, Bophal R. Ethnicity as a variable in epidemiological research. *BMJ* 1994;309:327-30.

4 McCue Horwitz S, Morgenstern H, Berkman LF. The use of pediatric medical care: a critical review. *J Chron Dis* 1985;38:935-45.