

Acute health effects of the Sea Empress oil spill

Ronan A Lyons, J Mark F Temple, Daphne Evans, David L Fone, Stephen R Palmer

Abstract

Study objective—To investigate whether residents in the vicinity of the Sea Empress tanker spill suffered an increase in self reported physical and psychological symptoms, which might be attributable to exposure to crude oil.

Design—Retrospective cohort study; postal questionnaire including demographic details, a symptom checklist, beliefs about health effects of oil and the Hospital Anxiety and Depression and SF-36 mental health scales.

Setting—Populations living in four coastal towns on the exposed south Pembrokeshire coast and two control towns on the unexposed north coast.

Patients—539 exposed and 550 unexposed people sampled at random from the family health services authority age-sex register who completed questionnaires.

Main results—Adjusted odds ratios for self reported physical symptoms; scores on the Hospital Anxiety and Depression and SF-36 mental health scales, in 1089 people who responded out of a possible 1585 (69%).

Conclusions—Living in areas exposed to the crude oil spillage was significantly associated with higher anxiety and depression scores, worse mental health; and self reported headache (odds ratio = 2.35, 95% CI 1.56, 3.55), sore eyes (odds ratio = 1.96, 95% CI 1.06, 3.62), and sore throat (odds ratio = 1.70, 95% CI 1.12, 2.60) after adjusting for age, sex, smoking status, anxiety, and the belief that oil had affected health. People living in exposed areas reported higher rates of physical and psychological symptoms than control areas. Symptoms significantly associated with exposure after adjustment for anxiety and health beliefs were those expected from the known toxicological effect of oil, suggesting a direct health effect on the exposed population.

(J Epidemiol Community Health 1999;53:306-310)

On the evening of the 15 February 1996, the Sea Empress laden with more than 130 000 tonnes of light crude oil, ran aground on rocks at the entrance of Milford Haven harbour in south west Wales. Over the next week an estimated 72 000 tonnes of crude oil and 360 tonnes of heavy fuel oil were released into the sea, contaminating around 200 km of coastline (fig 1) and causing strong smells and complaints of symptoms from residents of the coastal towns. Similar oil spills from the tankers Exxon Valdez¹ and Braer² were fol-

lowed by increases in physical and psychological symptoms in the general population. We undertook a population based retrospective cohort study to assess the acute physical and psychological health impact on the exposed population.

Methods

This study was commissioned four weeks after the incident. The exposed group was defined as people living on the most exposed south coast of Pembrokeshire (Milford Haven, Pembroke Dock, Tenby, and Saundersfoot) and the control group as similar but unexposed populations living on the north coast (Aberaeron and Fishguard) (fig 1).

Sample size calculations were based on the prevalence of symptoms in the control group reported in the Braer study.² A sample of 814 in both exposed and control areas was required for 80 per cent power to detect a statistically significant ($p < 0.05$) doubling of prevalence of symptoms in the exposed group against a background prevalence of 3 per cent. To allow for non-response, random samples of 1000 adults (of both sexes aged 18 to 65 years old) were selected from each of the exposed and control areas using the Dyfed-Powys family health services authority age-sex register.

A questionnaire, with a retrospective health diary covering the four weeks immediately after the incident, was devised that also incorporated a symptom check list used in the Braer investigation,² the Hospital Anxiety and Depression Scale (HAD),³ the mental health profile of the SF36⁴ as well as a series of questions relating to beliefs about the effects of the oil spill on the environment, employment, health, and finances. The HAD Scale is widely used in clinical settings and was included in the study because of the marked excess of cases of clinically relevant anxiety and depression after the Exxon Valdez oil spill.² The SF-36 mental health scale was included as population normative baseline data of SF-36 health scores was available from the Pembrokeshire health status study⁵ undertaken in 1994 in randomly sampled adults from North ($n = 659$) and South Pembrokeshire ($n = 597$).

Questionnaires were mailed to people on the 4 April (seven weeks after the grounding of the tanker). Non-responders were sent a second questionnaire one month later. Non-responders to the second questionnaire were contacted by telephone or visited at home by two of us (MT and DE) between 21 May and 18 June.

Statistical analysis was carried out using SPSS for Windows⁶ and Epi Info Logistic.⁷ Crude odds ratios for each self reported

Iechyd Morgannwg
Health
R A Lyons

Gwent Health
Authority
J M F Temple
D L Fone

University of Wales
Swansea
D Evans

Welsh Combined
Centres for Public
Health
S R Palmer

Correspondence to:
Professor S R Palmer, Welsh
Combined Centres for
Public Health, University of
Wales College of Medicine,
Abton House Wedal Road,
Cardiff CF4 6QX.

Accepted for publication
19 October 1998




Figure 1 South west Wales showing extent of oil pollution and locations of study and control towns.

symptom in the exposed group were calculated with 95 per cent confidence intervals. Other categorical data were analysed using the χ^2 test and continuous or ordinal data using either the t test or non-parametric analysis of variance as appropriate. We carried out a forwards logistic regression analysis for each symptom in turn as the dependent variable, adjusting firstly for age, sex, HAD anxiety score, exposed/control status and smoking status, and secondly, including the health belief effect variable. The Hosmer-Lemeshow goodness of fit test⁷ was used to assess the effect of inclusion of the interaction term between anxiety and health belief effect in each model. Inspection of the data showed substantially high rates of reported illness in the exposed population on day one (16 February) (32 of 539 and 6 of 550 in exposed and control areas respectively). We considered this most likely represented biased recall, and therefore logistic regression analysis was undertaken after exclusion of those subjects.

The dates of exposure to oil were estimated from the Sea Empress Environmental Evaluation Committee's Report.⁸

To estimate the impact of the oil on self reported illness in the exposed population the expected rate of illness was calculated from the control population. The observed minus the expected was plotted for the two groups of exposed populations.

Results

After excluding those who were positively confirmed as not resident at their home address during the relevant period (211 and 204 people in control and exposed populations respectively), the overall response rate was 1089 of 1585 (69%), with no significant difference between the exposed areas 68% (539 of 796) and control areas 70% (550 of 789). There was no difference in mean age (exposed mean age 40.3 years, control 40.6, range 18–65) or the female to male ratios (both 5.5: 4.5) between the two groups. The reported prevalence of "ever smokers" (six months or more) was higher in the exposed area (245 of 539: (45.5%) *v* 206 of 550 (37.5%) in the control area χ^2 7.186 $p = 0.0074$), with no significant difference for current smoking ((28.2%) *v* (25.3%), $p = 0.28$).

In the population resident in the exposed area, responders reported excess symptoms and non-specific illness on the days when oil exposure was at its highest. Figure 2 demonstrates this for all self reported illness. The oil came ashore within the Haven by 17 February, but did not approach the western shoreline of Carmarthen Bay until 22 February. The oil persisted in the western Camarthen Bay until after the study period. In the Haven area daily rates of reported headache peaked on 16 February, the day after the initial oil spill, and 23 February, one to two days after the major secondary spill.




Figure 2 Excess (observed-expected) of self reported illness in area exposed to oil from Sea Express.

In the exposed area 55 of 539 responders reported consulting their general practitioner about any symptoms, within four weeks of the Sea Express grounding, compared with 24 of 550 in the control area. (Yates's corrected $\chi^2 = 12.95$, $df = 1$, $p = 0.0003$). The crude relative risk of a person consulting was 2.34 (95% confidence intervals 1.47 to 3.72). No data are available on consulting rates before the incident.

After standardising for age and sex using 1994 baseline normative data, the difference in expected mean SF-36 mental health score for exposed (75.14) and control groups (74.83) was not statistically significant. After the incident the mean SF-36 mental health score was significantly lower in the exposed population (73.44 v 77.08, $p=0.002$) (table 1).

The four week period prevalence of almost all symptoms experienced between 16 February and 15 March was significantly increased in the exposed population (table 1). In the exposed area 39.3% reported one or more

symptoms compared with 20.4% in the control areas ($p<0.001$). The exposed population reported significantly more anxiety and depression, (HAD score >10 for each sub-scale) (table 1). Anxiety score >10 was associated with increased prevalence of most symptoms in both exposed and unexposed groups. In the exposed group 33 of 69 (47.8%) anxious subjects reported general illness, compared with 84 of 419 (20.0%) others ($\chi^2 = 13.3$ $df=1$ $p=0.0003$). In the control group the proportions were 9 of 50 (18.0%) compared with 30 of 455 (6.6%) (Fisher's exact test $p=0.03$). Adjustment for the effects of anxiety scores was made by logistic regression analysis. Odds ratios for all symptoms except diarrhoea, vomiting and skin blisters remained significantly raised in exposed subjects, adjusting for age, sex, HAD score and ever smoking status (table 2). The belief that the oil spill had affected their health was reported by 12 unexposed people (2.4%) compared with 112 (23.3%) of those exposed ($p < 0.001$). After inclusion of this

Table 1 Four week period prevalence of symptoms in exposed and unexposed population

Symptom	Exposed	Unexposed	O/Rs	95% CI	p Value
Generally ill	125/525 (23.8%)	40/541 (7.4%)	3.91	2.63, 5.85	<0.001
Headache	175/461 (38.0%)	66/469 (14.1%)	3.74	2.67, 5.24	<0.001
Nausea	68/419 (16.2%)	32/461 (6.9%)	2.60	1.63, 4.16	<0.001
Vomiting	15/396 (3.8%)	14/454 (3.1%)	1.24	0.55, 2.77	0.57
Diarrhoea	41/407 (10.1%)	25/455 (5.5%)	1.93	1.11, 3.35	0.01
Sore eyes	84/427 (19.7%)	27/457 (5.9%)	3.90	2.41, 6.35	<0.001
Runny nose	105/431 (24.4%)	62/470 (13.2%)	2.12	1.47, 3.05	<0.001
Sore throat	143/454 (31.5%)	58/471 (12.3%)	3.27	2.29, 4.68	<0.001
Cough	103/431 (23.9%)	53/466 (11.4%)	2.45	1.67, 3.58	<0.001
Itching skin	56/403 (13.9%)	26/455 (5.7%)	2.66	1.59, 4.48	<0.001
Rash	36/402 (9.0%)	16/453 (3.5%)	2.69	1.41, 5.18	<0.001
Blisters	6/389 (1.5%)	2/449 (0.4%)	3.50	0.63, 25.55	0.10
Shortness of breath	56/408 (13.7%)	24/454 (5.3%)	2.85	1.68, 4.86	<0.001
Weakness	117/432 (27.1%)	70/469 (14.9%)	2.12	1.50, 3.00	<0.001
Definite anxiety	72/502 (14.3%)	51/511 (10.0%)	1.51	1.01, 2.26	0.03
Mean anxiety score	5.44 SD 4.54 (SE 0.186)	4.82 SD 4.21 (SE 0.203)			0.036
Definite depression	27/505 (5.3%)	20/520 (3.8%)	1.41	0.75, 2.67	0.3
Mean depression score	3.33 SD 3.83 (SE 0.142)	2.70 SD 3.23 (SE 0.171)			0.049
Mean SF-36 mental health score	73.4 SD 19.5 (SE 0.869)	77.1 SD 18.39 (SE 0.814)			0.002

Table 2 Logistic regression odds ratios for symptoms in exposed population adjusted for age, sex, anxiety score, and smoking status

Symptom	Odds ratio (95% CI)	p value
Generally ill	3.5 (2.25, 3.50)	<0.0001
Headache	3.93 (2.70, 5.71)	<0.0001
Nausea	2.41 (1.44, 4.03)	<0.001
Vomiting	0.97 (0.40, 2.34)	0.95
Diarrhoea	1.56 (0.86, 2.83)	0.14
Sore eyes	3.59 (2.12, 6.07)	<0.0001
Runny nose	1.97 (1.33, 2.91)	<0.0001
Sore throat	2.89 (1.99, 4.20)	<0.001
Cough	1.94 (1.30, 2.91)	<0.001
Itching skin	2.31 (1.33, 4.03)	<0.01
Rash	2.26 (1.13, 4.51)	0.02
Blisters	5.99 (0.69, 52.0)	0.10
Shortness of breath	2.31 (1.27, 4.19)	<0.01
Weakness	2.04 (1.44, 2.89)	<0.001

Table 3 Logistic regression odds ratios adjusting for age, sex, smoking status, anxiety score, and health belief effect

Symptom	Odds ratio (95% CI)	Hosmer-Lemeshow goodness of fit p value*
Headache	2.35 (1.56, 3.55)	0.18 (0.11)
Sore eyes	1.96 (1.06, 3.62)	0.85 (0.002)
Sore throat	1.70 (1.12, 2.60)	0.15 (0.01)

*Goodness of fit p value without interaction term (anxiety \times health effect belief) in the model.

variable in each logistic regression model, the odds ratios for headache (OR = 2.35, 95% CI 1.56, 3.55, $p < 0.001$), sore eyes (OR = 1.97, 95% CI 1.08, 3.60, $p < 0.03$), and sore throat (OR = 1.65, 95% CI 1.08, 2.52, $p < 0.02$) remained significantly increased (table 3). In each of these three models the interaction term of anxiety score and health effect belief significantly increased the goodness of fit.

Discussion

This is the first reported study of urban populations exposed to a large oil spill. In previous studies^{1,2} the incident has occurred away from large centres of population. We found an increase in self reported psychological and physical symptoms in the exposed south Pembrokeshire population during the four weeks after the incident. Self reported headaches, sore eyes, and sore throats were significantly associated with exposure to crude oil after adjusting for age, sex, smoking status, anxiety, and health belief effects. The exposed population reported significantly more anxiety and depression measured by the Hospital Anxiety and Depression scale with mean SF-36 mental health scores 3.6 lower than the control population. The magnitude of this difference in the population mean mental health score is comparable to the difference between Pembrokeshire and the most socially disadvantaged areas of Wales as measured in the 1995 Welsh Health Survey.⁹

Before this excess of symptoms can be attributed to exposure to the oil spill it is necessary to consider possible biases in the study. Response rates were very similar in the exposed and control areas (68% v 70%); the rate of headaches in control non-responders would have to be more than five times the rate in responders to render the difference in control and exposed populations non-significant. The main threat to the validity of our findings

KEY POINTS

- People living in towns exposed to the oil spill had significantly higher self reported illness including anxiety and depression than those living in control towns.
- After adjustment for the effect of anxiety score, age, sex, and smoking status significant associations remained with headache, sore eyes, and sore throat.
- Headache, sore eyes, and sore throat are the symptoms predicted to be associated with exposure to crude oil on the basis of known toxicological effects.

comes from recall and rumination bias leading to exaggeration of symptoms in the exposed population. This may explain the excess of illness reported by the exposed population on day one of the incident; very little oil escaped within the first 12 hours of the ship running aground. So the excess of symptoms recorded on the first day cannot be attributed to the effect of the oil. However, even when persons reporting illness on day one were excluded from the analyses, most symptoms were still significantly more common in the rest of the exposed group. Increased anxiety because of fear, or the potential impact on employment might be expected to affect the reporting of symptoms but the increased prevalence of many symptoms was found to be independent of anxiety score.

Predictably, a higher proportion of people in the exposed areas reported that they believed the oil spill had affected their health (23% v 2%) and it is possible that those with this belief would tend to over-report symptoms. Adjustment for its effect by removing rumination bias may give a more reasonable estimate of risk. Rates of headaches, sore eyes and sore throat still remained statistically higher in the exposed population. We conclude that after conservative allowances are made for bias, the physical health of the exposed group remained significantly worse than the control population. Had data on individual exposures been available a dose response relation would have strengthened this conclusion.

Is the higher prevalence of symptoms (after adjusting for anxiety and health belief effects) a direct effect of exposure to oil or to the detergents used in the clean up, or a non-specific response to an environmental threat? Perceived exposure to an environmental threat, may result in an excess of reported physical symptoms in the absence of a chemical effect. For example, in Worcester, UK in 1995 the water supply to 160 000 people was contaminated with low doses of dioxanes.¹⁰ Within the group noticing an unusual taste to the water there was an excess of symptoms.

However, the distribution of the water supply was such that study subjects who did or did not notice an unusual taste were equally exposed to contaminated water. The authors concluded that the observed increase in reported symptoms was because of public anxiety caused by the incident rather than the

direct effect of the exposure. In Texas more than 1000 people attended the emergency departments of hospitals after 40 000 lb of hydrofluoric acid was released into the atmosphere.¹¹ High exposure was associated with an excess of physical symptoms but the presence of some physical symptoms was better explained by the degree of psychological distress. However, the physical symptoms significantly associated with exposure after adjusting for psychological distress were those for which there was a biologically plausible relation with hydrofluoric acid.

In our study, the increase in the reported prevalence of headaches, sore eyes, and sore throat after day one was independent of health beliefs and anxiety levels; these symptoms are compatible with both exposure to crude oil and to detergents (personal communication, Welsh National Poisons Unit). Furthermore, and importantly, our findings are similar to those found after the Braer oil spill in Shetland, Scotland in 1993.² The Braer study, based on 420 exposed people and 92 controls, reported significantly higher rates of headache, throat irritation, and itchy eyes particularly in the two days after the oil spill.

Little is known about the longer term psychological effects on the general population of coastal oil spills. One year after the Exxon Valdez oil spill in Alaska a survey of 599 people in 15 communities was carried out to measure psychological effects.¹ Residents from high exposure communities were 3.6 times as likely to have generalised anxiety disorder, 2.9 times as likely to have post-traumatic stress disorders, and 2.1 times as likely to have a high depression score. A follow up study of the Braer oil spill reported significantly higher scores on the general health questionnaire in the exposed group (23.6% *v* 3.4% above threshold scores for mental distress).¹²

In our study, the exposed groups also had significantly higher anxiety and depression scores as measured by the HAD scale and worse mental health as measured by the SF-36 mental health scale.³⁻⁴ Whether the psychological effects are because of a toxic effect of the oil or are a generalised response to an environmental disaster with potential impacts on health, employment, income, and perception of the environment is unknown. Given the experience of the Worcestershire water incident

and the Texas hydrofluoric acid episode it would seem probable that much, if not all, of the psychological effect is a generalised response to environmental disasters. These studies suggest that in environmental incidents adjustment should be made for psychological effects and health beliefs, as we did in our study.

Overall this study shows that, like previous similar incidents, the Sea Empress oil spill resulted in higher levels of psychological and physical symptoms in the exposed populations of south Pembrokeshire. Even after adjusting for the psychological effects and health beliefs of residents, there was a significant association found between exposure and biologically plausible symptoms. The plausibility is based on the known toxicological effects of crude oil. The coherence of these results suggests that the association is causal, resulting in a direct health effect on the exposed population. It also suggests that the method used in our study to adjust for psychological effects and health beliefs should be considered for use in epidemiological studies in future major environmental incidents. Further studies are planned to assess whether there are any longer term sequelae after this incident.

Conflicts of interest: none.

- 1 Palinkas LA, Peterson JS, Russel J, et al. Community patterns of psychiatric disorders after the Exxon Valdez oil spill. *Am J Psychiatry* 1993;150:1517-23.
- 2 Campbell D, Cox D, Crum J, et al. Initial effects of the grounding of the tanker Braer on health in Shetland. The Shetland Health Study Group. *BMJ* 1993;307:1251-5.
- 3 Zigmond AS, Snaith RP. The Hospital Anxiety and Depression Scale. *Acta Psychiatr Scand* 1983;67:361-70.
- 4 Ware JE, Sherbourne CD. The MOS 36 Items Short-Form Health Survey (SF-36): 1. Conceptual framework and item selection. *Med Care* 1992;30:473-80.
- 5 Monaghan S. The Pembrokeshire Health Status Survey. *South East Pembrokeshire Project: Technical Document 1*, Haverfordwest: Pembrokeshire Health Authority, 1994.
- 6 Norusis MJ. *SPSS for Windows Release 6.0*. Chicago: Statistical Package for the Social Sciences, 1993.
- 7 Dallal GE. LOGISTIC: A Logistic Regression Program for the IBM PC. *The American Statistician* 1988;42:272.
- 8 Sea Empress Environmental Evaluation Committee. *The environmental impact of the Sea Empress oil spill*. London: HMSO, 1997.
- 9 Welsh Office. *1995 Welsh Health Services*. Cardiff: Welsh Office, 1996.
- 10 Fowle SE, Constantine CE, Fone D, et al. An epidemiological study after a water contamination incident near Worcester, England in April 1994. *J Epidemiol Community Health* 1995;50:18-23.
- 11 Dayal HH, Baranowski T, Li Y, et al. Hazardous chemicals: psychological dimensions of the health sequelae of a community exposure in Texas. *J Epidemiol Community Health* 1994;48:560-8.
- 12 Campbell D, Cox D, Crum J, et al. Later effects of grounding of tanker Braer on health in Scotland. *BMJ* 1994;309:773-4.