RESEARCH REPORT

Variation of health status among people living on boats in Hue, Vietnam

Nguyen Khac Luong Quang, Takehito Takano, Keiko Nakamura, Masafumi Watanabe, Tomoko Inose, Yoshiharu Fukuda, Kaoruko Seino

J Epidemiol Community Health 2005;59:941–947. doi: 10.1136/jech.2005.034728

See end of article for authors' affiliations

Correspondence to: Professor T Takano, Health Promotion/International Health, Graduate School of Tokyo Medical and Dental University, Yushima 1-5-45, Bunkyo, Tokyo 113-8519, Japan; takano.hlt@tmd.ac.jp

Accepted for publication 16 May 2005

Objectives: To examine patterns of disease and injury in people living on boats in Hue City, Vietnam, and their relations to socioeconomic conditions, sanitary practices, disease prevention proficiency, and people's preference to continued living on boats.

Methods: The subjects were 3737 people aged 5 years and over living on boats in Hue City, Vietnam. Diseases and injuries were diagnosed according to ICD-10. The associations between disease/injury and socioeconomic conditions, sanitary practices, disease prevention proficiency, and preference to continued living on boats were analysed by logistic regression.

Main results: The prevalence rates of certain infectious and parasitic diseases, diseases of the respiratory system, diseases of the skin and subcutaneous tissue, diseases of the digestive system, and injuries were 85.3%, 78.0%, 51.2%, 15.4%, and 13.2%, respectively. Various associations were seen between diseases/injuries and socioeconomic conditions. Patterns of disease were strongly influenced by sanitary practices. Better disease prevention proficiency was significantly related to lower prevalence of the first three categories of diseases/injuries regardless of sex, age, or socioeconomic status ($p < 0.05$, $p < 0.001$, $p < 0.001$, respectively). Diseases were more prevalent among people who preferred not to continue living on boats.

Conclusions: This large scale comprehensive field study illustrated major diseases and injuries among people living on boats. Variations in health status showed a web-like relation of socioeconomic conditions, sanitary practices, disease prevention proficiency, and preference to continued living on boats. Measures to develop disease prevention proficiency reduce the risk of disease and injury.

The rapid urbanisation of the developing world is affecting health in these areas, as well as affecting the living environments and lifestyles that people have had for hundreds of years. The United Nations projects an average population growth of 2.3% per year in urban areas of the developing world during 2000–2030.¹ In Asia, 39% of the population currently lives in urban areas, and this percentage is expected to reach close to 55% by 2030.¹

In common with other cities in Asia, Hue City, Vietnam, is undergoing rapid urbanisation with a population growth of 100% in the past quarter of century.² The rapid urbanisation in Hue places a great deal of demand on urban infrastructure, housing, quality living environments, employment, health promotion, and community development projects in the area.^{2–3} Two long wars experienced in Hue resulted in much lasting damage to both infrastructure and the environment.²

In Hue, fragile living is common among people living on boats along the Huong River, which has traditionally provided a convenient place for life on the water. People living on boats are underprivileged with regard to health, education, income, household space, access to drinking water, and other living conditions, and have needs for improvements in their health.⁴

The health of the population is influenced by multiple factors, including socioeconomic concerns, the physical and social environment, disease prevention practices, personal skills, and people's attitudes towards health.^{5–6} In society, the associations between health and a wide range of health influencing factors identified in individual populations are characterised by multicausal relations.⁷ Evidence reflecting these web-like causal relations among specific factors influencing the health of a particular population will provide a basis for measures to improve the health of the population.

To consider the health of people under fragile living conditions, solid evidence on variation in health and disease in this population is required. However, systematic boat household studies have not been carried out to quantify disease and injury in Hue City. Such quantitative analyses and examinations of factors related to their health would provide evidence to facilitate the planning for health developments. In this study, we performed a household survey focusing on a wide range of variables, including demographic, socioeconomic, sanitary practices, disease prevention proficiency, and health status.

The objectives of this study were to determine disease and injury patterns of people living on boats along the Huong River in Hue City and their relations with socioeconomic conditions, sanitary practices, disease prevention proficiency, and people's preference regarding continued living on boats.

METHODS

Study area and subjects

The study area consisted of six quarters of Hue City—Phu Hau, Phu Hiep, Vy Da, Phuong Duc, Kim Long, and Phu Cat—which included 941 households living on boats according to the household registration records. The subjects of health examination were people aged 5 years and over from boat households.

Procedures

Ten interview teams were assigned to visit boat households and conducted interviews and health examinations from 4 August to 19 September 2003. A team of three people—one medical doctor, one nurse, and one civil servant—met the heads of household and carried out face to face interviews

using structured questionnaire forms. The medical doctor and nurse examined health conditions of household members aged 5 years and over. The interview team members participated in a two day training workshop in advance, to obtain appropriate knowledge and skills for the interview and health examination as well as to understand the objectives and protocol of the study.

Permission to implement the study was obtained from Hue People's Committee by consideration of ethical appropriateness. Informed consent from the heads of households and household members were obtained. The heads of the quarters and key people in the community were informed of the purposes and expected outcomes as well as general protocol of the study, in advance.

Questionnaire survey

Structured questionnaire forms were prepared in Vietnamese, and were pre-tested in the field to ensure that all questions could be understood clearly. The interview method was standardised through practice visits to boat households by the interview teams. The following items were assessed.

Demographic and socioeconomic conditions

Age, sex, family size, education level, income, and occupation. Family size was categorised as non-crowded (1–4 people), crowded (5–7), and overcrowded (8 or more). Education level was classified into three categories (never went to school, primary school, and secondary school or higher education). Economic level was divided into two categories (poor, non-poor) according to a national poverty line, less than 150 000 Vietnam dong (VND) (equivalent to 10 US dollars) or 150 000 VND or more per capita average monthly income.⁸ Individual occupations were identified and categorised as occupations requiring boats, occupations not involving boats, or inactive.

Sanitary practices

Water treatment before drinking; washing of hands before eating; heating foods before eating; sources of water for cooking, drinking, bathing and washing; defecation sites; and garbage disposal sites.

Disease prevention proficiency

A “disease prevention proficiency score” was developed based on the subjects' knowledge regarding preventable diseases, knowledge of disease prevention practices, and practices preferable for disease prevention. Subjects were categorised into three groups based on the disease prevention proficiency score: non-functional (0 point), minimally functional (1–4 points), and functional (5–11 points).

Preference to continued living on boats

Subjects were asked whether they would prefer to continue living on boats, and their response was categorised as either affirmative or negative.

Health examination

Diseases and injuries in the past 12 months and currently were assessed by clinical medical doctors and coded according to the 10th version of International Statistical Classification Diseases and Related Health Problems (ICD-10).⁹ A 12 month period was chosen to exclude the influence of variations in disease and injury prevalence relating to season. Four major disease categories (certain infectious and parasitic diseases, diseases of the respiratory system, diseases of the skin and subcutaneous tissue, and diseases of the digestive system) and injuries were recorded according to the disease prevalence records of Hue City. Selected specific

diseases and injuries among major diseases and injuries were also recorded.

Statistical analyses

The distributions of age, education, and occupation of the subjects were calculated. The distributions of occupation by education and income level were analysed by χ^2 test. The prevalence rates of the four major disease categories and injuries as well as selected specific diseases and injuries were quantified. Differences of the prevalence rates of major diseases and injuries by sex and age group children (5–14 years) compared with adults (15 years and over) and young adults (15–49) compared with older adults (50 years and over) were examined by χ^2 test. The sex and age adjusted odds ratios (OR) and 95% confident intervals (CI) of diseases and injuries for different family sizes, socioeconomic conditions, sanitary practices, and preference to continued living on boats were calculated by logistic regression analysis. ORs and 95% CIs of major diseases and injuries for increments of 1 in disease prevention proficiency score were calculated by logistic regression analysis. Three models were used: adjustment by sex and age; adjustment by sex, age, and occupation requiring/not requiring boats; and adjustment by sex, age, occupation requiring/not requiring boats, and education.

RESULTS

Profiles of the subjects

The results from 782 household visits were compiled for the analyses (83.1% response rate). A total of 3737 subjects participated in the health examination (73.8% of all reported family members aged 5 years and over). According to the reports made by the heads of the households, their households were made up of 2777 males and 2785 females with mean ages of 24.1 and 25.0 years, respectively. The median household size was seven people and 70.5% of the households were comprised of more than five people.

Table 1 shows age distribution, education, and occupation profiles of the subjects. Of the subjects aged 15 years and over, 39.2% had never attended school, while 33.8% had attended only primary school. Fishing, sand extraction, and transportation were the major occupations requiring boats, while vending, cyclo driver, and general labour were the major occupations that did not involve boats. “Poor” subjects made up 52.6% of the study population. The distribution of occupations requiring compared with those not requiring boats differed by education level; 49.7% of subjects who had participated in secondary school and above had occupations requiring boats, while 74.3% of those who had never attended school had such occupations ($p<0.001$). There was also another difference according to income level; 81.6% of the “non-poor” and 57.6% of “poor” subjects had occupations that required boats ($p<0.01$).

The percentages of households that used river water for cooking, drinking, bathing, and washing were 9.7%, 10.7%, 81.8%, and 91.7%, respectively. The percentages of households that reported they often or always boiled water before drinking, always washed their hands before eating and cooking, and always ensured foods were hot before eating were 42.6%, 26.3%, and 72.7%, respectively. The percentages of households that reported defecating directly into the river and throwing rubbish into the river were 85.4% and 54.0%, respectively.

The mean and standard deviation of the disease prevention proficiency score of the 3607 subjects were 3.6 and 2.4, respectively, and the median value was 3. The percentage distributions of the disease prevention proficiency by three categories—non-functional, minimally functional, and functional—were 3.1%, 64.2%, and 32.7%, respectively.

Table 1 Basic profiles of the subjects living on boats

Characteristics	Male		Female	
	n	%	n	%
Age				
5-9	321	17.5	331	17.4
10-14	348	18.9	337	17.7
15-19	215	11.7	184	9.7
20-24	124	6.8	141	7.4
25-29	133	7.2	137	7.2
30-34	146	7.9	141	7.4
35-39	106	5.8	119	6.3
40-44	97	5.3	123	6.5
45-49	95	5.2	127	6.7
50-54	73	4.0	78	4.1
55-59	36	2.0	41	2.2
Over 60	143	7.8	141	7.4
Total	1837	100.0	1900	100.0
Education level*				
Never gone to school	366	31.3	575	46.7
Primary school	474	40.6	337	27.4
Secondary school and above	187	16.0	111	9.0
Data not available	141	12.1	209	17.0
Total	1168	100.0	1232	100.0
Major occupation*				
Occupations requiring boats				
Sand extraction	447	38.3	398	32.3
Fishing	192	16.4	103	8.4
Transportation	34	2.9	30	2.4
Other occupations requiring boats	39	3.3	67	5.4
Occupations not involving boats				
Vending	7	0.6	185	15.0
Cyclo driver	172	14.7	2	0.2
Labour	43	3.7	48	3.9
Other occupations not involving boats	60	5.1	110	8.9
Inactive	174	14.9	289	23.5
Total	1168	100.0	1232	100.0

*For ages 15 years and over.

Most subjects (95.1%) reported that they would prefer not to continue living on boats, while the remaining subjects expressed no wish to change their living situation.

Disease and injury patterns

The prevalence rates of major diseases and injuries, both currently and over the past 12 months, among the subjects aged 5 years and over were 85.3% for certain infectious and parasitic diseases, 78.0% for diseases of the respiratory system, 51.2% for diseases of the skin and subcutaneous

tissue, 15.4% for diseases of the digestive system, and 13.2% for injuries. Table 2 shows the prevalence rates of major diseases and injuries divided according to sex and age group. Injuries were more prevalent among males than among females ($p < 0.001$). Certain infectious and parasitic diseases, diseases of the respiratory system and diseases of the skin, and subcutaneous tissue were more prevalent among children than among adults ($p < 0.05$, $p < 0.001$, $p < 0.01$, respectively). Diseases of the skin and subcutaneous tissue, diseases of the digestive system and injuries were more

Table 2 Prevalence (%) of major diseases and injuries in the past 12 months and currently among subjects by sex and age group

Major diseases/injuries (ICD-10 code)	Male	Female	Children	Adults
	n = 1837	n = 1900	n = 1337	n = 2400
Certain infectious and parasitic diseases (A.00-B.99)	86.2	84.5	87.2	84.3*
Diarrhoea and gastroenteritis of presumed infectious origin (A.09)	67.6	67.2	70.4	65.7**
Tuberculosis (A.15-A.19)	3.3	2.3	0.2	4.2***
Dengue fever/dengue haemorrhagic fever (A.90-A.91)	7.1	5.7	7.2	6.0
Dermatophytosis (B.35)	24.3	30.0***	11.7	35.8***
Malaria (B.50-B.54)	15.4	6.1***	0.6	16.3***
Ascariasis (B.77)	58.7	60.9	73.3	52.4***
Diseases of the respiratory system (J.00-J.99)	77.9	78.0	82.7	75.3***
Asthma (J.45)	3.8	3.1	4.6	2.8**
Bronchitis and asthma (J.40-J.43, J.45)	6.3	4.7*	8.4	3.8***
Diseases of the skin and subcutaneous tissue (L.00-L.99)	52.5	49.9	54.4	49.4**
Dermatitis and eczema (L.20-L.30)	12.8	10.7*	7.7	14.0***
Acne (L.70)	7.7	5.5**	6.1	6.8
Diseases of the digestive system (K.00-K.93)	16.4	14.5	1.8	23.0***
Injuries	18.0	8.5***	11.7	13.9
Injuries on boats	7.0	5.2*	5.3	6.5
Injuries on land	9.5	3.1***	6.5	6.1
Injuries at work	1.4	0.3***	-	-

* $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$.

Table 3 Adjusted odds ratios (OR) of major diseases and injuries in the past 12 months and currently by socioeconomic conditions among subjects living on boats

Major diseases/injuries (ICD-10 code)	OR (95% CI) in reference to subjects who completed secondary school (n=298), subjects 15 years and over		OR (95% CI) in reference to subjects with occupations not involving boats (n=620), subjects 15 years and over	OR (95% CI) in reference to non-poor† subjects (n=1418), subjects 5 years and over
	Primary school (n=811)	Never went to school (n=941)	Occupation requiring boats (n=1303)	Poor (n=2319)
Certain infectious and parasitic diseases (A.00–B.99)	1.43 (1.00 to 2.03)*	1.31 (0.90 to 1.90)	1.13 (0.87 to 1.47)	1.12 (0.93 to 1.34)
Diarrhoea and gastroenteritis of presumed infectious origin (A.09)	1.27 (0.96 to 1.68)	1.38 (1.03 to 1.87)*	0.99 (0.81 to 1.21)	1.05 (0.91 to 1.21)
Tuberculosis (A.15–A.19)	0.90 (0.40 to 2.04)	1.05 (0.46 to 2.42)	0.46 (0.28 to 0.74)**	1.62 (1.03 to 2.53)*
Dengue fever/dengue haemorrhagic fever (A.90–A.91)	0.69 (0.42 to 1.13)	0.62 (0.36 to 1.07)	1.08 (0.72 to 1.63)	0.71 (0.55 to 0.93)*
Dermatophytosis (B.35)	1.95 (1.42 to 2.69)***	2.44 (1.75 to 3.41)***	1.52 (1.24 to 1.86)***	0.81 (0.69 to 0.93)**
Malaria (B.50–B.54)	2.14 (1.37 to 3.33)***	2.17 (1.36 to 3.45)**	1.31 (0.99 to 1.72)	0.88 (0.70 to 1.10)
Ascariasis (B.77)	1.25 (0.96 to 1.65)	1.49 (1.12 to 1.99)**	1.12 (0.92 to 1.36)	1.01 (0.88 to 1.16)
Diseases of the respiratory system (J.00–J.99)	1.34 (0.99 to 1.82)	1.63 (1.18 to 2.26)**	1.07 (0.85 to 1.34)	0.96 (0.82 to 1.13)
Asthma (J.45)	1.76 (0.71 to 4.38)	1.53 (0.58 to 4.05)	1.12 (0.62 to 2.03)	0.76 (0.53 to 1.09)
Bronchitis and asthma (J.40–J.43, J.45)	1.92 (0.87 to 4.20)	1.51 (0.65 to 3.50)	1.05 (0.63 to 1.73)	0.86 (0.64 to 1.15)
Diseases of the skin and subcutaneous tissue (L.00–L.99)	1.38 (1.05 to 1.81)*	1.74 (1.30 to 2.33)***	1.10 (0.91 to 1.33)	1.18 (1.04 to 1.35)*
Dermatitis and eczema (L.20–L.30)	0.98 (0.64 to 1.52)	1.29 (0.83 to 2.01)	1.42 (1.06 to 1.90)*	0.80 (0.65 to 0.98)*
Acne (L.70)	1.25 (0.74 to 2.10)	1.28 (0.74 to 2.24)	0.74 (0.51 to 1.06)	1.17 (0.89 to 1.53)
Diseases of the digestive system (K.00–K.93)	1.98 (1.32 to 2.97)***	2.24 (1.48 to 3.40)***	0.94 (0.75 to 1.18)	0.91 (0.75 to 1.10)
Injuries	0.58 (0.41 to 0.84)**	0.60 (0.41 to 0.89)*	0.75 (0.56 to 0.99)*	1.11 (0.91 to 1.35)
Injuries on boats	0.99 (0.53 to 1.56)	0.82 (0.46 to 1.47)	1.34 (0.88 to 2.05)	0.81 (0.62 to 1.06)
Injuries on land	0.40 (0.24 to 0.65)***	0.44 (0.26 to 0.75)**	0.44 (0.29 to 0.66)***	1.62 (1.20 to 2.18)**
Injuries at work	1.23 (0.38 to 3.93)	1.62 (0.48 to 5.46)	0.80 (0.35 to 1.81)	0.74 (0.36 to 1.51)

*p<0.05; **p<0.01; ***p<0.001. ORs and 95% confidence intervals (95% CI) were adjusted for sex and age. †Non-poor was defined as subjects whose income per family member was 150,000 Vietnamese dong or more.

prevalent among older adults than among young adults (53.7% v 48.3%, p<0.05; 32.0% v 20.6%, p<0.001; 19.3% v 12.4%, p<0.001, respectively).

Table 3 shows the adjusted odds ratios of major diseases and injuries by education, occupation, and income. Diseases of the respiratory system, diseases of the skin and subcutaneous tissue, and diseases of the digestive system were more prevalent among subjects who had never gone to school than among those who had attended secondary school and above (p<0.01, p<0.001, p<0.001, respectively), while the inverse association was found for injuries (p<0.05). Regarding the relation between injuries and occupations, injuries were more prevalent among subjects with

occupations not requiring boats than among those with occupation requiring boats (p<0.05). Regarding the associations between major diseases and income, the diseases of the skin and subcutaneous tissue were more prevalent among “poor” subjects than among “non-poor” subjects (p<0.05). Diseases of skin and subcutaneous tissue among “overcrowded” households was more prevalent than that among “non-crowded” households (52.3% v 46.0%, p<0.01). Injuries among “overcrowded” households was more prevalent than that among “non-crowded” households, (13.4% v 10.4%, p<0.05).

Table 4 shows the adjusted odds ratios for major diseases and injuries by sanitary practices. Certain infectious and

Table 4 Adjusted odds ratios (OR) for major diseases and injuries in the past 12 months and currently by sanitary practices among subjects aged 5 years and over living on boats

Major diseases/injuries (ICD-10 code)	Certain infectious and parasitic diseases (A.00–B.99)	Diseases of the respiratory system (J.00–J.99)	Diseases of the skin and subcutaneous tissue (L.00–L.99)	Diseases of the digestive system (K.00–K.93)	Injuries
Water treatment before drinking					
Often or always boil (n=1539)	1.00	1.00	1.00	1.00	1.00
Sometimes boil (n=1709)	1.37 (1.14 to 1.65)***	1.07 (0.91 to 1.25)	1.13 (0.98 to 1.29)	1.09 (0.88 to 1.35)	0.91 (0.74 to 1.13)
Never boil (n=479)	4.70 (3.01 to 7.34)***	5.14 (3.47 to 7.62)***	4.49 (3.53 to 5.72)***	2.85 (2.18 to 3.73)***	1.67 (1.27 to 2.21)***
Wash hands before eating					
Always (n=899)	1.00	1.00	1.00	1.00	1.00
Often (n=2216)	1.52 (1.24 to 1.86)***	1.09 (0.91 to 1.31)	1.19 (1.02 to 1.39)*	0.93 (0.73 to 1.18)	0.61 (0.49 to 0.77)***
No (n=569)	3.99 (2.73 to 5.82)***	3.15 (2.30 to 4.30)***	4.01 (3.18 to 5.06)***	2.20 (1.66 to 2.93)***	1.20 (0.91 to 1.59)
Ensure foods are hot before eating					
Always (n=2665)	1.00	1.00	1.00	1.00	1.00
Sometimes (n=538)	5.32 (3.46 to 8.17)***	6.58 (4.55 to 9.50)***	3.13 (2.57 to 3.82)***	0.46 (0.33 to 0.65)***	0.99 (0.74 to 1.32)
No (n=481)	15.92 (7.52 to 33.70)***	13.76 (8.03 to 23.58)***	10.38 (7.82 to 13.77)***	2.94 (2.31 to 3.73)***	1.89 (1.46 to 2.44)***
Sources of water for washing					
Either tap or river (n=656)	1.00	1.00	1.00	1.00	1.00
Always from river (n=3038)	1.66 (1.33 to 2.06)***	1.48 (1.22 to 1.79)***	1.77 (1.49 to 2.10)***	1.22 (0.95 to 1.57)	1.10 (0.85 to 1.43)
Sources of water for bathing					
Either tap or river (n=599)	1.00	1.00	1.00	1.00	1.00
Always river (n=2532)	2.38 (1.91 to 2.97)***	1.26 (1.02 to 1.55)*	1.95 (1.62 to 2.33)***	1.68 (1.26 to 2.23)***	1.01 (0.77 to 1.31)

*p<0.05; ***p<0.001. ORs and 95% confidence intervals (95% CI) were adjusted for sex and age.

Table 5 Prevalence and adjusted odds ratios (OR) of major diseases and injuries in the past 12 months and currently among subjects 5 years and over living on boats according to disease prevention proficiency†

Major diseases/injuries (ICD-10 code)	Prevalence (%) by category of disease prevention proficiency			OR (95% CI)‡	OR (95% CI)§	OR (95% CI)¶
	Non-functional (n = 112)	Minimally functional (n = 2317)	Functional (n = 1178)			
Certain infectious and parasitic diseases (A.00–B.99)	98.2	85.5	83.8	0.76 (0.64 to 0.91)**	0.74 (0.58 to 0.94)*	0.72 (0.56 to 0.94)*
Diarrhoea and gastroenteritis of presumed infectious origin (A.09)	75.9	68.8	63.3	0.77 (0.67 to 0.88)***	0.73 (0.61 to 0.87)***	0.72 (0.59 to 0.87)***
Tuberculosis (A.15–A.19)	1.8	2.5	3.1	1.50 (1.01 to 2.24)*	1.28 (0.78 to 2.09)	1.34 (0.79 to 2.28)
Dengue fever/dengue haemorrhagic fever (A.90–A.91)	1.8	6.9	5.3	0.88 (0.67 to 1.14)	0.79 (0.55 to 1.13)	0.70 (0.47 to 1.04)
Dermatophytosis (B.35)	33.9	30.3	20.4	0.64 (0.56 to 0.75)***	0.69 (0.57 to 0.83)***	0.67 (0.55 to 0.82)***
Malaria (B.50–B.54)	8.9	11.0	10.2	1.04 (0.84 to 1.30)	1.12 (0.88 to 1.42)	1.10 (0.85 to 1.42)
Ascariasis (B.77)	84.8	58.9	60.4	0.87 (0.76 to 0.99)*	0.84 (0.70 to 1.00)*	0.77 (0.63 to 0.93)**
Diseases of the respiratory system (J.00–J.99)	93.8	78.8	74.5	0.70 (0.60 to 0.82)***	0.61 (0.50 to 0.75)***	0.57 (0.46 to 0.71)***
Asthma (J.45)	0.9	3.4	3.4	1.10 (0.78 to 1.56)	1.11 (0.67 to 1.83)	1.09 (0.63 to 1.87)
Bronchitis and asthma (J.40–J.43, J.45)	1.8	5.6	5.1	0.99 (0.75 to 1.31)	1.01 (0.65 to 1.56)	0.97 (0.61 to 1.54)
Diseases of the skin and subcutaneous tissue (L.00–L.99)	72.3	53.0	44.2	0.65 (0.58 to 0.74)***	0.63 (0.53 to 0.75)***	0.59 (0.49 to 0.72)***
Dermatitis and eczema (L.20–L.30)	17.9	13.3	6.8	0.54 (0.44 to 0.67)***	0.53 (0.40 to 0.69)***	0.52 (0.39 to 0.69)***
Acne (L.70)	7.1	5.7	7.7	1.26 (0.98 to 1.63)	1.12 (0.79 to 1.57)	0.97 (0.67 to 1.40)
Diseases of the digestive system (K.00–K.93)	14.3	15.8	15.0	1.06 (0.88 to 1.28)	1.07 (0.87 to 1.33)	1.08 (0.86 to 1.36)
Injuries	8.9	12.4	14.8	1.26 (1.04 to 1.52)*	1.20 (0.92 to 1.55)	1.20 (0.91 to 1.58)
Injuries on boats	5.4	6.3	5.5	0.92 (0.70 to 1.20)	0.92 (0.63 to 1.32)	0.90 (0.60 to 1.33)
Injuries on land	2.7	5.5	8.0	1.55 (1.20 to 2.01)***	1.43 (0.97 to 2.09)	1.45 (0.97 to 2.15)

* $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$. †Individual assessment items included the names of selected infectious and parasitic diseases, measures to prevent infectious and parasitic diseases, measures to prevent malnutrition, sanitary practices, and participation in health education programmes. ‡Odds ratio (OR) for an increment of 1 in score of disease prevention proficiency and 95% confidence interval (95% CI), adjusted by sex and age. §Odds ratio (OR) for an increment of 1 in score of disease prevention proficiency and 95% confidence interval (95% CI), adjusted by sex, age, and occupation requiring/not requiring boats. ¶Odds ratio (OR) for an increment of 1 in score of disease prevention proficiency and 95% confidence interval (95% CI), adjusted by sex, age, occupation requiring/not requiring boats, and education.

parasitic diseases, diseases of the respiratory system, diseases of the skin and subcutaneous tissue, and diseases of the digestive system showed significantly high prevalence rates among subjects who had improper sanitary practices such as never boiled water before drinking, never wash hands before eating, never ensure foods were hot before eating, and use river water for washing or bathing.

Table 5 shows the prevalence of major diseases and injuries divided according to disease prevention proficiency. ORs and p values calculated for three models with different combinations of adjustment variables were generally consistent except the results for injuries on land. Higher disease prevention proficiency was significantly related to lower prevalence of certain infectious and parasitic diseases ($p < 0.05$), diseases of the respiratory system ($p < 0.001$), and diseases of the skin and subcutaneous tissue ($p < 0.001$), after adjustment by sex, age, occupation, and education.

People who preferred not to continue living on boats were more likely to have higher prevalence rates of certain infectious and parasitic diseases ($p < 0.01$), diseases of the respiratory system ($p < 0.001$), and diseases of the skin and

subcutaneous tissue ($p < 0.01$) than those who expressed no wish to change their living situation.

DISCUSSION

The results of this quantitative study showed the prevalence of certain infectious and parasitic diseases, diseases of the respiratory system, diseases of the skin and subcutaneous tissue, diseases of the digestive system, and injuries among people living on boats in Hue City. The prevalence rates of the diseases and injuries were significantly associated with sanitary practices, disease prevention proficiency, and subjects' preference regarding continued living on boats. Disease prevention proficiency was shown to be conducive to reducing the prevalence rates of certain infectious and parasitic diseases, diseases of the respiratory system, and diseases of the skin and subcutaneous tissue regardless of the subjects' sex, age, or socioeconomic status.

In Vietnam, efforts have been made to provide people with access to primary health care services within five kilometres from their home.¹⁰ However, families living under fragile

What this paper adds

People living on boats are underprivileged with regard both to socioeconomic and living conditions. However, there have been no previous quantitative studies of disease patterns in such populations. This comprehensive field study performed in Hue City, Vietnam, illustrated the major diseases and injuries occurring among people living on boats and showed a web-like relation between health status, socioeconomic conditions, sanitary practices, disease prevention proficiency, and preference to continued living on boats.

Policy implications

- Our findings regarding variations in health status among people living on boats provide a basis for the development of measures to improve the health of populations living under specific fragile conditions.
- Resettlement of people living on boats onto the land and programmes to improve disease prevention proficiency by taking account of the peoples' experience, in addition to secure means of income, will benefit health development in this population under fragile conditions.

conditions have been reported to have fewer visits to health care facilities than those higher on the socioeconomic scale.¹¹ Therefore, disease statistics reported from health care facilities generally do not accurately reflect health situations of people living under fragile conditions. In this study, a response rate of 83.1% was achieved from a population living under fragile conditions. In the study area, along the Huong River, there is a long history of households living on boats. Almost all households living on boats in the area are on the household registration records. Therefore, this study was regarded as providing comprehensive quantitative data of the prevalence rates of disease and injury among people aged 5 years and over living on boats in Hue City. The report of the health status of children less than 5 years old in the study area, available from Hue-Bretagne boat clinic, will complement information regarding disease and injury patterns in the population as a whole. The participation of local medical doctors visiting boat households for onsite health examinations encouraged families to participate in the survey. The participation and cooperation of the heads of quarters in the preparation and implementation of the survey facilitated people's understanding of the survey activities. This process also contributed to improvement of security during the survey in the study area. Although most health problems were well diagnosed by local medical doctors, diagnosis of mental health problems, and diseases related to reproduction were regarded as unsuitable for a general community health survey; therefore, these diagnoses were excluded from this study. Future studies focusing specifically on these problems will widen our understanding of the disease and injury patterns of this population.

The prevalence rates of selected infectious and parasitic diseases were comparatively high in the study population. The prevalence of tuberculosis per 100 000 people was 580 per year, which was fivefold greater than the national estimate of 119¹² and fourfold greater than the areal estimate of 140.¹³ The proportion of poor people among the study population, 52.6%, was considerably higher than the national average, 17.2%.⁸ Poverty was considered to be among the factors responsible for the high prevalence of tuberculosis in this population.

The prevalence of ascariasis was 1.4-fold higher than the national estimate, 44.4%.¹⁴ Low communal services for people living in river villages are related to the high prevalence of ascariasis.¹⁵ The number of reported cases of malaria (10.3% per year) was higher than those in seven other regions in Vietnam (range, 0.3%–4.7% per year reported in 1998),¹⁶ although the incidence and prevalence of malaria have been reduced significantly in Vietnam because of malaria control efforts.¹⁷ The prevalence rate of dengue fever/dengue haemorrhagic fever (6.4% per year) was 10-fold higher than that reported in another region of Vietnam (0.44% per year).¹⁸ These differences were probably related to the increased frequency of exposure to mosquito breeding sites of those living on the Huong River and its branches.¹⁹

The prevalence rates of diseases of the respiratory system and diseases of the skin and subcutaneous tissue were comparatively high (sixfold and fivefold greater, respectively) as compared with those in the communities on land along the river in Hue.²⁰

The relations among education, occupation requiring/not requiring boats, income, and risks of disease and injury are complex in people living under fragile conditions. Statistical adjustment by sex and age group is useful in the analysis of these relations among people living on boats, because of the significant differences in the prevalence of diseases and injuries by sex and age group. The subjects in this study showed reduced rates of participation in school education, indicating that they had low education levels. Lack of

opportunities for school education have also been reported in some countries belonging to the Association of Southeast Asia Nations (ASEAN) where adult illiteracy rates were reported to range from 2.8% to 35.9% for men and 4.8% to 66.8% for women in 2000.²¹ Lower educational attainment was related to a higher risk of major diseases but lower risk of injury. The first relation has been seen in other low income countries.^{22–23} The second relation—the association between lower educational level and lower risk of injury—was because people with lower educational level were more likely to work on boats where the risk of injuries was considered to be comparatively small. In contrast, work on land such as cyclo driver, labour, and vending, is regarded as a risk of injuries in crowded urban spaces.

The prevalence rates of dermatophytosis, and dermatitis and eczema were high among non-poor subjects, most of whom had occupations involving boats. Occupations requiring boats, such as sand extraction, fishing, and transportation, brought higher incomes than occupations that do not involve boats. At the same time, occupations requiring boats reflect working environments, on the Huong River and its branches, exposure to river water contaminated with faeces and waste.¹⁰ These working environments are regarded as risks to the particular infectious and skin diseases mentioned above.

In contrast, tuberculosis was more prevalent among poor subjects than non-poor subjects. Poverty among tuberculosis patients may be explained as a background risk factor but also as a consequence of this long term illness that restricts patients' income generating activities and thus results in isolation from social activities.^{24–26} Most poor subjects were involved in occupation not requiring boats, such as cyclo driver, labour, and vending. These occupations were regarded as a risk of tuberculosis because they are associated with contacts with unspecified large number of people in urban crowded spaces.

People living on boats along the Huong River struggled with their means of income, working environments, and risk of diseases. To earn slightly better incomes, people chose occupations involving boats where the risks of particular diseases are high because of the river environment. People with slightly higher educational levels chose occupations on land and had higher risks of injury and tuberculosis but comparatively lower risks of some other infectious diseases.

People living on boats used the Huong River for defecation and rubbish disposal as well as a source of water for drinking, cooking, washing, and bathing thus leading to high risks of skin diseases and diseases transmitted orofecally. Associations between personal hygiene and both ascariasis and diarrhoeal disease have been reported previously.^{22–27–28} Sanitary practices were also critically related to other major diseases, as reported in other low income communities.^{26–29}

Functional disease prevention proficiency lowered the prevalence of a wide range of diseases, and this association was found even after excluding the influence of occupations requiring/not requiring boats and subjects' educational level. Improvements in health literacy in low income countries can be achieved by community based programmes to meet the needs of the communities.³⁰ Almost all communities in Vietnam have their own health stations. They deliver public health information, expanding health education and curative care.¹⁰ These health stations could be a focal point in programmes to improve disease prevention proficiency. The recent development of programmes in Hue City to increase disease prevention proficiency will contribute to decrease the risk of disease.

Fragile living conditions on boats may encourage more people to consider adopting a life on land, despite the uncertainty regarding the security of the means of income of

occupations requiring boats or those on land. Relocation from living on boats to settlement on land would improve sanitary conditions. Based on census data, we estimated that 2% of urban population in Vietnam live under fragile conditions along rivers or canals.^{4 31} Improvement of disease prevention proficiency of the urban boat population through appropriate interventions taking account of the person's experience must be considered in addition to a secure means of income when they begin living on land.

In conclusion, this large scale comprehensive field study in Hue City, Vietnam, quantified major diseases and injuries among people living on boats to provide solid evidence on which to base policy making decisions. Variations in health status in this population showed a complex relation with education, occupations requiring/not requiring boats, income, sanitary practices, disease prevention proficiency, and people's preference to continued living on boats. Programmes to improve disease prevention proficiency that take into account both way of life and occupation would reduce the risk of disease and injury in this fragile urban population.

ACKNOWLEDGEMENTS

We acknowledge members of the Hue People's Committee, Mr Nguyen Viet Tien, former mayor and Mr Nguyen Van Cao, present mayor; Dr Nguyen Nhen, Director of the Bureau of Foreign Affairs of Hue City; Dr Tran Bui, Director of the Hue Healthcare Center for supporting this study. We thank the local leaders of the six quarters, the medical doctors of the Hue Healthcare Center, the chiefs of the six health stations, and all of the government staff of the Bureau of Foreign Affairs of Hue City for their contribution to implementation of these field works.

Authors' affiliations

N K L Quang, T Takano, M Watanabe, T Inose, Y Fukuda, Health Promotion, Department of International Health Development, Graduate School of Tokyo Medical and Dental University, Tokyo, Japan
K Nakamura, K Seino, International Health and Medicine, Department of International Health Development, Graduate School of Tokyo Medical and Dental University

Funding: this study was partly supported by Grants-in-Aid for Scientific Study by Japan Society for Promotion of Science.

Conflicts of interests: none.

REFERENCES

- 1 **United Nations**. *World urbanization prospects, the 2003 revision*. New York: United Nations, 2003.
- 2 **Nhien N**. Community participation and urban policies on health and environment of the city of Hue: Vietnam. In: Takano T, ed. *Healthy cities and urban policy research*. London: Spon Press, 2003:261–4.
- 3 **Bui T**. Report on Healthy City Programme of Hue City. Workshop on healthy cities: preparing for the 21st century, WPR/EUD/EHE (O) (2)/99/INF, /14, Manila, Western Pacific Region, 1999:1–4.
- 4 **Hue Government**. *The project of residing people living on boat in Hue city*. Hue City: Hue Government, 2002.
- 5 **World Health Organisation**. Ottawa Charter for Health Promotion. *Health Promot* 1986;1:iii–v.
- 6 **Takano T, Nakamura K**. An analysis of health levels and various indicators of urban environments for Healthy Cities projects. *J Epidemiol Community Health* 2001;55:263–70.
- 7 **World Health Organisation**. Quantifying selected major risks to health. In: *World health report 2002*. Geneva: WHO, 2002:47–97.
- 8 **Vietnam**. *The comprehensive poverty reduction and growth strategy*, No. 2685/VP-CP-QHQT. Hanoi: Vietnam, 2002.
- 9 **World Health Organisation**. *International statistical classification of diseases and related health problems, tenth version*. Geneva: WHO, 1992.
- 10 **Committee for Population, Family and Children Vietnam**. *Vietnam demographic and health survey 2002*. Hanoi: General Statistical Office of Vietnam, 2003.
- 11 **Falkingham J**. Poverty, out-of-pocket payments and access to health care: evidence from Tajikistan. *Soc Sci Med* 2004;58:247–58.
- 12 **World Health Organisation**. *Vietnam country profile. Global tuberculosis control: surveillance, planning, financing*. Geneva: WHO, 2004:120–2.
- 13 **Hue Health Care Center**. *Hue health report 2004*. Hue City: Hue Health Care Center, 2004.
- 14 **van der Hoek W, De NV, Konradsen F, et al**. Current status of soil-transmitted helminthes in Vietnam. *Southeast Asian J Trop Med Public Health* 2003;34:1–11.
- 15 **Curtale F, Pezzotti P, Saad YS, et al**. An analysis of individual, household, and environmental risk factors for intestinal helminth infection among children in Qena Governorate, upper Egypt. *J Trop Pediatr* 1999;45:14–17.
- 16 **Laxminarayan R**. Dose reducing malaria improve household living standards? *Trop Med Int Health* 2004;9:267–72.
- 17 **Nam NV, de Vries PJ, Toi LV, et al**. Malaria control in Vietnam: the Binh Thuan experience. *Trop Med Int Health* 2005;10:357–65.
- 18 **Do QH, Nguyen TKT, Vu TQH, et al**. Dengue epidemic in Southern Vietnam, 1998. *Emerg Infect Dis* 2000;6:422–5.
- 19 **Birley MH**. *Guidelines for the forecasting of vector-borne disease implications of water resources development*. WHO/CWS/91.3. Geneva: WHO, 1991.
- 20 **Hau TD**. *ACWP mobile healthcare program, 2003 report*. Hue City: Hue Medical College, 2003.
- 21 **Lee ISM**. Baseline data in ASEAN member countries. In: Takano T, ed. *Healthy cities and urban policy research*. London: Spon Press, 2003:190–221.
- 22 **Holland CV, Taren DL, Crompton DWT, et al**. Intestinal helminthiasis in relation to the socioeconomic environment of Panamanian children. *Soc Sci Med* 1988;26:209–13.
- 23 **Traub RJ, Robertson ID, Irwin P, et al**. The prevalence, intensities and risk factors associated with geohelminth infection in tea-growing communities of Assam, India. *Trop Med Int Health* 2004;9:688–701.
- 24 **World Resources Institute**. *Changing environments, changing health. In: World resources 1998–1999: a guide to the global environment: environment change and human health*. Washington, DC: World Resources Institute, 1998:36–72.
- 25 **Long NH, Johansson E, Diwan VK, et al**. Fear and social isolation as consequences of tuberculosis in Vietnam: a gender analysis. *Health Policy* 2001;58:69–81.
- 26 **Gustafson P, Gomes VF, Vieira CS, et al**. Tuberculosis in Bissau: incidence and risk factors in an urban community in sub-Saharan Africa. *Int J Epidemiol* 2004;33:163–72.
- 27 **Atukorala TMS, Lanerolle P**. Soil transmitted helminthic infection and its effect on nutritional status of adolescent schoolgirls of low socioeconomic status in Sri Lanka. *J Trop Pediatr* 1999;45:18–22.
- 28 **Tumwine JK, Thompson J, Katua-Katua M, et al**. Diarrhea and effects of different water sources, sanitation and hygiene behaviour in East Africa. *Trop Med Int Health* 2002;7:750–6.
- 29 **Dogra S, Kumar B**. Epidemiology of skin diseases in school children: a study from northern India. *Pediatr Dermatol* 2003;20:470–3.
- 30 **Ohnishi M, Nakamura K, Takano T**. Improvement in maternal health literacy among pregnant women who did not complete compulsory education: policy implications for community care services. *Health Policy* 2004;72:157–64.
- 31 **Bolay JC, Cartoux S, Cunha A, et al**. Sustainable development and urban growth: precarious habitat and water management in Ho Chi Minh City, Vietnam. *Habitat Int* 1997;21:185–97.