

**P29** **OUT OF CONTROL? EXPERIENCES OF TRANSITIONS BETWEEN CARE SETTINGS AT THE END OF LIFE FOR OLDER ADULTS WITH HEART FAILURE: A QUALITATIVE STUDY**

doi:10.1136/jech.2010.120477.29

<sup>1</sup>L Holmes, <sup>2</sup>J Addington-Hall, <sup>3</sup>G Grande, <sup>4</sup>S Payne, <sup>5</sup>J Seymour, <sup>1</sup>B Hanratty. <sup>1</sup>Division of Public Health, University of Liverpool, Liverpool, UK; <sup>2</sup>School of Health Sciences, University of Southampton, Southampton, UK; <sup>3</sup>School of Nursing, Midwifery & Social Work, University of Manchester, Manchester, UK; <sup>4</sup>Division of Health Research, Lancaster University, Lancaster, UK; <sup>5</sup>School of Nursing, Midwifery & Physiotherapy, University of Nottingham, Nottingham, UK

**Background** Transitions between care settings have been associated with poor continuity and quality of care for older people. Movements between places of care occur frequently in the months before death, making them an important influence on well-being and health status. People with heart failure may be particularly likely to experience frequent, unplanned admissions to hospital.

**Objective** To explore and understand experiences of care transitions for older adults with heart failure at the end-of-life.

**Design** In-depth qualitative interviews with 35 people with heart failure, lung cancer or stroke. Framework methodology was used for data analysis.

**Participants** Fifteen adults aged over 75 years with heart failure, whose hospital clinician answered "no" to the question, would you be surprised if this patient dies within the next 12 months? Half the participants were from socially disadvantaged backgrounds.

**Findings** Participants described how they struggled to have control over many of their end of life experiences. Their ability to negotiate their role with professionals, family and the wider world was threatened by their biographical experiences and physical helplessness. Long-standing relationships with family doctors were associated with low expectations and a perceived loss of control, particularly for the most disadvantaged people. A majority had lost all trust in their GPs, and the doctor was allowed to share in symptom management only when severe—preserving some autonomy for the older person, but threatening physical health and future choices. The rest had taken on the role of passive recipients of advice and treatment. Frequent admissions to hospital were tainted by the perception of being a burden to staff reluctant to help with basic care, and families obliged to visit.

**Conclusions** Notions of choice and empowerment prevalent in current health policy discourses appeared to have little relevance to the lived experiences of the older adults in this study. Without an understanding of biographical and social context, analyses of how older people experience health and social care at the end-of-life risk promoting concepts of individual control that many older adults do not enjoy. If end-of-life care policies are to produce equitable outcomes, they must go beyond simple solutions to address the complexity of relationships older adults have developed with health services over their lifetimes.

**P30** **VALIDITY AND RELIABILITY OF TWO INDICATORS OF SOCIOECONOMIC POSITION IN A POPULATION SURVEY OF OLDER ADULTS**

doi:10.1136/jech.2010.120477.30

R J Lacey, J Belcher, P R Croft. *Arthritis Research Campaign National Primary Care Centre, Keele University, Keele, Staffordshire, UK*

**Objective** There is increasing interest in measuring the influence of socioeconomic position (SEP) across the life-course. However, there are few cohorts that encompass the whole life-course, so many studies use data from cross-sectional studies, hence the need for ensuring the accuracy of recall of the SEP measures used. The aim of

this study was to investigate the reliability and validity of two measures of SEP: education and longest job.

**Design** Prospective cohort study.

**Setting** Large population survey of older adults from the general population in North Staffordshire, UK.

**Main Outcome Measures** Health Questionnaire (HQ) including a four-part question on education and a five-part question about the longest job.

**Participant** In two pilot studies, random samples of patients aged 50+ registered with different general practices were mailed a HQ (nested test–retest repeatability studies were included; n=500; n=160). In the main study, all patients aged 50+ registered with three different general practices were sent a HQ at baseline. 6 years later, 3410 responders consenting to further contact were mailed a follow-up HQ; a subsample of these participants took part in independent qualitative interviews, which included a reconstruction of critical landmarks in their life-course. Ethical approval was granted for all stages of the study.

**Analysis** Validity was assessed by: percentage completion; internal percentage agreement within each set of questions; percentage agreement of qualitative and quantitative data for age left school and longest job questions; comparing age left school data with historical change in legal school leaving age. Reliability of both questions was calculated from the pilot repeatability studies using ICC and  $\kappa$ , with 95% CI.

**Results** The adjusted response to all stages of the study was good (62–84%). In the main surveys, completion of the age left school and longest job questions was excellent (94–98%). Internal agreement within each set of questions was good (education: 80–97%; longest job: 95–100%). Comparison of survey and interview data showed good agreement for age left school (85%) and longest job (95%). The change in age left school data concurred exactly with the change in legal school leaving age implemented in 1947. Reliability from pilot test–retests was excellent for recalled age left school (ICC=0.95; 95% CI 0.93 to 0.96) and substantial for longest job ( $\kappa$ =0.69; 95% CI 0.55 to 0.83).

**Conclusion** The age left school and longest job questions in this study are valid and reliable measures.

**Health behaviours/risk factors (obesity, smoking, physical activity, food)**

**P31** **SOCIAL CAPITAL AND HEALTH BEHAVIOUR**

doi:10.1136/jech.2010.120477.31

<sup>1</sup>T Nieminen, <sup>2</sup>T Martelin, <sup>1</sup>S Koskinen, <sup>1</sup>R Prättälä, <sup>3</sup>E Alanen, <sup>3</sup>M T Hyypä. <sup>1</sup>Department of Health, Functional Capacity and Welfare, National Institute for Health and Welfare (THL), Helsinki, Finland; <sup>2</sup>Thematic Programmes Unit, National Institute for Health and Welfare (THL), Helsinki, Finland; <sup>3</sup>Department of Chronic Disease Prevention, National Institute for Health and Welfare (THL), Turku, Finland

**Objective** To examine how different dimensions of social capital and health behaviour are associated.

**Design** Cross-sectional data of the Health 2000 Health Examination Survey including a personal interview and self-administered questionnaires.

**Setting** Representative sample (N=8028) of the Finnish adult population.

**Participants** 6986 (87%) adults aged 30 years or over.

**Main Outcome Measures** Non-smoking, non-excessive drinking, leisure-time physical activity, daily use of vegetables, adequate duration of sleep.

**Results** Social capital was found to be associated with health behaviour. The dimension of social participation and networks was positively associated with every type of health behaviour. High