



OPEN ACCESS

# Parallel contagion phenomenon of concordant mental disorders among married couples: a nationwide cohort study

Jong-Yi Wang,<sup>1</sup> Ming-Hong Hsieh,<sup>2,3</sup> Pei-Ching Lin,<sup>4</sup> Chiu-Shong Liu,<sup>5,6</sup> Jen-De Chen<sup>7</sup>

<sup>1</sup>Department of Health Services Administration, China Medical University, Taichung, Taiwan

<sup>2</sup>Department of Psychiatry, Chung Shan Medical University Hospital, Taichung, Taiwan

<sup>3</sup>School of Medicine, Chung Shan Medical University, Taichung, Taiwan

<sup>4</sup>Tele-Healthcare Center, Kaohsiung Medical University Chung-Ho Memorial Hospital, Kaohsiung, Taiwan

<sup>5</sup>Department of Family Medicine, China Medical University Hospital, Taichung, Taiwan

<sup>6</sup>Department of Medicine, China Medical University, Taichung, Taiwan

<sup>7</sup>Department of Sports, National Changhua University of Education, Changhua, Taiwan

## Correspondence to

Dr Jong-Yi Wang, Department of Health Services Administration, China Medical University, No.91, Hsueh-Shih Road, Taichung 40402, Taiwan; [ericwang@mail.cmu.edu.tw](mailto:ericwang@mail.cmu.edu.tw)

Received 25 August 2016

Revised 12 December 2016

Accepted 5 February 2017

Published Online First

20 February 2017

## ABSTRACT

**Background** The aggregation of mental disorders in couples, as reported by prior research, indicates the effect of familial environments and warrants attention. However, the concordant categories of mental disorders in couples remain unclear. This study investigated spousal concordance for the category of mental disorders among couples throughout Taiwan by using factors associated with such disorders.

**Methods** 5643 couples in the 2002–2013 Taiwan National Health Insurance Research Database were analysed and compared with propensity-matched 5643 non-couples. Twelve independent variables, including spouse and shared characteristics, and the category of mental disorders were analysed, mainly by using multinomial logistic regression.

**Results** The determined prevalence rates for concordant categories of mental disorder were 0.19% for affective disorders, 6.96% for anxiety disorders, and 3.15% of other mental disorders. Multinomial logistic regression results revealed that two spouses were significantly more likely to be diagnosed with the same category of the aforementioned mental disorders (ORs=2.914, 1.776 and 1.727, respectively). Other associated factors included gender, age, occupation, comorbidity and region. The odds of concordances were significantly higher in couples than in non-couples.

**Conclusions** A category of mental disorder in one spouse is a determinant of that in the other spouse. This study extended the emotional contagion theory to the phenomenon of parallel contagion to reflect the three concordances, suggesting a direction of family-based mental health intervention, particularly prevention for the same category of mental disorders in couples. Policymakers should strengthen the coping strategies of the caring spouse and external support system to psychiatrically vulnerable families.

## INTRODUCTION

Mental illness is a growing problem worldwide.<sup>1</sup> Genetic relatedness has been reported to be a determinant of mental disorders among family members.<sup>2–3</sup> Nevertheless, recent studies have reported the aggregation of mental disorders in couples who did not share genetic factors. A study revealed that depression is likely (OR=2.08) to occur in both married partners if either of them experienced depression.<sup>4</sup> A longitudinal study further indicated that somatic and psychiatric morbidity of one partner could affect the psychiatric condition of the other married partner.<sup>5</sup> Based on the existing literature, concordance of specific

mental disorders in couples includes depression,<sup>4–6</sup> anxiety<sup>7</sup> and substance abuse.<sup>8–10</sup> However, inconclusive and incomplete findings still exist among related research. Furthermore, existing concordance studies either aimed at a certain disorder or examined a relatively small number of associated factors. Hence, determining concordance for the category of mental disorders among couples out of a complete spectrum of mental disorders by using comprehensive factors is highly demanding.

Previous studies have explained the phenomenon of spousal concordance of diseases by using ‘assortative mating’ and ‘cohabitation effects’.<sup>11–13</sup> Assortative mating refers to the tendency of mate selection depending on the similarity of individual characteristics, including values and personality traits. Cohabitation effects explicate the significance of common household environmental influences, including the sharing of numerous aspects of lifestyle, after marriage. In addition, because marital relationship is reciprocal and interdependent, concordance of mental disorders in couples can be further explained by ‘emotional contagion’.<sup>14</sup> Marriage is an imperative resource during stress.<sup>15</sup> Accordingly, the emotional state of a spouse bidirectionally influences that of the partner.<sup>16</sup> Mutual experience of stress can delineate the resembling depression symptomatology between both spouses. In particular, higher levels of depression in husbands could predict the subsequent exacerbation of depressive symptoms in wives.<sup>17</sup> Furthermore, emotional similarity induced by the sharing of stressors and emotional support could unravel the phenomenon of mental illness transmission from one spouse to other spouse.<sup>18</sup> The theory of emotional contagion was primarily applied to depict the concordance of depression in couples.<sup>16–19</sup> However, it is conceivable that the initial similarity between both spouses in individual characteristics from the time of mate selection can expand to subsequently gradual conformity in health behaviours.<sup>20–22</sup> Affect and behaviour similarities could potentially lead to even an exposure to various types of disorders, other than depression, under the effects of durable cohabitation in the common ecological system, which involves family and work environment as the microsystem and mesosystem, respectively.<sup>23</sup> Therefore, the current research extended the concordance of depression to a hypothetical concordance of other categories of mental disorders, including anxiety and others, on the basis of assortative mating, cohabitation effects and emotional contagion jointly.



CrossMark

**To cite:** Wang J-Y, Hsieh M-H, Lin P-C, et al. *J Epidemiol Community Health* 2017;**71**:640–647.

For examining the effect of one mentally ill spouse on the other spouse, previous studies have adopted gender and age as two major related factors.<sup>4 5 14 24</sup> A higher risk of mental disorders has been typically associated with women and old ages.<sup>4 5</sup> Low household income has been closely tied to high psychiatric morbidity.<sup>25</sup> In addition, research has indicated that the presence of physical diseases such as diabetes and cancer was associated with the risk of mental disorders.<sup>26 27</sup> Therefore, comorbidity in each spouse was considered a factor associated with the occurrence of mental disorders. Moreover, a marked discrepancy in prevalence rates of mental disorders, including depression, anxiety and substance abuse, was present across various occupations.<sup>28–31</sup> Environmental characteristics might be associated with the occurrence of mental diseases.<sup>8 23</sup> However, the findings remained inconsistent. The association between the characteristics of geography where the family was located and the occurrence of mental disorders substantiates the need for investigation.

Scarce data have reported the concordant categories of mental disorders; therefore, this study investigated spousal concordance of the category of physician-diagnosed mental disorders by using all potential associated factors of mental disorders addressed in previous studies. Hence, the concordance of mental disorders in couples could be fully determined.

## METHODS

### Hypothesis and research design

This study hypothesised that, after controlling for the effects of related factors including individual and shared characteristics, the category of mental disorders diagnosed in one spouse is associated with that diagnosed in the other spouse. This hypothesis of categorical concordance was tested in a longitudinal population-based case–control design, analysing nationwide data of married couples in Taiwan. The study was approved by the institutional review board at China Medical University Hospital, Taiwan.

### Data source and study sample

The National Health Insurance (NHI) programme, launched in 1995, provides comprehensive healthcare to more than 99.7% of the residents in Taiwan (N=23.50 million). This retrospective study employed the 2002–2013 National Health Insurance Research Database (NHIRD), which includes data for one million randomly sampled beneficiaries and thus entirely represents all enrollees in Taiwan. The NHIRD is maintained by Taiwan's National Health Research Institutes and records all original medical claims under the universal NHI programme.

The couple could be identified from the NHIRD only under the condition of one spouse enrolling being a health insurance insured with a certain occupation and the other spouse being a dependent of this insured spouse. In addition, to ascertain the initial diagnosis of mental disorders, patients diagnosed with mental disorders (International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM): 290.x–319.x) in 2002 were excluded from the study. This study sought to retrieve all the identifiable couples. Accordingly, a total of 5643 married couples (5643 insured and dependent spouses) were identified for further analysis. To increase the strength of inference, the case group (couples) was matched by gender, age and comorbidity with the control group (non-couples) through 1:1 propensity score matching (PSM). This procedure was performed twice because there are two members in one couple (dual PSM). The results indicated that the couples and the non-couples were not significantly different in the three propensity

score-matched variables (all  $p=1$ ), indicating that the two groups qualified for comparison. Consequently, 5643 couples and 5643 non-couples were eligible for subsequent analysis.

### Variables

The category of mental disorders among dependent spouses served as the outcome variable that was categorically defined in this study. The category of mental disorders among insured and dependent spouses, with the former functioning as a major independent variable, both identically included affective disorders, anxiety disorders and other mental disorders, as addressed by previous studies. On the theoretical basis of emotional contagion and affect similarity, all affective disorders were applied, instead of merely depression. The categorisation of affective disorders involved any of the following ICD-9-CM codes: 296.0x, 296.2x, 296.3x, 296.4x, 296.5x, 296.6x, 296.7, 296.8x, 296.9x, 300.4, 301.13, 309.0, 309.1 and 311. The coding for anxiety disorders in this study included 300.0x, 300.2x, 300.3, 308.x and 309.81. The remaining 290.x–319.x diagnoses other than affective disorders and anxiety disorders were classified as other mental disorders. Mental disorders of low incidence, such as substance abuse, were merged into the category of 'other' to avoid unstable statistical estimations engendered by low frequency. Spouses were possibly not diagnosed with any mental disease throughout the observation period. Therefore, no mental disorders were added as one status category in the variables. Concordance in this study was defined as a clinical state in which insured and dependent spouses were diagnosed with any of the same category of mental disorders, including affective, anxiety and other mental disorders; otherwise, discordance or no concordance would be reported.

As the aforementioned studies indicated reciprocity of marital relationship in one common environment, the present research grouped the following independent variables that might be associated with the occurrence of mental disorder into three categories. (1) Insured spouse characteristics including category of mental disorders; (2) dependent spouse characteristics; (3) shared ecological characteristics. Age did not pass a normality test and thus was transformed into five ordinal levels according to the frequency distribution. Occupation, premium-based monthly salary, catastrophic illness and injury, and region were defined from the official classifications of the NHI programme. Since, Taiwan allows only opposite-sex marriage, only the gender of the dependent was adopted to eliminate predictable collinearity. Comorbidity was calculated using the Charlson Comorbidity Index (CCI),<sup>32</sup> a frequently used measure in clinical research. After original scoring from 0 to 6 by weighting ICD-9-CM codes for each patient, this study classified comorbidity into 0 (without comorbidity) and 1 (with comorbidity) because of the low-frequency distribution in CCI scores higher than 1. Urbanisation level was assessed using a 5-level scale, with levels 1 and 5 indicating the highest and lowest urbanisation levels, respectively. All the 13 independent variables were measured on a categorical or ordinal level.

### Data analysis

Statistical methods included the  $\chi^2$  test and multinomial logistic regression. The  $\chi^2$  test examined bivariate associations, with the observation values of prevalence reported. Since the outcome variable comprised four categories, namely affective disorders, anxiety disorders, other mental disorders and no mental disorders, multinomial logistic regression was used, instead of the commonly used binomial logistic regression that analyses a

**Table 1** Characteristics of couples, 2002–2013 (N=5643 couples)

Variables	Frequency	Per cent
<i>Insured spouse characteristics</i>		
Category of mental disorders		
No mental disorders	3404	60.32
Affective disorders	151	2.68
Anxiety disorders	1197	21.21
Other	891	15.79
Gender		
Female	1460	25.87
Male	4183	74.13
Age (years)		
16–34	379	6.72
35–44	1295	22.95
45–54	1607	28.48
55–64	1185	21.00
≥65	1177	20.86
Occupation		
First category (private employee and government employee)	2884	51.11
Second category (labour union member)	785	13.91
Third category (farmer and fisherman)	695	12.32
Fourth, fifth and sixth categories (soldier, social security insured, and veteran and religious group member)	1279	22.67
Premium-based monthly salary (US\$)		
≤576	1796	31.83
576–760	1833	32.48
760–960	413	7.32
960–1210	481	8.52
1210–1526	473	8.38
>1526	647	11.47
Comorbidity (CCI)		
0	3853	68.47
≥1	1774	31.53
Catastrophic illness and injury		
Absent	5106	90.48
Present	537	9.52
<i>Dependent spouse characteristics</i>		
Category of mental disorders		
No mental disorders	3230	57.24
Affective disorders	189	3.35
Anxiety disorders	1410	24.99
Other	814	14.42
Gender		
Female	4183	74.13
Male	1460	25.87
Age (years)		
16–34	654	11.59
35–44	1435	25.43
45–54	1391	24.65
55–64	1152	20.41
≥65	1011	17.92
Comorbidity (CCI)		
0	4102	72.91
≥1	1524	27.09
Catastrophic illness and injury		
Absent	5197	92.10
Present	446	7.90
<i>Shared ecological characteristics</i>		
Region		
Taipei	1980	35.17

Continued

**Table 1** Continued

Variables	Frequency	Per cent
Northern	869	15.44
Central	1076	19.12
Southern	752	13.36
Southeast	847	15.05
Eastern	105	1.87
Urbanisation level		
Level 1 (highest)	1745	31.04
Level 2	1665	29.62
Level 3	954	16.97
Level 4	739	13.14
Level 5 (lowest)	519	9.23

\*p&lt;0.05.

CCI, Charlson Comorbidity Index.

dichotomous outcome variable. Multinomial logistic regression was used for multivariate analysis, with all other covariates held constant. Since the case group was matched to compare with the control group through dual PSM, no further variables besides couple status were used in the couple-level analysis. Moreover, this study conducted collinearity diagnostics using indices including variance inflation and tolerance to detect any potentially high interrelation of characteristics between two spouses of one household. The unit of analysis in this study was primarily one individual. However, the one pair of couple or non-couple was used as the unit of analysis in the case-control analysis for concordant mental disorders. All tests were two sided and performed on an  $\alpha$  value of 0.05. Data were analysed using SAS V.9.4, SAS Institute, Cary, North Carolina, USA.

## RESULTS

**Table 1** presents the descriptive statistics of the study couples. The majority of the insured spouses were without any mental disorders (60.32%). Mental disorders of the insured spouses, ranked by the prevalence rate, included anxiety, other and affective disorders (21.21%, 15.79% and 2.68%, respectively). Most of the dependent spouses did not receive a mental disorder diagnosis (57.24%). The prevalence rates of anxiety, other and affective disorders among the dependents were 24.99%, 14.42% and 3.35%, respectively.

The  $\chi^2$  test revealed that, among the 13 independent variables, 11 were significantly associated with the category of mental disorders among the dependent spouses, including the category of mental disorders among the insured spouses (all  $p<0.01$ ) (**table 2**). Detailed results of the two major variables showed that no mental disorders of the insured spouses were significantly associated with no mental disorders of the dependent spouses (62.22%); affective disorders in the insured spouses were significantly associated with affective disorders in the dependent spouses (7.28%); anxiety disorders in the insured spouses were significantly associated with anxiety disorders in dependent spouses (32.83%); and other mental disorders of the insured spouses were significantly associated with other mental disorders of the dependent spouses (19.98%). The highlighted areas in **table 2** indicate this tendency of categorical concordance. In addition, the percentage of insured and dependent spouses, both not diagnosed with any mental disorders was 37.53% (aggregation of no mental disorders, data not shown in the table). Considering any psychiatric disorders, the prevalence rate of the aggregation of mental disorders in the couples

**Table 2** Category of mental disorders among dependent spouses, according to the characteristics of the couples, 2002–2013 ( $\chi^2$ , N=5643 couples)

Variables	No mental disorders		Affective disorders		Anxiety disorders		Other mental disorders		$\chi^2$ p Value
	n <sub>1</sub>	Per cent	n <sub>2</sub>	Per cent	n <sub>3</sub>	Per cent	n <sub>4</sub>	Per cent	
Total	3230	57.24	189	3.35	1410	24.99	814	14.42	
<i>Insured spouse characteristics</i>									
Category of mental disorders									<0.0001*
No mental disorders	2118	62.22†	102	3.00	760	22.33	424	12.46	
Affective disorders	80	52.98	11	7.28‡	35	23.18	25	16.56	
Anxiety disorders	569	47.54	48	4.01	393	32.83§	187	15.62	
Other	463	51.96	28	3.14	222	24.92	178	19.98¶	
Gender									<0.0001*
Female	907	62.12	36	2.47	278	19.04	239	16.37	
Male	2323	55.53	153	3.66	1132	27.06	575	13.75	
Age (years)									<0.0001*
16–34	245	64.64	17	4.49	69	18.21	48	12.66	
35–44	814	62.86	52	4.02	265	20.46	164	12.66	
45–54	1000	62.23	46	2.86	367	22.84	194	12.07	
55–64	643	54.26	34	2.87	316	26.67	192	16.20	
≥65	528	44.86	40	3.40	393	33.39	216	18.35	
Occupation									<0.0001*
First category (private employee and government employee)	1755	60.85	100	3.47	658	22.82	371	12.86	
Second category (labour union member)	471	60.00	25	3.18	177	22.55	112	14.27	
Third category (farmer and fisherman)	348	50.07	13	1.87	203	29.21	131	18.85	
Fourth, fifth and sixth categories (soldier, social security insured and veteran and religious group member)	656	51.29	51	3.99	372	29.09	200	15.64	
Premium-based monthly salary (US\$)									0.0133*
≤576	984	54.79	65	3.62	485	27.00	262	14.59	
576–760	1033	56.36	54	2.95	452	24.66	294	16.04	
760–960	259	62.71	15	3.63	84	20.34	55	13.32	
960–1210	275	57.17	15	3.12	123	25.57	68	14.14	
1210–1526	297	62.79	16	3.38	117	24.74	43	9.09	
>1526	382	59.04	24	3.71	149	23.03	92	14.22	
Comorbidity (CCI)									<0.0001*
0	2337	60.65	129	3.35	864	22.42	523	13.57	
≥1	883	49.77	60	3.38	543	30.61	288	16.23	
Catastrophic illness and injury									0.0003*
Absent	2965	58.07	170	3.33	1262	24.72	709	13.89	
Present	265	49.35	19	3.54	148	27.56	105	19.55	
<i>Dependent spouse characteristics</i>									
Gender									<0.0001*
Female	2323	55.53	153	3.66	1132	27.06	575	13.75	
Male	907	62.12	36	2.47	278	19.04	239	16.37	
Age (years)									<0.0001*
16–34	437	66.82	25	3.82	119	18.20	73	11.16	
35–44	886	61.74	63	4.39	313	21.81	173	12.06	
45–54	837	60.17	37	2.66	326	23.44	191	13.73	
55–64	620	53.82	32	2.78	328	28.47	172	14.93	
≥65	450	44.51	32	3.17	324	32.05	205	20.28	
Comorbidity (CCI)									<0.0001*
0	2655	64.72	133	3.24	799	19.48	515	12.55	
≥1	558	36.61	56	3.67	611	40.09	299	19.62	
Catastrophic illness and injury									<0.0001*
Absent	3023	58.17	170	3.27	1286	24.75	718	13.82	
Present	207	46.41	19	4.26	124	27.80	96	21.52	
<i>Shared ecological characteristics</i>									
Region									0.1089
Taipei	1177	59.44	73	3.69	459	23.18	271	13.69	
Northern	512	58.92	25	2.88	216	24.86	116	13.35	
Central	584	54.28	34	3.16	298	27.70	160	14.87	

Continued



Table 2 Continued

Variables	No mental disorders		Affective disorders		Anxiety disorders		Other mental disorders		$\chi^2$ p Value
	n <sub>1</sub>	Per cent	n <sub>2</sub>	Per cent	n <sub>3</sub>	Per cent	n <sub>4</sub>	Per cent	
Southern	429	57.05	20	2.66	201	26.73	102	13.56	0.3627
Southeast	466	55.02	33	3.90	211	24.91	137	16.17	
Eastern	61	58.10	3	2.86	20	19.05	21	20.00	
Urbanisation level									
Level 1 (highest)	1014	58.11	66	3.78	417	23.90	248	14.21	
Level 2	972	58.38	60	3.60	404	24.26	229	13.75	
Level 3	548	57.44	24	2.52	247	25.89	135	14.15	
Level 4	399	53.99	23	3.11	193	26.12	124	16.78	
Level 5 (lowest)	290	55.88	15	2.89	144	27.75	70	13.49	

\*p&lt;0.05.

†Cell percentage =37.53%.

‡Cell percentage =0.19%.

§Cell percentage =6.96%.

¶Cell percentage =3.15%.

CCI, Charlson Comorbidity Index.

reached 19.97% (data not shown). Furthermore, the prevalence rates of the concordant categories of mental disorders including affective disorders, anxiety disorders and other mental disorders were 0.19%, 6.96% and 3.15%, respectively (data not shown). No significant collinearity was detected.

The multinomial logistic regression used no mental disorders as a referent of the outcome variable for all multivariate analyses. After adjustment for all other covariates, seven variables were significantly associated with the category of mental disorders among dependent spouses (table 3, all p<0.05). The highlighted cells in table 3 present the highest adjusted ORs and lowest p values in each of the three vertical columns of a certain disorder category in dependent spouses that was significantly associated with a concordant category in the insured spouses (all p<0.01). Specifically, compared with no mental disorders, when the insured spouses were diagnosed with affective disorders, the dependent spouses were significantly more likely to be diagnosed with affective disorders (OR=2.914, 95% CI 1.484 to 5.722, p=0.0019); the insured and dependent spouses were significantly more likely to be concordantly diagnosed with anxiety disorders when the insured spouses were diagnosed with anxiety (OR=1.776, 95% CI 1.505 to 2.096, p<0.0001); when the insured spouses were diagnosed with other mental disorders, the odds of both spouses being diagnosed with other mental disorders were significantly higher (OR=1.727, 95% CI 1.397 to 2.134, p<0.0001). Other factors significantly associated with the category of mental disorders among the dependent spouses included insured occupation, dependent gender, dependent age, dependent comorbidity, catastrophic illness and injury of dependent, and region (all p<0.05).

The results of the couple-level propensity-matched analysis are presented in tables 4 and 5. The prevalence rates of concordant affective disorders, anxiety disorders and other mental disorders were significantly higher in couples than in non-couples (table 4; 0.19% vs 0.02%, 6.96% vs 0.02%, 3.15% vs 0.11%, respectively;  $\chi^2$  test, p<0.0001). The multinomial logistic regression results indicated that couple status (relation) was significantly associated with categorical concordance (all p<0.05). Compared with no concordance (including discordance and concordant no mental disorders), the odds of concordant affective disorders, anxiety disorders and other mental disorders were significantly higher in couples than in

non-couples (table 5; OR=12.248, 437.528 and 33.031; 95% CI 1.581 to 94.897, 61.462 to 3114.61 and 14.630 to 74.580, respectively).

## DISCUSSION

### High aggregation of mental disorders

The literature indicates the phenomenon of family clustering of mental disorders.<sup>2 4 8</sup> On this basis, this study initially conducted a 2×2  $\chi^2$  test for the association between the mental disorder status of insured spouses and that of dependent spouses. Consequently, the aggregation of mental disorders and that of no mental disorders in couples were both determined (37.53% and 19.97%, respectively). The annual prevalence of psychiatric disorders in the entire population of Taiwan increased from 7.68% in 2002 to 10.54% in 2013.<sup>33</sup> Compared with the prevalence of 33.90% (2002–2013) at the individual level, the finding that aggregate mental disorders reached the prevalence of 19.97% at the couple level warrants further attention. This high aggregation can be explained by the aforementioned cohabitation effects and emotional contagion. Predisposition similarity and familial resources for stress coping<sup>34</sup> may denote whether psychiatric diseases will aggregate in a couple. Accordingly, the occurrence of mental disorders in a spouse can predict the mental disorder status in the partner. Previous studies have revealed the crucial role of family in coping with the mental illness of a family member,<sup>35</sup> because a family with a psychiatric patient may not seek outside help.<sup>36</sup> Although being the most critical stress-coping resource, an informal carer of a family would experience a substantial adaptation process of personal change concerning his or her role in social and cultural contexts.<sup>37</sup> The familial context in which the first spouse with mental disorders resides is inextricably interconnected with the psychological well-being of the second spouse. Therefore, a higher risk of aggregation of mental disorders emerges in couples. Furthermore, mental illness in one spouse eliciting psychological predicaments in the other spouse can be examined using the stress process model.<sup>34</sup> When a spouse is mentally ill, the partner assumes the role of an informal carer who may undergo primary and secondary stressors. Primary stressors pertain to the direct experiences from the illness, including the perception of role captivity in the carer. When chronic stress perforates into other areas of the carer's family and work life,

**Table 3** Association between the category of mental disorders among insured spouses and that of mental disorders among dependent spouses, after adjustment for all other variables, 2002–2013 (multinomial logistic regression, N=5643 couples)

Variables	Affective disorders/no mental disorders			Anxiety disorders/no mental disorders			Other mental disorders/no mental disorders		
	Adjusted OR	95% CI	p Value	Adjusted OR	95% CI	p Value	Adjusted OR	95% CI	p Value
<i>Insured spouse characteristics</i>									
Category of mental disorders									
No mental disorders (referent)	–	–	–	–	–	–	–	–	–
Affective disorders	2.914	1.484 to 5.722	0.0019*	1.180	0.770 to 1.808	0.4475	1.572	0.979 to 2.525	0.0613
Anxiety disorders	1.770	1.219 to 2.569	0.0027*	1.776	1.505 to 2.096	<0.0001*	1.481	1.205 to 1.820	0.0002*
Other	1.181	0.761 to 1.831	0.4580	1.164	0.961 to 1.409	0.1196	1.727	1.397 to 2.134	<0.0001*
Occupation									
First category (private employee and public servant, referent)	–	–	–	–	–	–	–	–	–
Second category (labour union member)	0.793	0.435 to 1.446	0.4495	1.063	0.804 to 1.405	0.6694	0.887	0.642 to 1.225	0.4657
Third category (farmer and fisherman)	0.522	0.239 to 1.139	0.1024	1.285	0.937 to 1.762	0.1203	1.274	0.886 to 1.831	0.1921
Fourth, fifth and sixth categories (soldier, social security insured and veteran and religious group member)	1.582	0.848 to 2.954	0.1497	1.359	1.040 to 1.776	0.0245*	1.419	1.020 to 1.973	0.0375*
<i>Dependent spouse characteristics</i>									
Gender									
Female (referent)	–	–	–	–	–	–	–	–	–
Male	0.584	0.386 to 0.882	0.0106*	0.552	0.462 to 0.660	<0.0001*	0.980	0.803 to 1.197	0.8453
Age (years)									
16–34 (referent)	–	–	–	–	–	–	–	–	–
35–44	1.553	0.874 to 2.758	0.1331	1.329	1.000 to 1.766	0.0501	1.290	0.910 to 1.828	0.1523
45–54	0.997	0.483 to 2.057	0.9939	1.399	1.002 to 1.952	0.0484*	1.493	0.991 to 2.249	0.0553
55–64	0.899	0.366 to 2.206	0.8156	1.500	1.004 to 2.242	0.0479*	1.442	0.882 to 2.360	0.1448
≥65	0.852	0.291 to 2.495	0.7707	1.397	0.870 to 2.243	0.1659	1.808	1.017 to 3.213	0.0435*
Comorbidity (CCI)									
0 (referent)	–	–	–	–	–	–	–	–	–
≥1	2.078	1.438 to 3.002	<0.0001*	3.403	2.909 to 3.981	<0.0001*	2.285	1.888 to 2.764	<0.0001*
Catastrophic illness and injury									
Absent (referent)	–	–	–	–	–	–	–	–	–
Present	1.485	0.881 to 2.506	0.1381	0.883	0.686 to 1.137	0.3346	1.326	1.008 to 1.743	0.0434*
<i>Shared ecological characteristics</i>									
Region									
Taipei (referent)	–	–	–	–	–	–	–	–	–
Northern	0.860	0.509 to 1.455	0.5747	1.072	0.854 to 1.346	0.5473	0.976	0.739 to 1.289	0.8660
Central	1.006	0.622 to 1.627	0.9799	1.287	1.043 to 1.589	0.0188*	1.191	0.923 to 1.537	0.1789
Southern	0.837	0.464 to 1.507	0.5524	1.083	0.847 to 1.384	0.5257	0.975	0.723 to 1.315	0.8667
Southeast	1.256	0.791 to 1.995	0.3338	1.121	0.901 to 1.396	0.3051	1.288	0.996 to 1.664	0.0537
Eastern	0.864	0.251 to 2.970	0.8163	0.779	0.445 to 1.364	0.3828	1.522	0.866 to 2.675	0.1447

Only variables statistically significant are shown.

\*p<0.05.

CCI, Charlson Comorbidity Index.

secondary stressors emerge and affect the well-being of the carer.<sup>14 38</sup> Furthermore, a longitudinal study revealed that the association between depressive symptoms of husbands and the subsequent depressive symptoms of wives was stronger among couples reporting marital distress compared with those reporting higher marital satisfaction.<sup>17</sup> Briefly, mental disorders are costly and debilitating for ill and healthy spouses.<sup>39</sup> Therefore, the coping strategies of an informal carer and formal social support outside the family are pivotal to the mental health concerns in couples.<sup>18 36 37</sup> Releasing the care burden imposed on the healthy spouse is key in the intervention to prevent the new development of psychiatric morbidity and to provide the mentally ill spouse an opportunity to recover in a supportive family environment.<sup>5</sup>

### Phenomenon of parallel contagion

This study identified three concordant categories of mental disorders in married couples. The highlighted cells in table 2 present the row percentages of concordance for each category, including no mental disorders, affective disorders, anxiety disorders and other mental disorders. Table 3 further provides the adjusted ORs for the three concordances in a manner mirroring the findings presented in table 2. An additional analysis was performed using mental disorder status of dependent spouses as the independent variable and that of insured spouses as the outcome variable. This analysis which swapped the two major dependent and independent variables substantially generated same results (data not shown). Insured and dependent spouses were most likely to experience a concordant category of mental

**Table 4** Concordant mental disorders in couples versus non-couples, 2002–2013 (with dual propensity score matching for gender, age and Charlson Comorbidity Index;  $\chi^2$  test; N=11 286 pairs)

Variables	Concordant affective disorders		Concordant anxiety disorders		Concordant other mental disorders		No concordance		$\chi^2$ p Value
	n <sub>1</sub>	Per cent	n <sub>2</sub>	Per cent	n <sub>3</sub>	Per cent	n <sub>4</sub>	Per cent	
Relation									<0.0001*
Non-couple	1	0.02	1	0.02	6	0.11	5635	99.86	
Couple	11	0.19	393	6.96	178	3.15	5061	89.69	

\*p<0.05, with 1:1 propensity score matching according to gender, age and Charlson Comorbidity Index for the four individuals in the couples and non-couple pairs.

**Table 5** Association between couple status and concordant mental disorders, 2002–2013 (with dual propensity score matching for gender, age and Charlson Comorbidity Index; multinomial logistic regression; N=11 286 pairs)

Variables	Concordant affective disorders/no concordance			Concordant anxiety disorders/no concordance			Concordant other mental disorders /no concordance		
	OR	95% CI	p Value	OR	95% CI	p Value	OR	95% CI	p Value
Relation									
Non-couple	–	–	–	–	–	–	–	–	–
Couple	12.248	1.581 to 94.897	0.0165*	437.528	61.462 to 3115.25	<0.0001*	33.031	14.630 to 74.580	<0.0001*

\*p<0.05.

OR, with 1:1 propensity score matching according to gender, age and Charlson Comorbidity Index for the four individuals in the couples and non-couple pairs.

disorders among the three categories. If one spouse was diagnosed with affective disorders, the other spouse would most likely undergo affective disorders correspondingly. If one spouse experienced anxiety disorders, the same type of disease might occur in the other spouse. The phenomenon coherently applied to other mental disorders. This particular finding, further confirmed strongly in the matched case–control analysis, was named the phenomenon of parallel contagion, an empirical derivative from the theory of emotional contagion.<sup>14</sup> Signalling the concordance in mental illness, parallel contagion justifies the necessity of epidemiological prevention, particularly for the same type of mental disorders in couples. Notably, the contagion identified is not exclusive of different categories of mental disorders. Dependent spouses were also more likely to undergo affective disorders even if insured spouses were diagnosed with anxiety disorders (OR=1.770); a strong association was observed between anxiety disorders in insured spouses and other mental disorders in dependent spouses (OR=1.481). Although the association between the categories of mental disorders among insured spouses and those of dependent spouses could be nonparallel (discordant), the phenomenon of parallel contagion is evident and thus delivers essential clinical implications. Future research is required to explore the concordance within other categories of mental disorders in couples.

### Other characteristics predicting risk

Soldiers, social security insureds and veterans were closely related to higher odds of anxiety and other mental disorders, and this finding echoes the findings of previous studies.<sup>40–41</sup> As the possibility of two spouses of a household sharing common occupational stress is high,<sup>42</sup> the occupation of insured spouses is a determinant of the category of mental disorders among dependent spouses.

The present study has certain limitations that should be addressed. First, the secondary data used did not include information on health behaviour and perceived stress; therefore, the absence of such information might attenuate the statistical

testing power. Nevertheless, this study fully used all obtainable data for analysis. Second, the actual length of marriage and cohabitation could not be ascertained from the NHIRD, thus limiting further analysis in this study. Finally, spouses who could be identified from the database were limited to the insured–dependent relationship. The study findings must be cautiously extrapolated to all other scenarios.

### CONCLUSION

This study demonstrates that shared environmental factors contribute to the development of mental diseases in couples. When one spouse is diagnosed with mental disorders, the other spouse is at risk of mental illness. A family clustering of mental disorders was further determined to demonstrate the phenomenon of parallel contagion. The concordance of mental disorders in couples was identified in the aggregation form of affective–affective, anxiety–anxiety and other–other disorders. Although different from conventional contagion in epidemiology, the phenomenon of parallel contagion substantially carries clinical implications in developing the prospective direction of family-

### What this study adds?

- ▶ The prevalence rates for concordant categories of mental disorder are 0.19% for affective disorders, 6.96% for anxiety disorders and 3.15% of other mental disorders.
- ▶ Two spouses are significantly likely to be diagnosed with the same category of mental disorders including affective, anxiety and other disorders.
- ▶ Other associated factors include age, occupation, comorbidity and region.
- ▶ This study proposes 'the phenomenon of parallel contagion'. A category of mental disorder in one spouse is a determinant of that in the other spouse.

## What is already known on this subject?

- ▶ The aggregation of mental disorders in couples has been reported by prior research, which indicates the effect of shared environment on psychiatric epidemiology and warrants attention.
- ▶ The concordant categories of mental disorders in couples and the associated factors remain fragmented and inconclusive.

based mental health practice. Establishing an external support system toward mentally vulnerable family members is a policy priority. Policymakers should target spouses whose married partners were mentally ill by using the risk factors identified by this study for preventing mental disorders, particularly the same category of mental disorders.

**Acknowledgements** The authors are grateful to Taiwan's National Health Research Institutes for providing the NHIRD.

**Contributors** J-YW proposed the study idea, designed the study, interpreted the analysis and wrote the first draft of the manuscript. M-HH contributed to the theoretical construction and revised the manuscript. P-CL contributed to the literature review and performed the statistical analyses. C-SL provided guidance and aided in proofreading the manuscript. J-DC participated in critical discussion and edited the manuscript.

**Funding** This research was supported by Ministry of Science and Technology, Taipei, Taiwan (grant no. NSC 102-2410-H-039-006-SS2 and MOST 104-2410-H-039-006).

**Competing interests** None declared.

**Ethics approval** The institutional review board at China Medical University Hospital.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Open Access** This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

## REFERENCES

- 1 Lee JS, Joo EJ, Choi KS. Perceived stress and self-esteem mediate the effects of work-related stress on depression. *Stress Health* 2013;29:75–81.
- 2 Isomura K, Boman M, Ruck C, et al. Population-based, multi-generational family clustering study of social anxiety disorder and avoidant personality disorder. *Psychol Med* 2015;45:1581–9.
- 3 Mataix-Cols D, Boman M, Monzani B, et al. Population-based, multigenerational family clustering study of obsessive-compulsive disorder. *JAMA Psychiatry* 2013;70:709–17.
- 4 Hippisley-Cox J, Coupland C, Pringle M, et al. Married couples' risk of same disease: cross sectional study. *BMJ* 2002;325:636.
- 5 Joutsenniemi K, Moustgaard H, Koskinen S, et al. Psychiatric comorbidity in couples: a longitudinal study of 202,959 married and cohabiting individuals. *Soc Psychiatry Psychiatr Epidemiol* 2011;46:623–33.
- 6 Wittmund B, Wilms HU, Mory C, et al. Depressive disorders in spouses of mentally ill patients. *Soc Psychiatry Psychiatr Epidemiol* 2002;37:177–82.
- 7 Maes HH, Neale MC, Kendler KS, et al. Assortative mating for major psychiatric diagnoses in two population-based samples. *Psychol Med* 1998;28:1389–401.
- 8 Kendler KS, Ohlsson H, Sundquist K, et al. Environmental clustering of drug abuse in households and communities: multi-level modeling of a national Swedish sample. *Soc Psychiatry Psychiatr Epidemiol* 2015;50:1277–84.
- 9 Agrawal A, Heath AC, Grant JD, et al. Assortative mating for cigarette smoking and for alcohol consumption in female Australian twins and their spouses. *Behav Genet* 2006;36:553–66.
- 10 Reynolds CA, Barlow T, Pedersen NL. Alcohol, tobacco and caffeine use: spouse similarity processes. *Behav Genet* 2006;36:201–15.
- 11 Knuiaman MW, Divitini ML, Welborn TA, et al. Familial correlations, cohabitation effects, and heritability for cardiovascular risk factors. *Ann Epidemiol* 1996;6:188–94.
- 12 Knuiaman MW, Divitini ML, Bartholomew HC, et al. Spouse correlations in cardiovascular risk factors and the effect of marriage duration. *Am J Epidemiol* 1996;143:48–53.
- 13 Bookwala J, Schulz R. Spousal similarity in subjective well-being: the cardiovascular health study. *Psychol Aging* 1996;11:582–90.
- 14 Goodman CR, Shippy RA. Is it contagious? Affect similarity among spouses. *Aging Ment Health* 2002;6:266–74.
- 15 Hill R. Generic features of family under stress. *Social Casework* 1958;49:139–50.
- 16 Pierce GR. Social support in couples: marriage as a resource in times of stress (Book). *Journal of Marriage & Family* 1997;59:494–5.
- 17 Kouros CD, Cummings EM. Longitudinal associations between husbands' and wives' depressive symptoms. *J Marriage Fam* 2010;72:135–47.
- 18 Benazon NR, Coyne JC. The next step in developing an interactional description of depression? *Psychological Inquiry* 1999;10:279–304.
- 19 Tower RB, Kasl SV. Depressive symptoms across older spouses and the moderating effect of marital closeness. *Psychol Aging* 1995;10:625–38.
- 20 Falba TA, Sindelar JL. Spousal concordance in health behavior change. *Health Serv Res* 2008;43:96–116.
- 21 Homish GG, Leonard KE. Spousal influence on general health behaviors in a community sample. *Am J Health Behav* 2008;32:754–63.
- 22 Stimpson JP, Masel MC, Rudkin L, et al. Shared health behaviors among older Mexican American spouses. *Am J Health Behav* 2006;30:495–502.
- 23 Bronfenbrenner U. *The ecology of human development: experiments by design and nature*. Cambridge, MA: Harvard University Press, 1979.
- 24 Ildstad M, Roysamb E, Tambs K. The effect of change in mental disorder status on change in spousal mental health: the HUNT study. *Soc Sci Med* 2011;73:1408–15.
- 25 Sareen J, Afifi TO, McMillan KA, et al. Relationship between household income and mental disorders: findings from a population-based longitudinal study. *Arch Gen Psychiatry* 2011;68:419–27.
- 26 Dinkel A, Krensreiter K, Marten-Mittag B, et al. Comorbidity of fear of progression and anxiety disorders in cancer patients. *Gen Hosp Psychiatry* 2014;36:613–9.
- 27 Ducat L, Phillipson LH, Anderson BJ. The mental health comorbidities of diabetes. *JAMA* 2014;312:691–2.
- 28 Fan ZJ, Bonauto DK, Foley MP, et al. Occupation and the prevalence of current depression and frequent mental distress, WA BRFS 2006 and 2008. *Am J Ind Med* 2012;55:893–903.
- 29 Wulsin L, Alterman T, Bushnell P, et al. Prevalence rates for depression by industry: a claims database analysis. *Soc Psychiatry Psychiatr Epidemiol* 2014;49:1805–21.
- 30 Malard L, Chastang JF, Niedhammer I. Changes in major depressive and generalized anxiety disorders in the national French working population between 2006 and 2010. *J Affect Disord* 2015;178:52–9.
- 31 Ruiz-Grosso P, Ramos M, Samalvides F, et al. Common mental disorders in public transportation drivers in Lima, Peru. *PLoS ONE* 2014;9:e101066.
- 32 Deyo RA, Cherkin DC, Ciol MA. Adapting a clinical comorbidity index for use with ICD-9-CM administrative databases. *J Clin Epidemiol* 1992;45:613–9.
- 33 Ministry of health and welfare. Hospital and Clinical Statistics. 2016 (cited 15 August 2016). [http://www.mohw.gov.tw/EN/Ministry/Statistic.aspx?f\\_list\\_no=474](http://www.mohw.gov.tw/EN/Ministry/Statistic.aspx?f_list_no=474)
- 34 Pearlín LI, Mullan JT, Semple SJ, et al. Caregiving and the stress process: an overview of concepts and their measures. *Gerontologist* 1990;30:583–94.
- 35 Wouters E, Masquillier C, le Roux Booyesen F. The importance of the family: a longitudinal study of the predictors of depression in HIV patients in South Africa. *AIDS Behav* 2016;20:1591–602.
- 36 Thompson G, McBride RB, Hosford CC, et al. Resilience among medical students: the role of coping style and social support. *Teach Learn Med* 2016;28:174–82.
- 37 Moreno-Camara S, Palomino-Moral PA, Moral-Fernandez L, et al. [Problems in the process of adapting to change among the family caregivers of elderly people with dementia]. *Gac Sanit* 2016;30:201–7.
- 38 Goodman CR, Zarit SH, Steiner VL. Personal orientation as a predictor of caregiver strain. *Aging Ment Health* 1997;1:149–57. 9p.
- 39 Trivedi RB, Post EP, Sun H, et al. Prevalence, comorbidity, and prognosis of mental health among US veterans. *Am J Public Health* 2015;105:2564–9. 6p.
- 40 Hageman I, Pinborg A, Andersen HS. Complaints of stress in young soldiers strongly predispose to psychiatric morbidity and mortality: Danish national cohort study with 10-year follow-up. *Acta Psychiatr Scand* 2008;117:148–55.
- 41 Diala CC, Muntaner C, Walrath C. Gender, occupational, and socioeconomic correlates of alcohol and drug abuse among US rural, metropolitan, and urban residents. *Am J Drug Alcohol Abuse* 2004;30:409–28.
- 42 Tsukahara I. The effect of family background on occupational choice. *Labour* 2007;21:871–90.