are exposed to possible presentation bias resulting from the presence of the researcher and from bias inherent in relying upon a small self-selected sample of women who are motivated to participate in research and/or may have a particular interest in GDM. An alternative research approach is to opportunistically use 'found' data that is spontaneously user-generated within online discussion forums. The objective of this study was to explore the perceptions and beliefs surrounding GDM suggested within online parental-support forums for parents, that could have implications for future T2D prevention.

Methods A search was carried out using the Google search engine to identify online discussion forums that could be of relevance to women with GDM in the UK. We screened these 10 (out of 120 forums) to identify those containing relevant data that could be considered public, and selected Mumsnet and Netmums to be included in the study. They were comprehensively searched using the search term 'gestational diabetes or GD', and relevant posts from 1/1/2017 – 14/2/2019 were identified. Multiple messages from the same person were linked together as a single unit, analogous to a research participant, using an anonymised identifier. A theoretical framework derived previously was used to code and sort the data using framework approach.

Results A total of 646 posts in 137 threads from 282 unique users with current/previous GDM were analysed. Five prominent themes were identified: Emotional response to and understanding of diagnosis; personal responsibility; consequences and impact of GDM; lifestyle change; and Type 2 diabetes. Users' discussions around these themes highlighted three implicit 'messages' regarding GDM: that GDM is not a serious or even real diagnosis and the consequences are not severe; that women need not take personal responsibility for GDM; and there are minimal implications for T2D prevention.

Discussion These (partially) subliminal messages will all mitigate against efforts to encourage women to change their lifestyles for future T2D prevention. This is worrying as online networks have huge reach, and are viewed by millions of visitors who don't necessarily post themselves, but are signposted to these sites when they pose questions to global search engines. The data analysed were generated in a naturalistic setting and are perhaps closer than traditional qualitative research to women's real perceptions and beliefs surrounding GDM.

OP40 EXPLORING BARRIERS AND FACILITATORS FOR WOMEN TO SEEK HELP FOR REPRODUCTIVE SYMPTOMS IN THE UNITED KINGDOM: A RAPID REVIEW

¹AA Ayorinde*, ²F Boardman, ³L Porter, ³A Hadley, ³M Ludeke, ³A Sallis, ³S Mann, ¹ORO Oyeboade. ¹Warwick Centre for Applied Health Research and Delivery (W-CAHRD), University of Warwick, Coventry, UK; ²WMS – Social Science and Systems in Health, University of Warwick, Coventry, UK; ³Public Health England, London, UK

10.1136/jech-2020-SSMabstracts.40

Background Many women experience reproductive symptoms that impact on their wellbeing but do not seek help for a range of different reasons. In this rapid review we explored the factors that enable or hinder women to seek help for unwanted or debilitating reproductive symptoms. We also aimed to examine whether existing interventions address

these factors and identify opportunities for alternative interventions.

Methods We searched electronic databases MEDLINE, PsycINFO and CINAHL in October 2019 and contacted experts to identify grey literature and additional interventions. The search strategy included terms covering behaviour and intervention combined with terms relating to common reproductive symptoms such as, dysmenorrhoea and menopause. We limited to articles published in English since 2009. We included any study that presented facilitators and barriers to seeking help for reproductive symptoms that was conducted with women in the United Kingdom. We also included articles on relevant interventions. One reviewer screened titles and abstracts with 10% double screened by a second reviewer. Full texts were screened independently by two reviewers. We extracted study characteristics using a pre-piloted form and used the Mixed Methods Appraisal Tool for quality assessment. Data extraction and quality assessment were performed by one reviewer and a sample checked by a second reviewer. We coded information on barriers and facilitators from each study into themes using NVivo. Results were summarised descriptively.

Results We screened 12474 citations, 102 full-text articles and retained 22 articles for analysis. Ten help-articles focused on interventions. Of the 12 articles reporting barriers and facilitators, the most common barriers to seeking help were embarrassment, perception that symptom is too private to talk about, dissatisfaction with the information received from healthcare professionals, poor knowledge, perception that symptom is part of aging and difficulty explaining the symptoms to others. Facilitators were having access to useful information/information pitched at the right level, an information source women felt comfortable with and supportive work environments (including awareness among managers and flexible working).

Conclusion This review identified a range of barriers and facilitators spanning across capability, opportunity and motivation. We have categorised the barriers and facilitators into the Theory and Techniques Tool (TaTT) mechanisms of actions (MoAs). Our next step is to assess whether the existing intervention contents match those deemed theoretically appropriate using TaTT in order to identify missed opportunities for intervention. This will enable us to make recommendations for interventions to support women to seek help for reproductive symptoms when necessary.

OP41 ASSOCIATIONS BETWEEN MODE OF DELIVERY AND OFFSPRING OVERWEIGHT/OBESITY: FINDINGS FROM THE STUDYING LIFECOURSE OBESITY PREDICTORS (SLOPE) POPULATION-BASED COHORT

¹A Simpson*, ¹G Grove, ¹N Ziauddeen, ^{1,2}NA Alwan. ¹School of Primary Care, Population Sciences and Medical Education, University of Southampton, Southampton, UK; ²NIHR Southampton Biomedical Research Centre, University of Southampton and University Hospital Southampton NHS FT, Southampton, UK

10.1136/jech-2020-SSMabstracts.41

Background Childhood obesity affects one in ten children in England by age 5, and one in five by age 11. Existing evidence suggests a possible relationship between caesarean section (CS) birth and higher risk of overweight/obesity in childhood, however maternal obesity is a strong confounder