preconception planning is generally low. In this rapid evidence review, we examined the barriers and facilitators to women choosing to plan and prepare for a healthy pregnancy.

Methods We searched MEDLINE, PsycINFO and CINAHL from 2009 to October 2019 and limited to English language. We included any publications that presented facilitators and barriers for women choosing to plan and prepare for pregnancy. We also included papers presenting barriers and facilitators to health care professionals supporting this behaviour and articles on relevant interventions although these data are not presented here. We contacted experts to identify grey literature. We extracted study characteristics using a pre-piloted data extraction form and assessed the quality of individual studies using the Mixed Methods Appraisal Tool. One reviewer performed title and abstract screening, data extraction and quality assessment with a sample checked by a second reviewer. Two reviewers screened full texts independently. Using NVivo, we coded information on barriers and facilitators from each study into themes under two subheadings; a) information seeking and b) preparing for a healthy pregnancy.

Results We screened 2679 citations, 54 full-text articles and included 24 articles for analysis. 18 articles reported barriers and facilitators for women. The most frequently reported barriers to information seeking prior to conception were unintended pregnancy, information provoking anxiety, and belief that there is no need for preconception care. Facilitators included ad hoc prompts in health care settings, and opportunities to discuss pregnancy intentions (e.g. as part of care for a chronic condition). The most frequent barriers to preparing for a healthy pregnancy were: not knowing what recommended behaviours might be, lack of understanding or incorrect beliefs, information not appropriate for woman's context, and lived experience (of self or in social circle) that appears to contradict health advice. Facilitators included knowledge of recommended behaviours, feelings of responsibility towards a potential baby, and confidence in ability to achieve health goals prior to conceiving.

Conclusion We will conduct a behavioural analysis and categorise the identified barriers and facilitators into the Theory and Techniques Tool (TaTT) mechanisms of actions (MoAs). We will then examine whether existing intervention content matches what is theoretically appropriate, therefore identifying opportunities for improvement of existing interventions and novel development to promote preconception planning and ultimately, better maternal and neonatal outcomes.

P75

SYSTEMATIC REVIEW OF THE EFFECTIVENESS OF RIGHTS-BASED APPROACHES TO SEXUAL AND REPRODUCTIVE HEALTH IN LOW AND MIDDLE-INCOME COUNTRIES

¹M McGranahan*, ²J Nakyeeyune, ²C Baguma, ²NN Musisi, ²D Nsibirwa, ³S Sekalala, ¹O Oyeboode. ¹Warwick Medical School, University of Warwick, Coventry, UK; ²Center for Health, Human Rights and Development, Kampala, Uganda; ³School of Law, University of Warwick, Coventry, UK

10.1136/jech-2020-SSMabstracts.167

Background By 2030, the Sustainable Development Goals aim to achieve human rights for all. This involves empowering women and girls (Goal 5) and ensuring that everyone can access sexual and reproductive health rights (Goal 3). This is

the first systematic review of the effectiveness of rights-based approaches to sexual and reproductive health including gender-based violence (GBV), maternity, HIV and sexually transmitted infections (STI).

Aim To determine the effectiveness of rights-based approaches to sexual and reproductive health in low and middle-income countries (LMICs).

Methods Search Strategy EMBASE, MEDLINE and Web of Science were searched until 9/1/2020. Inclusion criteria were:

- Study design: any interventional study
- Population: adolescent and adult females
- Setting: LMICs
- Intervention: a 'rights-based approach' (defined by the author) and/or interventions that the author explicitly stated related to 'rights'.

Study selection, data extraction and risk of bias assessment was undertaken independently by two authors.

Synthesis A narrative synthesis of included studies was undertaken, and outcomes mapped to identify evidence gaps.

Risk of bias The RoB-2 tool and the ROBINS-I tool were used to assess risk of bias of cluster-randomised studies and non-randomised studies respectively. Uncontrolled before-after studies based on two cross-sectional surveys were given an overall assessment of serious or critical risk of bias.

Results Of 17,212 records identified through database searching, 13,404 records remained after de-duplication. Sixty-nine studies remained following title and abstract screening, of which seven were included after full-text screening. Reference list screening identified seventeen studies.

Rights-based interventions were effective for most included outcomes, but evidence was of poor quality. Testing uptake for HIV and/or other STIs improved with intervention but all relevant studies were at critical or serious risk of bias. Condom use improved with intervention (although one study showed no change), but all relevant studies were at high, serious or critical risk of bias. Awareness of rights improved with intervention, but all four studies were at critical or serious risk of bias.

Conclusion This is the first systematic review to evaluate the effect of rights-based approaches to sexual and reproductive health including maternity, GBV and STIs/HIV. Considerable risk of bias in all studies means results must be interpreted with caution.

Priority and relevance Rights-based approaches are often recommended but high-quality controlled studies are needed urgently to determine if they are effective for sexual and reproductive health in LMICs.

This systematic review was written through a collaboration between the University of Warwick and the Center for Health Human Rights & Development (CEHURD), Uganda.

P76

THE ASSOCIATIONS BETWEEN PREGNANCY LOSS AND CARDIOVASCULAR DISEASES IN THE UK BIOBANK

¹FM Simmons-Jones*, ²C Oliver-Williams. ¹Public Health, Essex County Council, Chelmsford, UK; ²Cardiovascular Epidemiology Unit, Department of Public Health and Primary Care, University of Cambridge, Cambridge, UK

10.1136/jech-2020-SSMabstracts.168

Rationale This study assessed the associations between pregnancy loss (miscarriage, stillbirth and abortion) and future

maternal cardiovascular diseases (CVD, myocardial infarction (MI), stroke, ischaemic stroke, intracerebral haemorrhage and subarachnoid haemorrhage) in women in the UK Biobank from 2006–2016.

Cardiovascular diseases (CVDs) are a high priority public health issue, presenting a high disease burden. Pregnancy has been described as a stress test for women's bodies, where complications with pregnancy, including pregnancy loss, may be an indicator of increased risk of future CVDs. However, findings regarding associations between pregnancy loss and future CVDs are mixed, with a scarcity of UK studies. Furthering understanding of female specific risk factors has the potential to increase the clinical utility of risk prediction and public health prevention.

Methods A prospective cohort study was conducted, including women in the UK Biobank cohort with self-reported exposure status recorded at baseline ($n=246,124$), with a median follow up time of 7.03 years (IQR 6.36–7.75) and a total of 1,762,729 person-years. Cardiovascular outcomes were ascertained through linkage of participant healthcare records and UK mortality databases. Cox proportional hazards models were used to ascertain Hazard ratios (HR) and 95% confidence intervals (CI). Patient and participant involvement (PPI) was included to inform understanding of public health impacts of study findings.

Results Within the study cohort, 79,482 (32.29%) women had experienced one or more pregnancy losses. During follow up 2,567 (1.04%) women developed CVD. A history of one or more stillbirths was associated with an increased risk of stroke (Adjusted HR 1.55, 95%CI 1.06–2.27), in particular subarachnoid haemorrhage (Adjusted HR: 1.52, 95%CI 0.62–3.74). Miscarriage was not consistently associated with an increased risk of any cardiovascular outcome and no association between a history of therapeutic abortion and increased risk of CVD was demonstrated.

Conclusion Analyses of the relationship between pregnancy loss and subtypes of CVD indicated that the direction and magnitude of associations were not universal in women in the UK Biobank or across populations. There is evidence of association between some types of pregnancy loss are associated with an increased risk of CVDs, though further research into associations with haemorrhagic stroke, and the associations between abortion and CVD, are needed. History of miscarriage and/or stillbirth should be considered for inclusion in cardiovascular risk assessment tools, potential pooled with other pregnancy complications, to identify and target support for women at increased risk.

P77

COMPARING THE RISK OF PREMATURE BIRTH FOLLOWING ABORTION WITH THE RISK AFTER MISCARRIAGE – A SYSTEMATIC REVIEW AND META-ANALYSES

¹M Eames*, ²G Gardner. ¹*The Acorns, Public Health Relations, Hatfield, UK;* ²*Primary Care, Birmingham Medical School, Birmingham, UK*

10.1136/jech-2020-SSMabstracts.169

Background Premature births have increased in the UK in the last decade. The incidence of PTB in Scotland is now at an all-time high. The cause remains a mystery. We aim to

quantify the risk of moderate PTB (mPTB<37 weeks), very PTB (vPTB<32 weeks) and extremely PTB (xPTB<28 weeks) allowing for other risk factors, following single and multiple abortions, and single and multiple miscarriages using a series of systematic meta-analysis of papers published 1990–2019.

Methods Systematic review* using PubMed, Cochrane, and Embase of papers published 1990–2019 and meta-analyses of observational studies were conducted separately for ToP and miscarriage. 67 separate studies were identified from 22 eligible papers for ToP; Further investigation allowed analysis relating to numbers of ToP, numbers of miscarriages and the degree of prematurity according to gestational birth age. Bias and Heterogeneity are measured and considered.

Results Miscarriage: AdjOR between 1.70, 95% CI (1.51–1.92) from Denmark and 1.12 (1.08–1.16) from Scotland for PTB after one miscarriage were noted from papers. These risks increased after two miscarriages with adjOR of between 2.20 (0.70–2.0) from Seattle and 1.36 (1.25–1.47) from Scottish data; AdjOR for xPTB after two or more miscarriages were between 4.0 (2.3–7.1) from Sweden and 2.81 (1.47–5.38) from Scotland. Iran reported Adj OR for xPTB of 4.10 (2.08–8.08) after three+ miscarriages.

Abortion and Meta-analysis: Fifty of the 67 studies demonstrated a significant increased risk of PTB related to abortion. Data was obtained from 22 countries worldwide. Risk of PTB after one+ abortion carried an Adj OR of 1.52 95% CI (1.43–1.62) compared to matched women who had no ToP (67 studies); Risk for corresponding miscarriage was 1.31 (1.18–1.45). Risk of vPTB after either ToP or miscarriage increased with increasing numbers of both ToP and miscarriage. The greatest risk increase noted was for xPTB after three+ ToP with Adj OR of 5.22 (1.58–17.21), whilst for three+ miscarriages it was 3.87 (2.85–5.26).

Risk of PTB was also measured according to method of abortion, or miscarriage treatment, and Adj OR compared for medical versus surgical treatments.

Conclusion The likelihood of xPTB increased after multiple Top (3 or more) and increased for miscarriages but to a lesser degree. The risk of any PTB also increases with multiple ToP and with several miscarriages. This is an important public health finding for women's choices for consent to ToP or to treatment after miscarriage in the UK. It has implications for costs, future research and reduction of premature births in the UK.

- * Included studies 6005 titles and abstracts were identified 1990–2019 and papers were screened for eligibility. 43 papers were selected for systematic review, from which 23 papers (with 3,796,010 participants) met the inclusion criteria for the meta-analysis for ToP.
- Papers which did not distinguish miscarriage from abortion, and studies with overlapping data were excluded.
- The search strategy used MESH terms for Abortion or ToP (('Abortion, Induced' or 'Abortion, Legal' or 'Abortion, Therapeutic', 'Termination of pregnancy') AND ('infant, premature' or 'obstetric labor, premature' or 'premature birth' or 'preterm birth' 'fetal membranes, premature rupture' or 'Pregnancy complications, or 'pregnancy outcome'.
- For Miscarriage: ('miscarriage'; 'spontaneous abortion'; 'pregnancy loss')