frequency in the past four weeks respectively, at baseline. Calculating change scores over time, we assessed whether changes in social media belonging, were associated with changes in drinking status, and whether changes in frequency of use of social media sites were related to changes in drinking frequency, using multinomial logistic and linear regression model. Analyses were repeated with number of times binge drinking (5+ drinks) in the past four weeks, on a sub-sample of participants aged 16–19. Models adjusted for sex, age, household tenure, urban/rural area, educational/work status, baseline social media use/frequency and number of friends.

Results At baseline, those who belonged to a social media site were over four times more likely (OR=4.62 95% CI (3.61–5.91)) to be a drinker versus non-drinker. Those who chatted for longer on social media sites were more likely to have drunk at least once a week compared with not drinking in the past month (none vs. 1–3 hours (2.31(1.45–3.68), 4–6 hours 2.60(1.44–4.72), 7 hours or more 5.92 (2.86–12.28)). Among participants aged 16–19, a similar gradient was found with higher odds of binge drinking three or more times in the past month, with greater chat frequency (e.g. no hours chatting vs. 7 hours or more 6.29 (2.50–15.82)). Over time, becoming new social media users increased the odds of becoming a new drinker (1.77 (1.04–3.03)) versus no change in drinking status. Participants who became more frequent users of social media sites, were more likely to drink more frequently ($\beta=0.13$, 95% CI (0.09–0.17)), and binge drink ($\beta=0.21$, 95% CI (0.12–0.31)) more frequently across waves.

Conclusion Increasing social media use was associated with greater alcohol consumption; mechanisms behind this relationship should be investigated. Alcohol-related content on the internet should be monitored.

P89

TACKLING LOCAL ALCOHOL-RELATED HARMS IN THE NIGHT TIME ECONOMY: A PROCESS EVALUATION WITH A COMPLEX SYSTEMS PERSPECTIVE

¹ET McGill*, ²D Marks, ³M Petticrew, ³M Egan. ¹Department of Health Services, Research and Policy, London School of Hygiene and Tropical Medicine, London, UK; ²Department of Public Health, Environments and Society, London School of Hygiene and Tropical Medicine, London, UK

10.1136/jech-2020-SSMabstracts.181

Background English local authorities (LAs) have certain powers they can utilise to help them tackle alcohol-related harms. One such power is the Late Night Levy (LNL) which uses a 'polluter pays' principle whereby the LA charges alcohol retailers a fee for selling alcohol after midnight. Public health evaluators are increasingly considering how concepts from complexity science and systems thinking might shed light on how interventions can affect health in complex, real world settings such as local alcohol systems. This study aims to explore how the LNL acts as an event within the alcohol system, with a particular emphasis on understanding how the intervention, actors and the system as a whole adapt and co-evolve in response.

Methods We conducted a process evaluation, using a complex systems perspective, in one LA. We included a range of system actors and utilised qualitative data generation methods. We interviewed LNL implementers and alcohol consumers

(n=21), conducted observations in pubs and during community-safety patrols (34.5 hours) and reviewed documents on the LNL and the broader system (n=15). We conducted a deductive thematic analysis, applying concepts from systems thinking and complexity science to: 1) describe the system into which the LNL is introduced; and 2) identify emergent system outcomes.

Results The LNL operates within a complex system consisting of diverse actors with competing goals; these include the local political system and statutory functions, the retail sector, the third sector, residents and users of the night-time economy. Early analyses identified a number of system responses stemming from the intervention, which include: 1) retailer adaptation to the levy; 2) new and evolving relationships between retailers, the police, community-safety officers and LA officers; 3) changes in venue management; and 4) the evolution of the intervention itself in response to broader system feedback. These adaptations are hypothesised to affect alcohol-related health and social harms. However, the intervention also interacts with wider system influences on public health, for example local government budgetary constraints during a period of central government cuts.

Discussion Conceptualising the LNL as an event within a complex system allows us to analyse its trajectory as it embeds within, and influences the broader system in which it is located. Further fieldwork and analyses are ongoing and focus on sampling additional system actors and exploring the influence of the vertical systems (e.g. national policy) which shape local responses to alcohol-related harms.

This study is funded by the National Institute for Health Research (NIHR) School for Public Health Research (Grant Reference Number PD-SPH-2015). The views expressed are those of the author(s) and not necessarily those of the NIHR or the Department of Health and Social Care.

P90

HEALTH-JUSTICE PARTNERSHIPS: A SYSTEMATIC SCOPING REVIEW OF INTERNATIONAL EVIDENCE ON IMPACT

¹S Beardon*, ²C Woodhead, ¹S Cooper, ³H Genn, ¹R Raine. ¹Department for Applied Health Research, University College London, London, UK; ²Department of Psychological Medicine, King's College London, London, UK; ³Faculty of Laws, University College London, London, UK

10.1136/jech-2020-SSMabstracts.182

Background Social welfare legal problems impact negatively on mental and physical health. These include issues with welfare benefits, debt, homelessness, family and employment. Services providing legal assistance have developed collaborations with healthcare across the UK and the globe. These 'health-justice partnerships' aim to support health of the most disadvantaged groups, tackling interconnected challenges of health inequalities and access to justice. International evidence on the impacts of these partnerships has never previously been reviewed. This study interrogates the state of current evidence and provides new insights from international practice.

Methods A systematic scoping review of international literature was undertaken. Both academic and grey literature was included from OECD countries, 1995–2018. The review

aimed to describe different service models and summarize evidence on impacts. Data from included publications were extracted and analysed narratively. Quality of included publications was assessed. Evidence on impact is presented according to each service objective.

Results Health-justice partnerships were diverse in their characteristics. Target populations focused on low income, vulnerable or underserved groups. Approaches to service coordination included co-location, referral, or integration of legal advisors into care teams and care pathways. The strongest evidence on impact was for increased access to legal assistance and improvements in individuals' financial and social circumstances. Despite this role in addressing social determinants of health, evidence was lacking on the wider issues of prevention and health inequalities. There was strong qualitative evidence for improvements in mental wellbeing, reflected in some quantitative research. Studies suggested positive outcomes in supporting healthcare professionals and contributing to high quality patient care, although findings on healthcare utilisation and costs were mixed.

Discussion This is the first international review on the delivery of health-justice partnerships, bringing together evidence from across the world to map current knowledge on service models and impacts. Integration of health and legal services aligns with policy priorities in both sectors, addressing social welfare legal issues which are leading causes of inequality and underlying determinants of poor health. The evidence supports health-justice partnerships as a means of improving access to justice for vulnerable groups and alleviating health-harming legal needs. Some key service objectives require a stronger knowledge base, while others remain un-evidenced. Cross-sector alignment and collaboration presents challenges, and further research is needed on successful delivery of integrated services. This review has informed a primary research study exploring implementation of health-justice partnerships in different settings across England to inform future policy and practice.

P91 IMPACT OF THE STATUTORY CONCESSIONARY TRAVEL SCHEME ON BUS TRAVEL AMONG OLDER PEOPLE: A NATURAL EXPERIMENT FROM ENGLAND

E Whitley*, P Craig, F Popham. *Social and Public Health Sciences Unit, University of Glasgow, Glasgow, UK*

10.1136/jech-2020-SSMabstracts.183

Background In the context of ageing populations worldwide, increasing numbers of older people are lonely, isolated and excluded, with serious implications for their health, cognitive and physical functioning. Access to good public transport can improve mobility, access to services and social participation among older adults. Policies that improve access and encourage the use of public transport are therefore potentially important in promoting healthy and successful ageing. Concessionary travel schemes for older people are in place in many countries but are under threat following the global financial crisis. Evidence regarding the success of these schemes in increasing activity and social participation is generally positive but based largely on qualitative or observational associations. In particular, robust evaluations of the schemes are limited by the lack of appropriate comparison groups as they generally represent a universal benefit.

Methods We use changes to the English statutory concessionary travel scheme over time, in particular the rising eligibility age from 2010 onwards, as a natural experiment to explore its impact on older people's travel. A difference-in-difference analysis of the annual National Travel Surveys (2002–2016) compares three age groups differentially affected by the eligibility criteria: (i) those aged 50–59 years who were never eligible for the scheme (N~2,500 per year); (ii) those aged 60–64 years who were decreasingly eligible for the scheme from 2010 onwards (N~1,000 per year); and (iii) those aged 65–74 years who were consistently eligible for the scheme throughout the period of interest (N~2,000 per year).

Results Compared with 50–59 year-olds, bus travel among 60–74 year-olds increased year-on-year from 2002 to 2010, particularly from 2006 when the scheme offered free travel. The frequency of bus travel in both the older groups then fell following rises in eligibility age (annual change in the proportion traveling by bus at least weekly: -2.9% (-4.1%, -1.7%) in 60–64 and 65–74 compared with 50–59 year-olds). Results were consistent across gender, occupation and urban/rurality.

Conclusion Our results indicate that access to, specifically, free travel increases bus use and access to services among older people, potentially improving mobility, social participation and health. However, the rising eligibility age in England has led to a reduction in bus travel in older people, including those not directly affected by the change, demonstrating that the positive impact of the concession goes beyond those who are eligible. Future work should explore the cost/benefit trade-off of this and similar schemes worldwide.

P92 A MATCHED ANALYSIS OF THE IMPACT HAVING A CARER HAS ON AN INDIVIDUAL'S HEALTH AND SOCIAL CARE UTILISATION ACROSS FIVE SETTINGS OF CARE FOR ADULT RESIDENTS OF BARKING AND DAGENHAM

¹J Shand*, ¹M Gomes, ²S Morris. ¹Department of Applied Health Research, UCL, London, UK; ²Department of Public Health and Primary Care, University of Cambridge, Cambridge, UK

10.1136/jech-2020-SSMabstracts.184

Background Across the UK today, an estimated 6.8 million people are carers, supporting friends and family who are older, disabled or seriously ill. The economic value of informal carers contributions is an estimated £132 billion a year. This is calculated by multiplying the total hours of care provided by carers (using responses from the 'Personal Social Services Survey of Adult Carers in England 2014–15' to determine numbers of carers and average hours of care provided) by the unit cost of an hour of replacement home care for an adult (£17.20). This assumes informal carers provide substitution for formal care. Reliance on carers appears to be increasing, with fewer people getting access to formal support as a result of local authority budget cuts.

This study aimed to understand if people with a carer have different levels of service use across different settings of care when compared to those who do not have a carer but have similar characteristics.

Methods A quantitative study using person-level data in Barking and Dagenham (B&D), a London borough, to assess the impact of having a carer in terms of the differences in cost-weighted utilisation relative to a matched control group. Nearest neighbour matching was used, matching on age,