behaviour (e.g. driving, active travel, public transit, walking, and cycling) from adults in the general population. Interventions were categorized into functions (how the intervention seeks to affect behaviour). Depending on whether gains or losses of functions could occur, interventions were classified as carrots (e.g. new bikeshare programs), combined carrot and stick (e.g. traffic calming), or stick interventions (e.g. congestion charging). Harvest plots were used to visually summarize the findings weighted by study quality. Where possible, outcomes were converted into standardized mean differences (SMD) and random-effects meta-analyses were conducted.

Results We extracted data from 83 publications reporting 98 interventions. From these, we identified 20 intervention types and eight function categories. The majority of interventions were carrots (n=64), followed by carrot and stick (n=17) and stick (n=17). Harvest plots demonstrated that most evaluations, particularly those classified as higher quality, found changes in favour of the intervention. Results for carrot interventions, however, were more less consistent than for stick or combined interventions. This was consistent with findings from the meta-analysis, which were statistically nonsignificant but had point-estimates of greater magnitude for driving outcomes for sticks (SMD -0.21; 95%CI -0.43, 0.01) and combined carrot and stick interventions (-0.17; -0.65, 0.31) compared to carrots (-0.09; -0.21, 0.03). Likewise, for active travel outcomes, combined carrot and stick interventions had a higher SMD (0.39; -0.01, 0.78) compared to carrot interventions (0.10; -0.06, 0.25). Financial functions were found to be the most effective for driving outcomes, whereas access, convenience, safety, and space were more effective for active travel outcomes.

Discussion This is the first review to compare whether positive, negative, or combined strategies and their functions differ in terms of effectiveness on travel behaviour, which can aid policymakers in designing sustainable transportation policies. Further research is needed for interventions with a stick component, which suggest greater effectiveness yet remain less well-studied, possibly because they are less conducive to experimental manipulation.

OP51

ACTIVE GROUP-BASED PERFORMING ARTS INTERVENTIONS FOR PARKINSON'S DISEASE: SYSTEMATIC REVIEW AND META-ANALYSIS

¹Maxwell Barnish*, ²Susannah Barran. ¹Peninsula Technology Assessment Group (PenTAG), University of Exeter, Exeter, UK; ²Children and Young People's Speech and Language Therapy, Guy's and St Thomas' NHS Foundation Trust, London, UK

10.1136/jech-2021-SSMabstracts.51

Background Parkinson's disease (PD) is a common age-related neurodegenerative condition associated with a wide range of motor and non-motor symptoms. There has been increasing interest in performing arts interventions for PD, due to the limitations of pharmacological therapies. As no prior systematic review had addressed a wide range of artistic modalities, we systematically evaluated the potential benefit of all active group-based performing arts interventions in PD.

Methods In order to systematically evaluate the benefit of performing arts interventions in PD, searches on PsycINFO, AMED, CINAHL, EMBASE, and MEDLINE were performed in February 2020. Supplementary web searches and citation chasing were conducted. Included studies used any quantitative design to assess the benefit of any active group-based

performing arts intervention on quality of life, functional communication, speech, motor function or cognitive status for people with diagnosed PD. Following standardised duplicate screening and data extraction, data synthesis was conducted using thematic narrative synthesis and random effects meta-analyses using Review Manager 5.3. Risk of bias was evaluated using SURE checklists (Cardiff University).

Results Searches identified 2669 records, of which 260 proceeded to initial full-text review following de-duplication, 129 to detailed full-text review and 67 to inclusion in the systematic review, representing 56 separate studies. Data from 10 separate studies were included in meta-analyses. Risk of bias assessment revealed limitations across many studies. Dance was the most studied artistic modality (39 studies), followed by singing (12 studies), music therapy (4 studies) and theatre (2 studies). Narrative synthesis showed evidence for a benefit of dance, music therapy and singing on quality of life; singing on speech; dance and music therapy on cognition; and dance, music therapy, singing and theatre on motor function. There was no evidence of benefit on functional communication. In meta-analyses, tango-based dance predicted superior UPDRS motor than usual care (Z=2.87, p=0.004) and superior timed-up-and-go than exercise (Z=3.77, p=0.0002), while PD-specific dance predicted superior PDQ-39 quality of life than usual care (Z=3.77, p=0.0002).

Conclusion Performing arts interventions may offer benefit in PD. However, it is difficult to assess which performing arts modalities may be most beneficial for which outcomes, due to a lack of studies directly comparing different performing arts interventions. Key limitations included small sample sizes, differences in comparator interventions, differences in intervention frequency and duration, as well as differing disciplinary backgrounds of session leaders. Further research is needed with greater methodological rigour before firm conclusions can be drawn.

OP52

IDENTIFYING LOCAL AUTHORITY NEED FOR, AND UPTAKE OF, SCHOOL-BASED PHYSICAL ACTIVITY INTERVENTIONS IN ENGLAND – A CLUSTER ANALYSIS USING ROUTINE DATA

¹Tishya Venkatraman*, ¹Kate Honeyford, ¹Bina Ram, ²Esther MF van Sluijs, ¹Céire E Costelloe, ¹Sonia Saxena. ¹Department of Primary Care and Public Health, Imperial College London, London, UK; ²MRC Epidemiology Unit and Centre for Diet and Activity Research (CEDAR), University of Cambridge, Cambridge, UK

10.1136/jech-2021-SSMabstracts.52

Background Schools in England need effective, simple interventions to address shortfalls in children's physical activity. Areas in England where physical activity is low among children may particularly benefit from school-based physical activity interventions. Further, stark geographical inequalities for children's physical activity exist and have widened during Covid-19 lockdowns. The Daily Mile (TDM) is a school-based running programme that has been widely adopted across England. However, there is limited evidence of its equitability and whether it reaches children in most need. We aimed to i) identify local authorities in England with the highest need for children's physical activity promotion and ii) determine whether TDM reaches schools in areas of high need.

Methods This was a cross-sectional study linking routinely collected data from a range of sources including the Active Lives Surveys and national datasets. Characteristics of need were

identified following a literature review to identify determinants of children's physical activity. Our assessment of local authority 'need' included measures of children and adult's physical activity and excess weight, children's mental health needs, access and utilisation of outdoor space for exercise, and deprivation. We grouped local authorities using a hierarchical cluster analysis. The optimal number of clusters for our data was three clusters. We estimated the association between 'need' and registration to TDM with a Poisson regression.

Results We identified three distinct clusters of need for children's physical activity interventions representing 4.5 million primary school children. High-need, medium- need, and low-need local authorities included 24%, 56% and 20% of 123 upper-tier English local authorities respectively. Schools in local authorities of high-need were more likely to be registered to TDM (IRR: 1.25, 95%CI:1.12–1.39) compared with schools in low-need local authorities, where registration to TDM ranged from one in five schools in the low-need cluster to one in four in the high-need cluster.

Conclusion Our study suggests that TDM is an equitable intervention reaching schools in areas with the highest need. Areas of highest need are concentrated in particular areas in England, for example the North East. Matching public health interventions to the needs of the population is more efficient and equitable than blanket policies directed at whole populations. This identification of clusters of local authorities that share patterns of need may guide shared learning between local authorities that share similar challenges and contextual features.

OP53

EXPLORING VIEWS ON POTENTIAL COMPONENTS OF A DIET AND PHYSICAL ACTIVITY INTERVENTION WITH PARENTAL INVOLVEMENT AMONG CHILDREN IN NIGERIA

Oritseweyinmi Orighoye*, Maria Maynard, Tanefa Apekey. *Clinical and Applied Sciences, Leeds Beckett University, Leeds, UK*

10.1136/jech-2021-SSMabstracts.53

Background There is limited evidence on diet and physical activity (PA) interventions to prevent childhood over- and under-nutrition in Nigeria, and none focused on parental involvement. Knowledge of socio-cultural and environmental contexts, prioritising views of target populations, is needed to inform intervention strategies. The aim of this study was to explore parents, children and other stakeholders' views on the factors that might enable or hinder participation in diet and PA interventions and parental involvement, and potential intervention components likely to be feasible and acceptable.

Methods A grounded theory, qualitative cross-sectional study was conducted in culturally diverse local government areas of Lagos State, Nigeria. Participants were identified through purposive and theoretical sampling, and data collected over three iterative phases. Eleven boys and girls aged 8–11 years; 19 women and 14 men aged 19–60 years who were parents, teachers/school heads, community leaders, health workers, and health or education civil servants, took part in 25 semi-structured one-to-one interviews and three focus groups. Discussions were digitally recorded and transcribed verbatim. Manual thematic analysis and independent coding of the transcripts generated key themes and reduced bias in the analysis.

Results Three overarching themes were identified: 1. Active community collaboration 2. Strategies for involving families; and 3. Schools as key settings for interventions. Adult participants voiced active partnership between communities and schools as essential to addressing barriers to diet and physical activity interventions, such as inconsistent funding and lack of safe outdoor space for PA. Children reported concerns about school meal quality and poor access to clean water and menstrual hygiene products impacting PA participation. Suggested solutions achieved by community partnerships included security for outside play areas, and infrastructure improvements using recycled/locally sourced materials, and health campaigns funded by local dignitaries. Suggested activities for engaging families in interventions included health literacy teaching for parents, using learning aids tailored to literacy levels and local dialects, and involving religious leaders.

Conclusion This study highlighted the challenges for consideration in childhood diet and PA intervention development in Nigeria. The feasibility and acceptability of grassroots suggestions for intervention components and parental involvement could usefully be explored in future pilot studies.

OP54

ASSOCIATIONS BETWEEN SOCIOECONOMIC POSITION ACROSS LIFE AND GRIP STRENGTH AT AGE 46 YEARS: FINDINGS FROM THE 1970 BRITISH COHORT STUDY

Mohamed Yusuf*, Jamie McPhee, Gallin Montgomery, Rachel Cooper. *Department of Sport and Exercise Sciences, Manchester Metropolitan University, Manchester, UK*

10.1136/jech-2021-SSMabstracts.54

Background Muscle weakness is a key criterion for important age-related conditions including sarcopenia and frailty. Research suggests lower childhood socioeconomic position (SEP) contributes to muscle weakness in later life but there is little evidence in younger adults closer to peak muscle strength from more recently born cohorts. We aimed to examine the relationships between indicators of SEP in childhood and adulthood and grip strength at age 46 years.

Methods A total of 3,113 men and 3,132 women from the 1970 British Cohort study, with data on paternal occupational class and parental education levels at age 5 and own occupational class, education level, grip strength and covariates including height, body mass index and occupational activity at age 46, were included in analyses. Interactions between sex and each SEP indicator were formally assessed, and models were sex-stratified if evidence of interaction was found. Linear regression models were used to test associations of childhood and adulthood SEP with maximum grip strength.

Results Among women, there was evidence of associations between lower SEP in childhood and adulthood and weaker grip strength. For example, women whose fathers were in the lowest occupational classes had 1.14kg (95%CI: -1.74,-0.54) weaker grip strength than women whose fathers were in the highest occupational classes, and these associations were not fully explained by covariates (fully-adjusted regression coefficient: -0.81kg (-1.39,-0.22)). Among men, different patterns of association were observed (*p*-values for sex interactions <0.05). In unadjusted models, lower SEP in both childhood and adulthood was associated with stronger grip, and these associations strengthened after adjustment for height. For example, men whose fathers were in the lowest occupational classes had stronger grip strength 1.01kg