receipt and care-provision by the older population as maintaining these activities could be challenging with travel restrictions and recommended physical distancing. This study aimed to determine if prevalence of informal caring changed during the COVID-19 pandemic, and examine longitudinal trends in quality of life, stress, and depressive symptoms in carers.

Methods We analysed the COVID-19 self-completion questionnaire (SCQ) sub-study (June–November 2020) of The Irish Longitudinal Study on Ageing (TILDA), a nationally representative study of people aged ≥ 50 years ongoing since 2009. A total of 3,043 participants aged ≥ 60 years were included. Participants were asked if they cared for someone during the COVID-19 pandemic, their relationship to the recipient, and number of hours/week. Outcome measures collected were CES-D8, Perceived stress Scale (PSS) and CASP12. Change in prevalence of caring from previous waves was examined. Multi-level regression analyses examined longitudinal trends in CES-D8, PSS and CASP12 scores by caring status since Wave 3 (2014), adjusting for sociodemographic variables. Analyses were conducted in Stata 14.

Results Prevalence of informal caring tripled during the COVID-19 pandemic (15% compared to 5% in 2018). While 26% of carers reported they had stopped caring since the pandemic, most of those providing care were new carers (68% of men and 66% of women carers). Becoming a new carer during the pandemic was associated with lower CASP12 scores ($\beta = -0.582$, 95% CI $-1.018, -0.146$) and higher PSS ($\beta = 0.541$, 95% CI $0.316, 0.765$) compared to no caring. Caring for ≥ 50 hours in the past week was associated with lower CASP12 ($\beta = -0.878$, 95% CI $-1.345, -0.412$), higher PSS scores ($\beta = 0.383$, 95% CI $0.066, 0.700$) and higher depressive symptoms ($\beta = 0.371$, 95% CI $0.251, 0.490$). This association was moderated by carer status; new carers with increased hours had higher CES-D8 compared to non-carers, with no difference for those who stopped or continued caring.

Discussion This study demonstrated that the reduction of availability of both formal and informal care for older people during the COVID-19 pandemic, led to increased caring by older household members and was associated with lower well-being and mental health. This study provides further evidence of the detrimental indirect effects of the pandemic for older people and emphasizes for policymakers the importance of prioritising formal and informal care arrangements for the older population.

OP62

CARING IN THE TIME OF COVID-19, LONGITUDINAL TRENDS IN WELLBEING AND MENTAL HEALTH IN CARERS IN IRELAND: EVIDENCE FROM THE IRISH LONGITUDINAL STUDY ON AGEING (TILDA)*

¹Christine McGarrigle*, ¹Mark Ward, ¹Aisling O'Halloran, ¹Celine DeLooze, ^{1,2}Rose Anne Kenny. ¹The Irish Longitudinal Study on Ageing (TILDA), Trinity College Dublin, Dublin, Ireland; ²Mercer's Institute for Successful Ageing, St James's hospital, Dublin, Ireland

10.1136/jech-2021-SSMabstracts.62

Background The initial public health response to the COVID-19 pandemic in Ireland recommended that older people aged ≥ 70 years remain at home, and physically isolate from those outside their household. This may have affected both care-

OP63

THE MENTAL HEALTH OF UK WORKERS DURING THE COVID-19 PANDEMIC: A LONGITUDINAL ANALYSIS OF DIFFERENTIAL IMPACTS ACROSS INDUSTRIES AND SOCIAL CLASS CATEGORIES*

¹Theocharis Kromydas*, ¹Michael Green, ¹Peter Graig, ¹Vittal Katikereddi, ¹Alastair Layland, ²Claire Niedzwiedz, ¹Anna Pearce, ¹Rachel Thomson, ¹Evangelia Demou. ¹MRC-CSO Social and Public Health Sciences Unit, Institute of Health and Wellbeing, University of Glasgow, Glasgow, UK; ²Institute of Health and Wellbeing, College of Medical, Veterinary and Life Sciences, University of Glasgow, Glasgow, UK

10.1136/jech-2021-SSMabstracts.63

Background The COVID-19 pandemic has substantially affected mental health but its impacts are likely to differ across industries and social class groups. The 'new normal' in employment opportunities, patterns and conditions is likely to include a degree of fear, uncertainty, employment instability and job loss. It is imperative we identify and understand how

workers have been disadvantaged by the pandemic and current working modes. We aimed to examine the impact of COVID-19 on the mental health of UK workers by industry and social class category and to investigate whether any observed impacts differed by age, gender, and country of residence within the UK.

Methods We used the UK Household Longitudinal Study (Understanding Society) to analyse data from adults in paid employment, who participated in pre-COVID-19 Waves [9 (2017/19) and 10/11(2018/20)] and in at least one COVID-19 wave. Our primary outcome was probable psychological distress, defined by 'caseness' (a score of 4+) in the General Health Questionnaire-12 (GHQ-12) and exposures included industrial sectors (classified using the UK Standard Industrial Classification) and social class (using the Eight Class National Statistics Socio-economic Classification). We fitted mixed-effects logistic regression models, adjusting for age, gender, employment type, financial situation and UK country of residence and used inverse probability weights to account for attrition and non-response, to estimate odds ratios (ORs) with 95% confidence intervals (95%CI). Subsequently, we stratified by gender, UK country of residence, and age group.

Results Our final sample includes 44,500 observations (Pre-pandemic period=13,314, pandemic period=31,186) across 7,075 individuals. GHQ-caseness increased for workers in most industries during the pandemic; especially in 'Accommodation and Food Services' (OR:3.2; 95%CI:1.9,5.4). Analysis by social class showed that 'Small employers and own account' workers were the most affected (OR:2.9; 95% CI:2.1,4.0). Our stratified analysis by industry demonstrated the largest increase in GHQ-caseness between the pre and pandemic periods to be by gender in the 'Accommodation and Food Services' sector (OR:9.6; 95%CI:3.2,29.0 for men; OR:2.0; 95%CI:1.2,3.4 for women); whereas by social class the largest increase was observed when stratifying by UK country of residence for 'Small employers and own account' workers (non-England: OR:8.8; 95%CI:4.1,18.8; England: OR:2.5; 95%CI:1.6,3.9).

Conclusion Our findings indicate that the pandemic has significantly affected the mental health of UK workers. Policies with a decentralised element are needed, prioritising support to certain industrial sectors and the self-employed/small business owners, as well as for particular demographic groups (e. g. men in the 'Accommodation and Food Services' industry) who are in an excessively high risk.

OP64

COMMUNICATION ACROSS MATERNAL SOCIAL NETWORKS DURING THE UK'S NATIONAL LOCKDOWN AND ITS ASSOCIATION WITH POSTNATAL DEPRESSIVE SYMPTOMS

¹Emily Emmott*, ^{1,2}Sarah Myers. ¹UCL Anthropology, UCL, London, UK; ²BirthRites Independent Max Planck Research Group, Max Planck Institute for Evolutionary Anthropology, Leipzig, Germany

10.1136/jech-2021-SSMabstracts.64

Background Postnatal depression (PND) is highly detrimental for both mother and baby, with a pre-COVID-19 estimated prevalence of up to 23% in Europe. Low social support is a key risk factor for developing PND. Social distancing measures designed to limit COVID-19 transmission likely created unprecedented barriers for mothers to access social support. We explore how social support networks existed during the

UK's first national lockdown amongst London mothers with infants, and their associations with postnatal depressive symptoms.

Methods Using self-reported social network data from 162 London-based mothers with infants aged ≤6 months collected in May-June 2020 (during the first lockdown), we conduct a concurrent design mixed-method study. We quantitatively describe communication within maternal support networks, using quasi-Poisson regression models to analyse associations with postnatal depressive symptoms using the Edinburgh Postnatal Depression Scale (EPDS). In parallel, we thematically analyse open-text data to explore and understand the experiences of mothers with infants during England's lockdown.

Results *Quantitative findings:* Levels of depressive symptoms were high in our sample, with 47.5% of participants meeting the ≥11EPDS cut-off for potential postnatal depression. On average, women kept in contact with 10–11 people remotely (via WhatsApp, Zoom etc), and saw one family member other than their partner, suggesting high compliance with lockdown rules. The number of seen in person, and remote communication with a higher proportion of those not seen, negatively predicted depressive symptoms (RRR = 0.950 – 0.996 across models). However, a higher proportion of relatives seen in person was associated with increased depressive symptoms (RRR=1.003; 95% CI 1.001–1.006), suggesting family members may have broken lockdown rules to support mums in need. *Qualitative findings:* We identified four themes. Some mothers in our sample felt lockdown 'protected' family time which led to better bonding, often facilitated by high levels of support from the partner (theme 1). However, many experienced the 'burden of constant mothering' without anyone around to help (theme 2). Mothers felt remote contact was 'not as good' and led to reduced or inadequate support (theme 3). Mothers grieved for lost opportunities for them and their baby, and worried about the developmental impact on their baby (theme 4). Overall, these experiences coexisted with feelings of isolation, exhaustion, worry, inadequacy, guilt, and increased stress.

Conclusion Results suggest the strict lockdown in the UK may have increased the risk of postnatal depression, reflected in the high levels of PND symptoms and poor maternal experiences in our sample.

OP65

MENTAL HEALTH INEQUALITIES IN DISRUPTIONS TO HEALTHCARE, ECONOMIC ACTIVITY AND HOUSING DURING COVID-19: FINDINGS FROM 12 UK LONGITUDINAL POPULATION SURVEYS

¹Michael J Green*, ²Giorgio Di Gessa, ³Jane Maddock, ⁴Eoin McElroy, ^{5,6}Ellen J Thomson, ^{3,7}Praveetha Patalay, ⁶Helena L Davies, ⁶Jessica Mundy, ⁸Anna J Stevenson, ^{9,10}Alex SF Kwong. ¹MRC/CSO Social and Public Health Sciences Unit, University of Glasgow, Glasgow, UK; ²Institute of Epidemiology and Health Care, UCL, London, UK; ³MRC Unit for Lifelong Health and Ageing, UCL, London, UK; ⁴Department of Neuroscience, Psychology and Behaviour, University of Leicester, Leicester, UK; ⁵Department of Twins Research and Genetic Epidemiology, Kings College London, London, UK; ⁶Social, Genetic and Developmental Psychiatry Centre, Kings College London, London, UK; ⁷Centre for Longitudinal Studies, UCL, London, UK; ⁸Centre for Genomic and Experimental Medicine, University of Edinburgh, Edinburgh, UK; ⁹Division of Psychiatry, University of Edinburgh, Edinburgh, UK; ¹⁰Population Health Sciences, University of Bristol, Bristol, UK

10.1136/jech-2021-SSMabstracts.65

Background The COVID-19 pandemic with its associated virus suppression measures have disrupted many domains of