

Supplementary File 1. Baseline comparison between the study sample and participants who dropped out at 3-month follow-up

Predictors	N = 3,438 Drop-out sample	N= 3,169 Study sample	Test of difference
	Raw counts (Weighted %)/ Weighted Mean (SD)	Raw counts (Weighted %)/ Weighted Mean (SD)	<i>t, <math>\chi</math></i>
Age	34.95 (10.79)	41.17 (13.66)	<i>t</i> (9998)=41.08***
Gender			
Male	1748 (50.2)	1224 (37.0)	$\chi$ (1)=174.01***
Not male	1690 (49.8)	1945 (63.0)	
Race/Ethnicity			
Non-Hispanic White	2050 (54.1)	2277 (73.7)	$\chi$ (4)=619.18***
Non-Hispanic Black	324 (8.1)	297 (7.1)	
Hispanic	673 (27.7)	276 (9.5)	
Asian	247 (4.2)	238 (3.3)	
Other	144 (5.9)	81 (6.4)	
Education			
Some college or below	906 (27.5)	947 (32.8)	$\chi$ (2)=62.40***
Associate/bachelor's degree	1712 (48.2)	1643 (48.8)	
Advanced degree	820 (24.3)	579 (18.4)	
Student status			
Not a student	2308 (68.7)	2702 (91.2)	$\chi$ (2)=838.20***
Part-time	346 (9.1)	175 (3.6)	
Full-time	784 (22.1)	292 (5.2)	
Marital Status			
Single	1117 (26.7)	1048 (23.5)	$\chi$ (2)=648.86***
D/S/W	214 (8.7)	410 (29.4)	
Married/LWP	2107 (64.5)	1711 (47.1)	
Minors in household			
No	1584 (45.9)	1994 (74.4)	$\chi$ (1)=842.68***
Yes	1854 (54.1)	1175 (25.6)	
Work Status			
Half/full-time	2607 (74.2)	2018 (47.4)	$\chi$ (2)=725.55***
Self-employed	230 (6.6)	316 (12.3)	
Not working	601 (19.2)	835 (40.2)	
Personal income	33417.52 (23560.66)	35188.18 (22288.63)	<i>t</i> (9998)=5.65**
State of residence			
Northeast	630 (16.8)	607 (16.7)	$\chi$ (3)=48.73***
Midwest	625 (18.8)	655 (21.9)	
South	1299 (37.4)	1251 (40.1)	
West	884 (26.9)	656 (21.2)	

Veteran status	618 (20.5)	198 (11.4)	$\chi(1)=156.29^{***}$
Medical Outcomes Study Social Support Survey	21.11 (5.66)	21.27 (6.67)	$t(9998) = 1.28^{***}$
Financial distress score	0.87 (1.33)	0.37 (0.93)	$t(9998) = -29.88^{***}$
COVID-19 status W1			
Untested	2040 (58.2)	2394 (79.1)	$\chi(2)=654.53^{***}$
Positive	305 (8.8)	49 (1.1)	
Negative	1093 (33.0)	726 (19.8)	
# of medical conditions	1.62 (2.58)	1.48 (1.82)	$t(9998) = 4.60^*$
History of psychiatric disorders			
SSD	213 (6.7)	47 (1.0)	$\chi(1)=239.38^{***}$
PTSD	415 (12.2)	269 (6.4)	$\chi(1)=101.04^{***}$
Bipolar disorder	389 (12.3)	159 (3.7)	$\chi(1)=268.32^{***}$
Anxiety disorder	1076 (29.6)	966 (22.4)	$\chi(1)=67.20^{***}$
Major depression	630 (18.0)	483 (11.6)	$\chi(1)=80.78^{***}$
AUD	625 (19.8)	162 (3.9)	$\chi(1)=647.00^{***}$
Drug use disorder	238 (7.1)	89 (1.9)	$\chi(1)=165.42^{***}$
TBI	91 (2.8)	42 (0.8)	$\chi(1)=64.44^{***}$
Positive screen for COVID-19 era-related stress	1825 (52.5)	789 (17.1)	$\chi(1)=1407.41^{***}$
Positive screen for major depression	1718 (49.1)	927 (22.2)	$\chi(1)=786.69^{***}$
Positive screen for generalized anxiety disorder	1743 (50.1)	945 (21.5)	$\chi(1)=894.58^{***}$
Past 2-week suicidal ideation	1487 (42.6)	577 (11.5)	$\chi(1)=1272.23^{***}$
Positive screen for AUD	1587 (45.9)	1019 (27.6)	$\chi(1)=356.85^{***}$
Any illicit drug use in past month	1023 (30.7)	452 (14.8)	$\chi(1)=364.31^{***}$

Note: D/S/W= Divorced/Separated/Widowed; SSD= Schizophrenia-spectrum disorder, PTSD= Posttraumatic stress disorder, AUD= Alcohol use disorder, TBI= Traumatic brain injury; \* $p<.05$ , \*\* $p<.01$ , \*\*\* $p<.001$