

## Closing the health inequity gap during the pandemic: COVID-19 mortality among racial and ethnic groups in Connecticut, March 2020 to December 2021

COVID-19 has disproportionately burdened racial and ethnic minority groups within the USA.<sup>1-5</sup> However, prior studies relied predominately on aggregated data from populations residing in congregate and non-congregate settings.<sup>1-5</sup> This approach may underestimate outcome disparities in the community, especially in states that experienced a large COVID-19 burden in nursing homes.<sup>5</sup> Herein, we identified COVID-19-related deaths among Connecticut residents of congregate and non-congregate settings and characterised the evolution of racial and ethnic disparities in COVID-19-related mortality among residents of non-congregate settings.

We retrospectively identified COVID-19-related deaths reported between 1 March 2020 and 13 December 2021 from the Connecticut Electronic Disease Surveillance System and determined residential status according to recorded address (online supplemental file).<sup>5</sup> Using Poisson regressions and 2019 census population estimates, we estimated age-adjusted mortality for composite race and ethnicity categories (Hispanic, non-Hispanic (NH)

black and NH white) and age-adjusted mortality rate ratios (MRR) for Hispanic and NH blacks, relative to NH whites, for three periods (1 March 2020 to 25 August 2020, 26 August 2020 to 12 July 2021 and 13 July 2021 to 13 December 2021; figure 1C; online supplemental file).<sup>5</sup>

During the study period, congregate facility-associated deaths accounted for 74.6% (3205/4299), 37.7% (1341/3557) and 14.2% (84/590) of deaths in the first, second and third periods, respectively (figure 1A). Among congregate facility-associated deaths, 5.6% (259), 10.0% (463) and 84.4% (3908) were Hispanic, NH black and NH white, respectively.

Overall, age-adjusted mortality among residents of non-congregate settings was highest for Hispanics (19.6/100 000 population-months), followed by NH blacks (17.3/100 000 population-months) and NH whites (6.4/100 000 population-months). During the initial pandemic period, we observed large disparities in COVID-19 mortality within residents of non-congregate settings (Hispanic MRR: 3.9 (95% CI, 3.3 to 4.5); NH black MRR: 5.2 (CI: 4.5 to 5.9)). However, by the third wave, the disparities had attenuated and mortality between NH blacks and NH whites no longer differed significantly (Hispanic MRR: 1.9 (1.5 to 2.5); NH black MRR: 1.3 (0.9 to 1.7); figure 1B,C/online supplemental eTable).

Our findings suggest that attenuation of racial and ethnic disparities is an

achievable public health goal. Future analyses investigating the role public health interventions had on reducing these disparities should be conducted to inform effective, generalisable policies targeting the outstanding inequities in COVID-19-related outcomes.

The limitations of our analysis include the use of a COVID-19-related death endpoint and the potential for misclassification of race, ethnicity and residency status. However, it was strengthened by the disaggregation of congregate facility events, through which we may have revealed levels of disparity not previously appreciated.<sup>1 2 4 5</sup>

Olivia Schultes,<sup>1</sup> Margaret L Lind,<sup>2</sup> Jessica Brockmeyer,<sup>3</sup> Peri Sosensky,<sup>2</sup> Derek A T Cummings,<sup>4,5,6</sup> Albert I Ko<sup>2,7,8</sup>

<sup>1</sup>Department of Epidemiology, School of Public Health, University of Washington, Seattle, Washington, USA

<sup>2</sup>Department of Epidemiology of Microbial Diseases, Yale University School of Public Health, New Haven, Connecticut, USA

<sup>3</sup>Connecticut Department of Public Health, Hartford, Connecticut, USA


<sup>4</sup>Department of Epidemiology, Center for Teaching and Learning, Johns Hopkins Bloomberg School of Public Health, Baltimore, Maryland, USA

<sup>5</sup>Department of Biology, University of Florida, Gainesville, Florida, USA

<sup>6</sup>Emerging Pathogens Institute, University of Florida, Gainesville, Florida, USA

<sup>7</sup>Instituto Gonçalo Moniz, Salvador, Bahia, Brazil

<sup>8</sup>Department of Internal Medicine, Section of Infectious Diseases, Yale School of Medicine, New Haven, Connecticut, USA



**Figure 1** Temporal trends in COVID-19-related deaths and mortality among the Connecticut population residing in congregate and non-congregate settings, according to composite race and ethnic groups, March 2020 to December 2021. (A) Monthly COVID-19 deaths by race/ethnicity and residency status. (B) Age-adjusted monthly COVID-19 mortality (deaths per 100 000 population-months) by composite race and ethnic groups in the population residing in non-congregate settings. (C) Ratios comparing mortality among non-Hispanic black (NH black) and Hispanic to non-Hispanic white (NH white) populations residing in non-congregate settings during each of the three epidemic periods. Congregate and non-congregate residency status was defined based on recorded residential address at the SARS-CoV-2 infection was reported.

**Correspondence to** Dr Margaret L Lind, School of Public Health, Yale University, New Haven, CT 06520, USA; margaret.lind@yale.edu

**Acknowledgements** The authors thank Charles Powell for accessing the data and Matthew Cartter, Lynn Sosa and Karyn Backus for helpful comments on the manuscript.

**Contributors** OS and MLL had full access to all the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis. Concept and design: AIK, OS and DATC. Acquisition, analysis or interpretation of data: OS, MLL and JB. Drafting of the manuscript: MLL, OS, AIK and PS. Critical revision of the manuscript for important intellectual content: all authors. Statistical analysis: MLL. Administrative, technical or material support: JB. Supervision: AIK and DATC.

**Funding** Connecticut Department of Public Health Emerging Infections Program: COVID-19 contract (DPH log # 2021-0071-3).

**Competing interests** None declared.

**Patient consent for publication** Not applicable.

**Ethics approval** This study involves human participants and was not deemed as human research by the Yale University Institutional Review Board (IRB). All data were retrospective and per the requirements of the overseeing IRB, no retroactive consent was needed.

**Provenance and peer review** Not commissioned; internally peer reviewed.

**Supplemental material** This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are

not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.



## OPEN ACCESS

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/jech-2022-218975>).

OS and MLL contributed equally.



**To cite** Schultes O, Lind ML, Brockmeyer J, *et al.* *J Epidemiol Community Health* 2022;**76**:695–696.

Received 1 March 2022

Accepted 5 March 2022

Published Online First 15 April 2022

*J Epidemiol Community Health* 2022;**76**:695–696.  
doi:10.1136/jech-2022-218975

### ORCID iD

Margaret L Lind <http://orcid.org/0000-0002-1572-4074>

### REFERENCES

- 1 Laurencin CT, Wu ZH, Grady JJ, *et al.* Changes in COVID-19-Associated deaths during a year among blacks and Hispanics compared to whites in the state of Connecticut. *J Racial Ethn Health Disparities* 2021. doi:10.1007/s40615-021-01143-z. [Epub ahead of print: 28 Sep 2021].
- 2 Mude W, Oguoma VM, Nyanhanda T, *et al.* Racial disparities in COVID-19 pandemic cases, hospitalisations, and deaths: a systematic review and meta-analysis. *J Glob Health* 2021;11:05015.
- 3 Lawton R, Zheng K, Zheng D, *et al.* A longitudinal study of convergence between black and white COVID-19 mortality: a County fixed effects approach. *Lancet Reg Health Am* 2021;1:100011.
- 4 Mackey K, Ayers CK, Kondo KK, *et al.* Racial and Ethnic Disparities in COVID-19-Related Infections, Hospitalizations, and Deaths : A Systematic Review. *Ann Intern Med* 2021;174:362–73.
- 5 Hayes L, Abdellatif E, Olson J. *Connecticut COVID-19 mortality: demographic comparisons and disparities, March 2020–February 2021*. Hartford, CT: Connecticut Department of Public Health, 2021.