


Stigma: the social virus spreading faster than COVID-19

Kim Robin van Daalen ¹, Mark Cobain,² Oscar H Franco,³ Rajiv Chowdhury⁴

Novel epidemics and pandemics are inherently plagued by scientific uncertainties and a rapidly evolving nature. This can lead to widespread fear and confusion—directly proportional to the level of disease impact and subsequent media coverage, which perpetuates panic.¹ Evolutionary psychologists argue that as response to fear humans tend to distance from those considered a source of danger (infections), guided by self-preservation, desire to find control or unfounded beliefs that those infected bear responsibility. Yet, this leads to misplaced reactions and stigmatisation.²

Originally from ancient Greek, the term ‘stigma’ was used to ‘permanently mark people as criminals, traitors or slaves’. Nowadays, it describes negative associations and discrimination against people with certain attributes (social, physical, behavioural).³ Examples of stigmatisation can be found throughout infectious disease history, affecting communities, caregivers, family and those infected. Among the most stigmatised have been those affected by or associated with HIV/AIDS.⁴ Many other examples exist, towards Jewish immigrants during the 1892 typhus fever and cholera outbreaks in New York City,¹ towards ‘African-ness’ during Ebola outbreak(s), towards ‘Asian-ness’ during SARS⁴ and towards those of Mexican/Latin American descent during 2009 H1N1 influenza pandemic (inadequately called ‘Mexican flu’). These stigmas significantly limited the ability to control diseases and affected mental wellbeing.

This is not different for COVID-19. The pandemic provoked worldwide discriminatory behaviours—including violence—and social stigma towards those (perceived) to have or been in contact with the virus or those of specific ethnic backgrounds

(especially those of Asian descent and immigrants). This includes deplorable examples such as the use of ‘Chinese virus’ by government representatives or use of pejorative expressions towards healthcare workers.⁵

Such stigmatisation can increase transmission, have critical hidden human costs on those stigmatised and exacerbate already difficult circumstances of the most vulnerable populations (especially in low- and middle-income countries).^{4–8} First, people with or at risk of disease may be more reluctant to seek healthcare when needed or may try to hide the disease or misreport symptoms, reducing early detection and treatment.⁴ Second, certain health behaviours are required to prevent disease, while stigma may hinder adherence due to avoidant coping strategies or mistrust in public health agencies by those stigmatised.⁶ Third, experiencing and anticipating stigma can lead to stress, elevated depressive symptoms and suicidal ideation.⁹ Similarly, internalised stigma may result in individuals seeing themselves as inferior or not worthy.⁶ Fourth, social stigma can often lead to neglect, catastrophic costs and poverty, increasing susceptibility to disease.⁷ Fifth, stigma may lead towards distorted perceptions of risk and disproportionate allocation of health resources.⁷ Worryingly, social stigmatisation often persists even after outbreaks, epidemics or pandemics have

ended (eg, continuous social rejection of Ebola survivors).⁸ Finally, stigma from multiple sources (eg, infectious disease, mental health, disability) can co-occur and interact with each other and other dimensions (eg, ethnicity, employment), increasing impact and inequalities.¹⁰

Furthermore, like previously described by Logie and Turan, tensions exist between COVID-19 mitigation, containment and the prevention of stigma. For example, while physical distancing is essential to prevent COVID-19 spread, othering can result in mistreatment and social avoidance of stigmatised individuals. Similarly, travel restrictions may prevent COVID-19 spread, while potentiating xenophobia or stigma by generating ideas of ‘foreign invasion’. Consequently, it can be difficult for governments to balance between appropriate mitigation and preventing fear and stigmatisation.¹¹

Social stigma is not inevitable. Various strategies reducing disease-related stigma during pandemics exist—with vital roles for government, media, private sector and citizens (Figure 1).⁵ These strategies should be used to tackle sources of stigma and to assist those stigmatised. First, timely, honest risk communication, addressing misinformation and improving awareness, is deemed essential for all involved. Stigma reducing messaging can foster empathy while appropriately reflecting evolving COVID-19 patterns and normalising physical distancing.^{6, 11} Communication should be framed both for the general population as for the affected population in appropriate channels and language.¹ Second, strategies improving employment sick leave and access to testing have potential to address underlying social inequalities




Figure 1 Social stigma during the COVID-19 pandemic: what can individuals do? (Based on World Health Organization (WHO), International Federation of Red Cross and Red Crescent Societies (IRFC) and United Nations Children's Fund (UNICEF) guidance.)⁵

¹Cardiovascular Epidemiology Unit, Department of Public Health and Primary Care, Cambridge University, Cambridge, UK

²Department of Cardiovascular Medicine, Imperial College London, London, UK

³Institute of Social and Preventive Medicine, University of Bern, Bern, Berne, Switzerland

⁴Cardiovascular Epidemiology Unit, Department of Public Health and Primary Care, Cambridge University, Cambridge, UK

Correspondence to Ms Kim Robin van Daalen, Department of Public Health and Primary Care, Cambridge University, Cambridge, UK; krv22@cam.ac.uk

aiding stigmatisation.¹¹ Third, involving community leaders, their communities⁶ and those most affected by COVID-19¹¹ provides crucial information on contextually specific approaches. Furthermore, messages designed to foster a sense of altruism in behaviours¹² should feature diverse audiences, showing their contribution. Fourth, when public outreach and information campaigns are not enough, implementation of skill training and educational programmes may be beneficial.⁶ Nonetheless, further research is needed to clarify COVID-19-specific drivers of stigma at individual/structural levels and its impact on health.

Efforts should be focused on providing an adequate science-based COVID-19 response that includes the prevention of social stigma derived from COVID-19.

Twitter Kim Robin van Daalen @daalenkim

Contributors KRvD wrote the manuscript and designed the figure with support from RC, OHF and MC. All authors provided critical feedback and helped shape the presented commentary.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests The authors have no competing interest.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; internally peer reviewed.

This article is made freely available for use in accordance with BMJ's website terms and conditions for the duration of the covid-19 pandemic or until otherwise determined by BMJ. You may use, download and print the article for any lawful, non-commercial purpose (including text and data mining) provided that all copyright notices and trade marks are retained.

© Author(s) (or their employer(s)) 2021. No commercial re-use. See rights and permissions. Published by BMJ.



To cite van Daalen KR, Cobain M, Franco OH, *et al.* *J Epidemiol Community Health* Epub ahead of print: [please include Day Month Year]. doi:10.1136/jech-2020-214436

Received 29 April 2020

Revised 21 November 2020

Accepted 11 December 2020

J Epidemiol Community Health 2021;**0**:1–2.
doi:10.1136/jech-2020-214436

ORCID iD

Kim Robin van Daalen <http://orcid.org/0000-0001-6955-9708>

REFERENCES

- 1 Person B, Sy F, Holton K, *et al.* Fear and stigma: the epidemic within the SARS outbreak. *Emerg Infect Dis* 2004;10:358–63.

- 2 Smith RA, Hughes D. Infectious disease stigmas: maladaptive in modern society. *Commun Stud* 2014;65:132–8.
- 3 Goffman E. *Stigma: notes on the management of spoiled identity*. Englewood Cliffs, NJ:Prentice Hall, 1963.
- 4 Des Jarlais DC, Galea S, Tracy M, *et al.* Stigmatization of newly emerging infectious diseases: AIDS and SARS. *Am J Public Health* 2006;96:561–7.
- 5 Social stigma associated with COVID-19. IRFC, who, UNICEF 2020.
- 6 Fischer LS, Mansergh G, Lynch J, *et al.* Addressing disease-related stigma during infectious disease outbreaks. *Disaster Med Public Health Prep* 2019;13:989–94.
- 7 Barrett R, Brown PJ. Stigma in the time of influenza: social and institutional responses to pandemic emergencies. *J Infect Dis* 2008;197 Suppl 1:S34–7.
- 8 Overholt L, Wohl DA, Fischer WA, *et al.* Stigma and Ebola survivorship in Liberia: results from a longitudinal cohort study. *PLoS One* 2018;13:e0206595.
- 9 Zeng C, Li L, Hong YA, *et al.* A structural equation model of perceived and internalized stigma, depression, and suicidal status among people living with HIV/AIDS. *BMC Public Health* 2018;18:138.
- 10 Jackson-Best F, Edwards N. Stigma and intersectionality: a systematic review of systematic reviews across HIV/AIDS, mental illness, and physical disability. *BMC Public Health* 2018;18:1–19.
- 11 Logie CH, Turan JM. How do we balance tensions between COVID-19 public health responses and stigma mitigation? learning from HIV research. *AIDS Behav* 2020;24:2003–6.
- 12 Brooks SK, Webster RK, Smith LE, *et al.* The psychological impact of quarantine and how to reduce it: rapid review of the evidence. *The Lancet* 2020;395:912–20.