eTable 1. Linear regression detailing change in MVPA in restricted complete cases sample (n=3 209)

	Adjusted for sex and wear-		Adjusted for sex, wear-		Mutually-adjusted model c	
	time ^a		time, SEP 1	b	withthany-aujusted	mouer
PARENTAL OCCUPATIONAL CLASS						
I Professional, II Managerial/Technical	Ref				Ref	
III Manual or Non-manual	-3.17 (-5.34, -1.00)	< 0.005			-2.13 (-4.34, 0.07)	0.10
IV Partly skilled, V unskilled	-4.36 (-7.19, -1.53)				-2.86 (-5.78, 0.05)	
PRE-NATAL MATERNAL RISK FACT	ORS					
Maternal age at birth, per 1 year	0.34 (0.17, 0.51)	< 0.001	0.31 (0.14, 0.48)	< 0.001	0.28 (0.08, 0.48)	< 0.01
Previous pregnancies						
0	Ref		Ref		Ref	
1-2	1.14 (-0.77, 3.05)	0.32	1.14 (-0.77, 3.04)	0.25	0.51 (-1.48, 2.51)	0.25
3+	1.88 (-0.86, 4.63)		2.17 (-0.57, 4.91)		0.55 (-2.59, 3.69)	
Smoking during pregnancy	-3.15 (-4.92, -1.37)	< 0.001	-2.84 (-463, -1.05)	< 0.005	-2.92 (-4.76, -1.09)	< 0.005
Clinical diabetes	-5.35 (-15.5, 4.83)	0.30	-5.32 (-15.49, 4.84)	0.31	-5.71 (-15.84, 4.42)	0.27
POST-NATAL INFANT RISK FACTOR	S					
Gestational age						
BOYS						
Pre-term (<37 weeks)	-2.23 (-8.73, 4.26)		-2.04 (-8.53, 4.45)		-3.49 (-10.27, 3.29)	
Term (37-41 weeks)	Ref	< 0.005	Ref	< 0.005	Ref	< 0.005
Post-term (>41 weeks)	-8.30 (-12.83, -3.77)		-8.00 (-12.53, -3.47)		-7.28 (-11.81, -2.76)	
GIRLS						
Pre-term (<37 weeks)	-3.81 (-10.30, 2.67)		-2.91 (-16.15, 10.33)		-4.82 (-11.53, 1.88)	
Term (37-41 weeks)	Ref	0.35	Ref	0.42	Ref	0.65
Post-term (>41 weeks)	-1.71 (-5.59, 2.17)		-1.59 (-11.40, 8.21)		-1.00 (-4.88, 2.88)	
Birth weight, per 1kg	-1.25 (-2.96, 0.46) ^d	0.15	-1.38 (-3.09, 0.33) ^e	0.11	-2.61 (-4.47, -0.74)	< 0.01
Breastfeeding >3months, n(%)	2.91 (0.39, 5.43)	0.02	2.36 (-0.19, 4.90)	0.07	1.62 (-0.93, 4.18)	0.21
Infant health concerns, n(%)						
0	Ref		Ref		Ref	
1	0.08 (-2.12, 2.27)	0.61	0.13 (-2.07, 2.32)	0.67	-0.27 (-2.49, 1.95)	0.57
2+	-2.27 (-6.83, 2.29)		-2.03 (-6.59, 2.52)		-2.53 (-7.23, 2.17)	

^b 8 individual models, each adjusted for sex, wear time and paternal occupational class

^c A single combined model including sex, wear time, paternal occupational class, maternal age at birth, previous pregnancies, maternal smoking during pregnancy, maternal clinical diabetes, gestational age, standardised birth weight, breastfeeding and infant heath concerns

^d Estimate for birth weight in model adjusted for sex and gestational age: -1.57 (-3.39, 0.24); p=0.09

^e Estimate for birth weight in model adjusted for sex, wear time, SEP and gestational age: -1.69 (-3.50, 0.12); p=0.07

eTable 2. Comparison of pre- and post-natal factors between the analytical sample (n=5 011), those who participated in age 46 data collection but did not have MVPA data (n=3 570) and those who were lost to follow-up before age 46 (n=9 425)

,		Participated in age	
	Analytical	46 data collection	Lost to follow-up
	sample	but no MVPA	before age 46
	-	data	
	(n=5 011)		(n=9 425)
CITIES (CC)		(n=3 570)	
SEX, n(%)	2207 (47.0)	15(2(40.4)	5005 (55.1)
Male	2396 (47.8)	1762 (49.4)	5087 (55.1)
Female	2615 (52.2)	1808 (50.6)	4147 (44.9)
PARENTAL OCCUPATIONAL			
CLASS, n(%)			
I Professional, II Managerial/Technical	999 (21.6)	683 (20.7)	1428 (15.6)
III Manual or Non-manual	2730 (59.1)	1966 (59.7)	5245 (57.3)
IV Partly skilled, V unskilled	891 (19.3)	645 (19.6)	2474 (27.1)
PRE-NATAL MATERNAL RISK			
FACTORS			
Age at birth, mean ±SD	26.0 ± 5.3	25.9 ± 5.4	26.0 ± 5.7
Previous pregnancies, n(%)			
0	1707 (36.8)	1138 (34.4)	2975 (32.2)
1-2	2264 (48.7)	1609 (48.7)	4283 (46.3)
3+	674 (14.5)	559 (16.9)	1987 (21.5)
Smoking during pregnancies, n(%)	1986 (43.0)	1443 (43.9)	4470 (48.6)
Clinical diabetes, n (%)	39 (0.9)	26 (0.8)	65 (0.7)
POST-NATAL INFANT RISK			
FACTORS			
Gestational age: males, n(%)			
	73 (4.0)	70 (5.2)	315 (8.1)
	, ,	, ,	, ,
	79 (3.9)	51 (3.7)	225 (7.1)
	(- 10 (- 110)	(***)
	3382 + 534	3361 +522	3282 +0.65
	333 (13.3)	517 (12.2)	310 (0.5)
, , ,	3468 (75.7)	2432 (74.6)	6555 (74.1)
		` ,	
Smoking during pregnancies, n(%) Clinical diabetes, n (%) POST-NATAL INFANT RISK	1986 (43.0)	1443 (43.9)	4470 (48.6)

^a N varies due to missing data in early life factors